

[Images in...](#)

Cardiac lymphoma: ECG images pre-treatment and post-treatment

Daniel Xavier Augustine,¹ Anu Garg,¹ Christopher Knechtl²

¹ Department of Cardiology, RUH Bath NHS Foundation Trust, Bath, UK

² Department of Haematology, RUH Bath NHS Foundation Trust, Bath, UK

Correspondence to Daniel Xavier Augustine, dxaugustine@hotmail.com

DESCRIPTION

The chest radiograph demonstrating cardiomegaly and severe widening of the mediastinum (figure 1) is from a 55-year-old lady who presented with several months of breathlessness. Her ECG revealed Mobitz type II second-degree heart block.

A 2-year history of pruritus was noted. CT scan revealed mediastinal lymphadenopathy.

Transthoracic ECG showed thickening of the atrial septum consistent with an infiltrative process. It extended through the tricuspid valve and into the right ventricular

outflow tract (figure 2, upper panel). The infiltration was also seen to encase the aorta (figure 2, upper panel).

Biopsy of the mediastinal nodes yielded a histological diagnosis of nodular sclerosing Hodgkin's disease. Due to concerns relating to the possibility of cardiac rupture, treatment was initiated in the form of prednisolone 50 mg/m² daily for 11 days prior to adriamycin, bleomycin, vinblastine and dacarbazine (ABVD) chemotherapy.

Six months later, following eight cycles of ABVD chemotherapy, repeat ECG has shown remarkable regression of the Hodgkin's disease (fig 2, lower panel).




Figure 1 Chest x-ray on admission.




Figure 2 ECG pre-chemotherapy (upper panel) and post-chemotherapy (lower panel).

Learning points

- ▶ Autopsy series show that cardiac invasion by non-Hodgkin's lymphomas is as high as 25%,¹ whereas cardiac involvement by Hodgkin's disease is distinctly rare.²
- ▶ This case is more unusual in that the cardiac involvement by Hodgkin's disease has caused heart block.
- ▶ Pruritus is a feature in >25% of patients presenting with Hodgkin's disease.³ Earlier consideration of this may have avoided the ensuing cardiac complications.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Roberts WC**, Glancy DL, DeVita VT Jr. Heart in malignant lymphoma (Hodgkin's disease, lymphosarcoma, reticulum cell sarcoma and mycosis fungoides). A study of 196 autopsy cases. *Am J Cardiol* 1968;**22**:85–107.
2. **Filly R**, Bland N, Castellino RA. Radiographic distribution of intrathoracic disease in previously untreated patients with Hodgkin's disease and non-Hodgkin's lymphoma. *Radiology* 1976;**120**:277–81.
3. **Cavalli F**. Rare syndromes in Hodgkin's disease. *Ann Oncol* 1998;**9**(Suppl 5):S109–13.

This pdf has been created automatically from the final edited text and images.

Copyright 2010 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Augustine DX, Garg A, Knechtli C. Cardiac lymphoma: ECG images pre-treatment and post-treatment. *BMJ Case Reports* 2010;10.1136/bcr.05.2010.3018, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow