

Images in...

# Electrocardiographic changes in an athlete before and after detraining

Saqib Ghani, Sanjay Sharma

Cardiovascular Sciences Department, St George's University London, London, UK

Correspondence to Dr Saqib Ghani, saqibghani@hotmail.com


## DESCRIPTION

A young Caucasian female professional athlete underwent a 12-lead ECG as a part of preparticipation cardiovascular screening. She trained and exercised for an average 20 h per week. There were no symptoms

or relevant family history reported, and cardiovascular examination was normal. The ECG exhibited T-wave inversion in the inferior and lateral leads (figure 1). These ECG findings were considered abnormal, and warranted further investigations based on recent guidelines from



**Figure 1** ECG with T-wave inversions in inferior and lateral leads (arrows).



**Figure 2** Repeat ECG after a period of detraining shows resolution of T-wave inversions.

the European Society of Cardiology.<sup>1</sup> The echocardiogram showed normal left ventricular cavity size of 42 mm in diastole, maximal wall thickness of 10 mm and preserved bi-ventricular function. Further investigations including exercise stress test, 24-h ECG and cardiac MR scan with gadolinium enhancement were also normal, and did not suggest any phenotypic features of an underlying cardiomyopathy. A repeat ECG after a 6-week period of detraining (figure 2) was normal with resolution of T-wave inversions. The athlete continues to compete at professional level and is being followed-up with annual ECG and echocardiogram. Regular physical exercise is associated with electrical, structural and functional cardiac adaptation,<sup>2</sup> referred to as 'athletes heart'. However, T-wave inversions in inferior and lateral leads are uncommon even in highly trained athletes, and may be feature of underlying cardiomyopathy. In circumstances where further investigations do not reveal any significant abnormality, a period of detraining can be offered to resolve the dilemma due to ECG abnormalities. This case highlights the role of detraining in differentiating pathology from physiological cardiac adaptation to exercise.<sup>3</sup>

## Learning points

- ▶ T-wave inversions in inferior and lateral leads are uncommon in athletes, and warrant further investigations.
- ▶ Re-evaluation after a period of de-training can demonstrate resolution of ECG changes, and can be a useful tool in differentiating pathology from physiology.
- ▶ In absence of features of cardiomyopathy on extensive evaluation athletes should not be disqualified from sports, but regular follow-up with repeat tests is recommended.

**Competing interests** None.

**Patient consent** Obtained.

## REFERENCES

1. **Corrado D**, Pelliccia A, Heidbuchel H, *et al.*; Section of Sports Cardiology, European Association of Cardiovascular Prevention and Rehabilitation. Recommendations for interpretation of 12-lead electrocardiogram in the athlete. *Eur Heart J* 2010;**31**:243–59.
2. **Baggish AL**, Wood MJ. Athlete's heart and cardiovascular care of the athlete: scientific and clinical update. *Circulation* 2011;**123**:2723–35.
3. **Basavarajaiah S**, Wilson M, Junagde S, *et al.* Physiological left ventricular hypertrophy or hypertrophic cardiomyopathy in an elite adolescent athlete: role of detraining in resolving the clinical dilemma. *Br J Sports Med* 2006;**40**:727–9; discussion 729.

This pdf has been created automatically from the final edited text and images.

Copyright 2012 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.  
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Ghani S, Sharma S. Electrocardiographic changes in an athlete before and after detraining. *BMJ Case Reports* 2012;10.1136/bcr.01.2012.5520, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact [consortiasales@bmjgroup.com](mailto:consortiasales@bmjgroup.com)

Visit [casereports.bmj.com](http://casereports.bmj.com) for more articles like this and to become a Fellow

Keep up to date with all published cases by signing up for an alert (all we need is your email address) <http://casereports.bmj.com/cgi/alerts/etoc>