

Acute abdominal pain and diarrhoea in healthy adult

Bradford McGwire

Division of Infectious Diseases,
The Department of Internal
Medicine, The Ohio State
University, Columbus, Ohio,
USA

Correspondence to
Dr Bradford McGwire,
brad.mcgwire@osumc.edu

DESCRIPTION

An adult woman presented with 2 weeks of mild fatigue followed by 1 day of acute abdominal pain and explosive watery diarrhoea. Upon finding an object from her anus, slow extraction revealed a tape-like structure (see figure 1). Although living in the mid-west USA, she has travelled throughout the Middle East and sub-Saharan Africa as well as frequently consuming raw fish. The specimen identified as *Diphyllobothrium latum*, known as the broad tapeworm because the proglottids of this worm are broader than they are long (arrow figure 1B) as opposed to *Taenia* genus tapeworms. The parasite is acquired through the consumption of sparganum stage (plerocercoid larva) found in raw fish, when released attaches to the intestinal lining



Figure 1 (A) Specimen of *Diphyllobothrium latum* expelled from patient described is estimated at least 200 cm in length and is a partial specimen in that the head of the worm was not found and remained in the patient's intestinal tract along with an additional portion of the worm. Inset in (B) shows close-up of wide proglottids that are characteristic of this genus (black arrow).

and mature in some cases for years.^{1 2} Diagnosis can be made by the identification of eggs or portions of the adult parasite shed in the stool or by visualisation by endoscopy.³ Prolonged infection can lead to B₁₂ deficiency through scavenging of this vitamin by the parasite. This patient was given a single dose of praziquantel, the preferred therapy for diphyllobothriasis.

Learning points

- ▶ *Diphyllobothrium* infection is associated with the consumption of raw freshwater fish.
- ▶ Patients can be asymptomatic, present with abdominal pain and/or diarrhoea and/or megaloblastic anaemia.
- ▶ *Diphyllobothrium* is morphologically distinguished from other tapeworms by the presence of wide proglottids and an extremely long adult stage.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Scholz T, Garcia HH, Kuchta R, et al. Update on the human broad tapeworm (genus *Diphyllobothrium*), including clinical relevance. *Clin Microbiol Rev* 2009;22:146–60.
- 2 Church C, Neill A, Schotthoefer AM. Intestinal infections in humans in the rocky mountain region, United States. *J Parasitol* 2010;96:194–6.
- 3 Nomura Y, Fugiya M, Ito T, et al. Capsule endoscopy is a feasible procedure for identifying a *Diphyllobothrium nihonkaiense* infection and determining the indications for vermicide treatment. *BMJ Case Rep* 2010;10.1136/bcr.05.2010.3023 31 Aug 2010

To cite: McGwire B. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-009441

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmjjournals.com for more articles like this and to become a Fellow