

# Unilateral proptosis in a woman with asthma

Saori Sakaue, Shuji Sumitomo, Keishi Fujio, Kazuhiko Yamamoto

Department of Allergy and Rheumatology, The University of Tokyo Hospital, Bunkyo-ku, Tokyo, Japan


**Correspondence to**  
Dr Shuji Sumitomo,  
sumitos-ky@umin.ac.jp

Accepted 26 January 2015


## DESCRIPTION

A 47-year-old woman with a history of bronchial asthma presented with progressive proptosis of the right eye and diplopia. Her right eye showed apparent exophthalmos, impaired movement, conjunctival hyperaemia and episcleritis. Laboratory analysis revealed an increased titre of MPO-ANCA (4.3 IU/mL) and hypereosinophilia (1900 cells/ $\mu$ L, 22.5% on white cell count). MRI showed significant swelling of the extraocular muscles of the right eye and right ethmoid sinusitis (figure 1). Since a biopsy of extraocular muscles is unfeasible due to a risk of complications, the right bulbar conjunctiva was biopsied and dense infiltration of eosinophils with a granuloma formation (figure 2, arrows) was found. There were several phagocytised Charcot-Leyden crystals (figure 2, arrowheads) that were created by the destruction of eosinophils and release of their inner granules. The patient was diagnosed with eosinophilic granulomatosis with polyangiitis (EGPA) based on this histology, along with asthma, sinusitis and eosinophilia. It was speculated that unilateral proptosis was caused by the eosinophilic orbital myositis. Thirty milligrams of prednisone dramatically improved her proptosis and conjunctivitis.

Orbital myositis is an inflammatory process that involves the extraocular muscles. The typical symptoms are orbital pain, ocular movement impairment, diplopia and proptosis.<sup>1</sup> Orbital myositis is mainly idiopathic but sometimes caused by infection, thyroid dysfunction or systemic inflammatory diseases including antineutrophil cytoplasmic antibody-associated vasculitis.<sup>2</sup> However, the



**Figure 1** MRI showing swollen extraocular muscles of the right eye.



**Figure 2** Histopathology of right conjunctiva (H&E stain) showing granuloma formation (arrows) and Charcot-Leyden crystals (arrowheads).

prevalence of orbital manifestations in EGPA is much lower than that in granulomatosis with polyangiitis.<sup>3</sup> Our case demonstrates the diagnostic utility of bulbar conjunctival biopsy in such a major diagnostic challenge.

## Learning points

- ▶ Orbital myositis is sometimes caused by systemic diseases including infection, thyroid dysfunction or systemic inflammatory diseases such as antineutrophil cytoplasmic antibody-associated vasculitis.
- ▶ Eosinophilic granulomatosis with polyangiitis (EGPA) is a rare but important differential diagnosis in a patient with orbital myositis.
- ▶ Biopsy of bulbar conjunctiva is a useful diagnostic technique in EGPA with an eye presentation.

**Contributors** SSa and SSu wrote the manuscript and provided the figures. KF and KY supervised the manuscript.

**Competing interests** None.

**Patient consent** Obtained.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

- 1 Yuen SJ, Rubin PA. Idiopathic orbital inflammation: distribution, clinical features, and treatment outcome. *Arch Ophthalmol* 2003;121:491–9.
- 2 Schmidt J, Pulido JS, Matteson EL. Ocular manifestations of systemic disease: antineutrophil cytoplasmic antibody-associated vasculitis. *Curr Opin Ophthalmol* 2011;22:489–95.
- 3 Gordon LK. Orbital inflammatory disease: a diagnostic and therapeutic challenge. *Eye* 2006;20:1196–206.



**To cite:** Sakaue S, Sumitomo S, Fujio K, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-207532

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.  
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact [consortiasales@bmjgroup.com](mailto:consortiasales@bmjgroup.com)

Visit [casereports.bmj.com](http://casereports.bmj.com) for more articles like this and to become a Fellow