

Suspected case of lipoedema in Japanese woman with a characteristic histology in skin biopsy

Hiroyuki Koyama, Tomohiro Tanaka, Kenro Imaeda

Department of Gastroenterology and Metabolism, Nagoya City University Graduate School of Medical Sciences, Aichi, Japan

Correspondence to

Dr Tomohiro Tanaka,
ttnaka@kuhp.kyoto-u.ac.jp

Accepted 16 August 2017

DESCRIPTION

A 42-year-old Japanese woman with a body mass index of 42, presented with a long history of bilateral swelling of buttocks and lower extremities. The upper trunk, upper extremities and feet were spared of excessive fat deposition without a complete loss of adipose tissues (figure 1), lowering the likelihood of partial lipodystrophy. Physical examination revealed that the edema was dry, hard and non-pitting. Stemmer's sign was negative, and five out of seven criteria for the diagnosis of lipoedema¹ were met. A CT scan showed massive circumferential enlargement of subcutaneous tissues with the same CT value as fat, further suggesting the diagnosis of lipoedema (figure 1). Other conditions causing lower-extremity swelling or fat excess were excluded. We performed skin punch biopsy of the right hip (figure 2). Chief pathological findings were as follows: (1) the size of adipocytes was generally large (with a diameter up to 100 μ m); (2) the contour of adipocytes was maintained, but with membrane breakage; and (3) infiltration of plasma cells or lymphocytes typically observed in lymphoedema was scarce.

While the diagnostic pathology of lipoedema has not been established, hypertrophic and hyperplastic adipocytes are common pathological features.² We presume that hypertrophic and ruptured adipocytes in our specimen highly suggest a rare case of lipoedema in Japanese woman.



Figure 1 General appearance and CT scan of the lower extremities.




Figure 2 Histology of adipocytes with H&E.

Learning points

- Unexplained bilateral leg swelling presents potential diagnosis of lipoedema.
- Skin punch biopsy may be helpful for the diagnosis of lipoedema.

Contributors HK is an attending physician during hospitalisation and in charge of drafting the article. TT is the doctor in charge when the patient attends our hospital as an outpatient and he polished the article. KI is the head of our department. He gather everyone together and argue about the patient to achieve a correct diagnosis and treatment. He also had final approval of the article to be published.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Wold LE, Hines EA, Allen EV. Lipedema of the legs: a syndrome characterized by fat legs and edema. *Ann Intern Med* 1951;34:1243–50.
- 2 Buck DW, Herbst KL. Lipedema: a relatively common disease with extremely common misconceptions. *Plast Reconstr Surg Glob Open* 2016;4:e1043.



CrossMark

To cite: Koyama H, Tanaka T, Imaeda K. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-221049

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow