

Wellens' syndrome: a pattern to remember

Tiago Gama Ramires, Joana Sant'Anna, Joao Pais, Bruno Cordeiro Picarra

Hospital do Espirito Santo EPE,
Evora, Portugal

Correspondence to
Dr Tiago Gama Ramires,
tiago.raposo.ramires@gmail.com

Accepted 1 April 2018

DESCRIPTION

Electrocardiographic changes in asymptomatic individuals may be a diagnostic challenge. The knowledge and early identification of these patterns and their correct association with clinical picture shall have an urgent and specific management.

A 75-year-old Caucasian man with medical history of hypertension and peripheral arterial disease presented at the emergency department, transported by the prehospital emergency team, complaining of an intense chest pain, which woke him up, with irradiation to the left upper limb and diaphoresis. The patient denied nausea, vomiting, palpitations and previous complaints of angina as well. It was administered, during transport, sublingual nitroglycerine, with resolution of the pain.

The patient was vigilant, oriented and haemodynamically stable. Physical examination was unremarkable.

The admission ECG, without pain, showed sinus rhythm, heart rate of 60 bpm, right bundle branch block pattern and deep negative symmetrical T waves in the precordial derivations (V2 to V6) suggestive of type 2 Wellens' pattern (figure 1). Blood tests documented a mild elevation of troponin I (1.76 ng/mL (normal <0.012 ng/mL)), without other abnormalities.

The patient was admitted to the coronary intensive care unit with the diagnosis of acute coronary syndrome without ST-elevation.

During hospital stay, patient complained of chest pain. A new ECG was performed showing sinus rhythm, heart rate of 70 bpm, right bundle branch block pattern, no changes in ST segment and biphasic T waves in V3 (figure 2). The echocardiogram revealed mid-apical segment hypokinesia of the anterior wall conditioning mild depressed global systolic function with an estimated ejection fraction of 49%.




Figure 1 Hospital admission ECG, without chest pain, showing sinus rhythm, heart rate of 60 bpm, right bundle branch block pattern and deep negative symmetrical T waves in the precordial derivations V2 to V6 (arrow)—type 2 Wellens' pattern.




Figure 2 ECG performed during recurrent chest pain showing sinus rhythm, heart rate of 70 bpm, right bundle branch block pattern (arrow), no changes in ST segment and biphasic T waves in V3 (arrowhead).

Cardiac catheterisation documented a stenosis of 99% of the proximal segment of the left anterior descending (LAD) coronary artery. Angioplasty was performed with placement of one drug-eluting stent. The patient improved rapidly and was discharged home at fourth day of hospital admission.

Wellens' syndrome was first described by Zwann and Wellens in 1982 in a subgroup of patients with substantial risk of develop anterior myocardial infarction.^{1,2} This syndrome is characterised by T wave abnormalities in the anterior chest leads and reflects a critical obstruction in the proximal LAD coronary artery.¹⁻³

The clinical, laboratorial and ECG criteria for Wellens' syndrome is a history of angina, minimal or no elevation of cardiac enzymes, minimal or no ST-segment elevation (<1 mm), no pathological precordial Q waves and biphasic T waves in leads V2 and V3 (type 1) or deep, symmetrical and inverted T waves in leads V2 and V3, occasionally V1, V4, V5 and V6 (type 2).^{1,2}

In a symptomatic patient, T abnormalities can normalise or evolve into ST-segment elevation.^{1,2}

The ECG shall be the only indication of an imminent massive anterior infarction.² So, a timely diagnostic and therapeutic approach will be able to

Learning points

- ▶ Wellens' syndrome is characterised by T wave abnormalities in the anterior chest leads and reflects a critical obstruction in the proximal left anterior descending coronary artery.
- ▶ In the symptomatic patient, T changes can normalise or evolve into ST-segment elevation.
- ▶ Prompt diagnosis and therapeutic management will prevent a massive anterior myocardial infarction.



To cite: Ramires TG, Sant'Anna J, Pais J, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-224582

prevent a massive infarction in LAD coronary artery territory¹⁻³ and reduce the high morbidity and mortality associated with coronary artery disease.

Contributors Conception and design of the work; data collection and analysis: TGR. Drafting the article: TGR, JP and BCP. Critical revision of the article: JP, JS and BCP.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Parikh KS, Agarwal R, Mehrotra AK, *et al*. Wellens syndrome: a life-saving diagnosis. *Am J Emerg Med* 2012;30:255.e3–255.e5.
- 2 Pires de Morais G, Lopes A, Machado C, *et al*. Síndrome de Wellens—o padrão esquecido. *Revista portuguesa de cardiologia* 2009;28:641–4.
- 3 Rhinehardt J, Brady WJ, Perron AD, *et al*. Electrocardiographic manifestations of Wellens' syndrome. *Am J Emerg Med* 2002;20:638–43.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow