

# Purpura fulminans causing acute cortical necrosis in postpartum period

Mohan Kumar, Kundan Mishra, Vikas Suri, Savita Kumari

Internal Medicine, Post Graduate Institute of Medical Education and Research, Chandigarh, India

**Correspondence to**  
Dr Mohan Kumar,  
monu8501@gmail.com

Accepted 10 March 2018

## DESCRIPTION

A 22-year-old woman underwent normal vaginal delivery; 3 days later, she developed fever and rashes over all four limbs followed by decreased urine output. Examination revealed pedal oedema, non-palpable purpuric lesions over extremities with erosions at few places, and blackish discolouration of second and third toe of right foot (figure 1A). On investigation, haemoglobin was 6.7 g/dL, white blood cell 26 200/mm<sup>3</sup>, platelet 39 000/mm<sup>3</sup>, prothrombin time 25 s (control 12–14 s), activated thromboplastin time 38 s (control 24–30 s) and serum creatinine 9.1 mg/dL. Urine showed albumin 1+. Ultrasound was suggestive of bilateral acute renal parenchymal disease. Work-up for tropical infections was negative. Clinical diagnosis of puerperal sepsis causing purpura fulminans and acute kidney injury was made.

Patient was managed with haemodialysis and packed red blood cell infusion. She had persistent anuria and contrast-enhanced CT showed acute bilateral renal cortical necrosis (figure 1B). Skin biopsy showed erythema multiforme. She underwent amputation for digital gangrene. Hypercoagulable state work-up was done 6 weeks after the acute insult showed normal protein C, protein S, factor V and homocysteine level. Six months later, she continues to be anuric, and skin lesions have disappeared with mild hypopigmentation.

Renal cortical necrosis is a rare entity, caused by significantly diminished arterial perfusion of



**Figure 1** Clinical photograph showing multiple non-palpable purpuric lesions over extremities with erosions at few places and blackish discolouration of toes (A). CT scan of abdomen showing acute bilateral renal cortical necrosis (B).

## Learning points

- ▶ The clinical photograph shows the typical lesions associated with purpura fulminans.
- ▶ CT image shows the diagnostic appearance of acute cortical necrosis precluding renal biopsy, which has an increased risk of complications in the presence of Disseminated Intravascular Coagulation (DIC).
- ▶ Early and aggressive management pending evaluation is the key to successful outcome in patients with acute cortical necrosis.

the kidneys due to spasms of the feeding arteries, microvascular injury or disseminated intravascular coagulation and is the pathological progression of acute tubular necrosis.<sup>1 2</sup> Contrast-enhanced CT also shows characteristic findings such as lack of cortical enhancement, medullary enhancement and absent renal excretion.<sup>3</sup>

Early initiation of dialysis is the key to improved outcome. However, in most of the cases, recovery is not complete. If untreated, mortality rate exceeds 50%.<sup>2</sup>

**Contributors** MKH, KM, VS and SK were involved in the management of the case. MKH and KM wrote the manuscript, which was vetted by all authors.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent** Obtained.

**Provenance and peer review** Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

## REFERENCES

- 1 Matlin RA, Gary NE. Acute cortical necrosis. Case report and review of the literature. *Am J Med* 1974;56:110–8.
- 2 Prakash J, Singh VP. Changing picture of renal cortical necrosis in acute kidney injury in developing country. *World J Nephrol* 2015;4:480–6.
- 3 Agarwal A, Sakhuja V, Malik N, et al. The diagnostic value of CT scan in acute renal cortical necrosis. *Ren Fail* 1992;14:193–6.

Check for updates

**To cite:** Kumar M, Mishra K, Suri V, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-224669

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.  
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact [consortiasales@bmjgroup.com](mailto:consortiasales@bmjgroup.com)

Visit [casereports.bmj.com](http://casereports.bmj.com) for more articles like this and to become a Fellow