

Incidentally detected adrenocortical carcinoma in familial adenomatous polyposis: an unusual presentation of a hereditary cancer syndrome

Samarth Agarwal, Ashish Sharma, Deepanshu Sharma, Satyanarayan Sankhwar

Department of Urology, King George's Medical University, Lucknow, Uttar Pradesh, India

Correspondence to
Dr Ashish Sharma,
ashishuc343@gmail.com

Accepted 20 August 2018

DESCRIPTION

A 30-year-old woman presented to us with complaints of intermittent bleeding per rectum, vague abdominal pain and altered bowel habits for last 3 months. There was no history of headache, palpitations, diarrhoea, weight loss, skin lesions, haematuria and hypertension. Her grandmother had history of colostomy for acute intestinal obstruction and died of unexplained cause at age of 55 years. Her laboratory investigations were normal. On digital rectal examination, multiple polypoidal firm masses were felt approximately 1–5 cm from anal verge. On colonoscopy, number of polyps were seen in the colon around 40–50 in number, ranging from few millimetres to largest measuring around 3 cm. These polyps spread over ascending to sigmoid colon and rectum in increasing number. The biopsy from these polyps was suggestive of tubular adenomas. A definitive diagnosis of familial adenomatous polyposis (FAP) was made. FAP is associated with a myriad of presentations, and hence, a thorough work up was done to elicit associated lesions. An upper gastrointestinal endoscopy was performed, which was unremarkable. On slit-lamp examination, discrete flat multiple pigmented lesions were present in retina suggestive of congenital hypertrophy of retinal pigment epithelium (CHRPE). However, the patient did not complain of any eye symptoms.

Further evaluation with contrast-enhanced CT scan (CECT) of abdomen showed multiple




Figure 2 (A) Intraoperative picture showing total proctocolectomy with permanent end ileostomy. (B) The total proctocolectomy specimen showing multiple polypoidal lesions throughout the rectum and sigmoid.

enhancing polypoidal mass lesions (largest of size 3×2 cm) throughout the colon and rectum with no evidence of any regional or distant metastasis. To our surprise, CECT also revealed a heterogeneously enhancing, well-defined adrenal mass measuring $8.5 \times 8.5 \times 6.5$ cm with areas of necrosis suggestive of localised adrenocortical carcinoma (ACC) (figure 1). Functional adrenal work-up (serum aldosterone, serum cortisol and 24 hours urinary metanephrine and normetanephrine) results were within normal range.

After proper counselling, she underwent radical adrenalectomy with total proctocolectomy with permanent end ileostomy as shown in figure 2A. The total proctocolectomy specimen was examined and revealed multiple polypoidal lesions as shown in figure 2B. Blood sample of patient was analysed for truncated APC gene product, and in vitro protein synthesis assay was positive. Histopathology and immunohistochemistry (synaptophysin and inhibin) examination revealed high-grade ACC with negative surgical margin with pT3N0M0 staging. The postoperative period was uneventful. She was discharged on postoperative day 8. She was doing well at 4 weeks follow-up with no significant complains.

The FAP is an autosomal-dominant colorectal cancer syndrome, which is caused by mutation of APC gene on chromosome 5q21. It is characterised by occurrence of hundreds of adenomatous colorectal polyps and almost inevitable progression to colorectal cancer by the fourth decade of life. Other associated features of FAP include




Figure 1 Contrast-enhanced CT scan abdomen (coronal and transverse sections) showing heterogeneously enhancing, well-defined adrenal mass measuring $8.5 \times 8.5 \times 6.5$ cm with areas of necrosis suggestive of localised adrenocortical carcinoma.



© BMJ Publishing Group Limited 2018. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Agarwal S, Sharma A, Sharma D, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-226799

Images in...

upper gastrointestinal polyps, desmoid tumours, CHRPE and other extraintestinal malignancies. Uncommonly, endocrine tumours like parathyroid, pituitary, pancreatic and adrenal neoplasms have been described in these patients.¹

Due to widespread utilisation of modern imaging modalities like CT and MRI, the detection rates of adrenal incidentalomas has increased during the last decades. About 7% of patients with FAP or its variants have adrenal masses, as compared with 3% in general populations.² The biological behaviours of adrenal neoplasm in patients with FAP are comparable with sporadic

variety but its required longer follow-up.³ Mostly adrenal lesions are non-functional and benign, while functioning adrenal tumour mostly secretes cortisol. These non-functional adrenal tumours are asymptomatic and are usually detected incidentally or at autopsy. ACC is a rare neoplasm of adrenal gland with poor prognosis with incidence of one to two per million populations. The ACC may be associated with Li-Fraumeni, Lynch syndrome, Beckwith-Wideman, MEN type 1 and FAP. There are only few reported cases showing association between FAP and ACC.

Contributors SA: concept, design, supervision, processing, writing manuscript and critical analysis. AS: supervision, processing, writing manuscript and critical analysis. DS: concept, supervision, writing manuscript and critical analysis. SS: concept, supervision, writing manuscript and critical analysis.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Learning points

- ▶ Familial adenomatous polyposis (FAP) is an autosomal-dominant colorectal cancer syndrome, which is characterised by occurrence of hundreds of adenomatous colorectal polyps and almost inevitable progression to colorectal cancer by the fourth decade of life.
- ▶ The extraintestinal manifestation includes desmoid tumour, retinal lesions and endocrine tumours like parathyroid, pituitary, pancreatic and adrenal neoplasms.
- ▶ The clinical presentation and biological behaviours of adrenal neoplasm in FAP is comparable with sporadic cases, and same protocols are followed in these cases.
- ▶ Adrenocortical carcinoma is a rare, mostly non-functional neoplasm of adrenal gland with poor prognosis.

REFERENCES

- 1 Felner EI, Taweevisit M, Gow K. Hyperaldosteronism in an adolescent with gardner's syndrome. *J Pediatr Surg* 2009;44:e21–e23.
- 2 Smith TG, Clark SK, Katz DE, *et al.* Adrenal masses are associated with familial adenomatous polyposis. *Dis Colon Rectum* 2000;43:1739–42.
- 3 Will OC, Hansmann A, Phillips RK, *et al.* Adrenal incidentaloma in familial adenomatous polyposis: a long-term follow-up study and schema for management. *Dis Colon Rectum* 2009;52:1637–44.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow