Images in...

A subcutaneous cord over the chest in Mondor's disease

Joydeep Chakraborty, 1 Shashikiran Umakanth, 2 Kushaal Vikram, 1 BM Yashodhara 3

Correspondence to Dr Shashikiran Umakanth, shashikiranu@gmail.com

DESCRIPTION

A middle-aged man presented with pain over the right anterior aspect of chest. He gave history of doing heavy exercise (extreme yoga postures) few days ago. On examination, there was a tender area on right anterolateral part



Figure 1 The vertical cord and groove in the anterolateral chest, seen when the patient's right arm was raised above the shoulder.

of chest with a vertical cord like swelling just under the skin. On overhead abduction of the right arm, the cord like structure and adjacent longitudinal groove became prominent (figure 1). A clinical diagnosis of superficial thrombophlebitis – Mondor's disease was made based on the preceding history of strenuous exercise and the abovementioned findings. Complete blood count, erythrocyte sedimentation rate, C-reactive protein, coagulation parameters, liver and renal function tests, performed to rule out other aetiologies, were within normal limits. Patient was treated with ibuprofen and heparinoid ointment. He recovered completely within 2 weeks.

Mondor's disease is superficial thrombophlebitis of anterior chest wall seen more often in women. It is an uncommon condition that presents as sudden onset of chest wall pain. The phlebitic vein appears as a red, tender cord like structure. It may be preceded by local trauma, surgical trauma, muscular strain (strenuous exercise) or be associated with underlying connective tissue diseases, hypercoagulable states and malignancy. The diagnosis is usually based on typical history and examination findings. However, laboratory investigations can be done to rule out underlying aetiologies. Treatment includes hot fomentation, nonsteroidal anti-inflammatory drugs (NSAIDs) and local anticoagulants. It is self-limiting with spontaneous resolution within weeks.

Competing interests None.

Patient consent Obtained.

REFERENCES

- Kikano GE, Caceres VM, Sebas JA. Superficial thrombophlebitis of the anterior chest wall (Mondor's disease). J Fam Pract 1991;33:643

 –4.
- Giovanni André PV, Fabrício MO. Superficial thrombophlebitis (Mondor's disease) after breast augmentation surgery. *Indian J Plast Surg* 2008:41:219–21.
- Cesarone MR, Belcaro G, Agus G, et al. Management of superficial vein thrombosis and thrombophlebitis: status and expert opinion document. Angiology 2007;58:7S–14S.

¹Department of Medicine, Kasturba Medical College, Manipal, Karnataka, India;

²Department of Medicine, Melaka-Manipal Medical College, Udupi, India;

³Department of Medicine, Melaka-Manipal Medical College, Melaka, Malaysia

BMJ Case Reports

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit http://group.bmj.com/group/rights-licensing/permissions.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Chakraborty J, Umakanth S, Vikram K, Yashodhara BM. A subcutaneous cord over the chest in Mondor's disease. BMJ Case Reports 2011; 10.1136/bcr.04.2011.4131, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ► Submit as many cases as you like
- ► Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
 Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow