

A Baker's dozen

Katrina Sheikh,¹ Keith Siau²

¹Queen Elizabeth Hospital
Birmingham, West Midlands,
UK

²Department of
Gastroenterology, Queen
Elizabeth Hospital,
Birmingham, UK

Correspondence to
Dr Katrina Sheikh,
kat008@hotmail.com

Accepted 9 February 2015

DESCRIPTION

A 94-year-old woman with chronic left knee osteoarthritis was referred to the deep vein thrombosis (DVT) clinic with a 2-day history of non-traumatic left calf pain and swelling, with mild restriction in knee flexion. On examination, she had focal swelling extending from mid-calf to the back of her knee, without warmth or erythema and required support with a frame. Serum D-dimer was elevated at 439 µg/L (<250), with normal inflammatory markers. Doppler ultrasonography of the limb excluded DVT but revealed an effusion in the left popliteal fossa in keeping with Baker's cyst, interestingly, with the presence of round hyperechoic lesions (figure 1). Knee radiographs confirmed osteoarthritis with 12 irregular radiolucent fragments posterior to the knee joint, coined Baker's dozen (figures 2 and 3). Appearances were in keeping with synovial osteochondromatosis (SOC), probably secondary to long-standing osteoarthritis. Following orthopaedic review, the patient was managed conservatively with analgesia, leg elevation and physiotherapy and discharged with general practitioner (GP) follow-up.

SOC is a benign condition characterised by synovial hyperplasia and neoplasia into cartilage forming cells, resulting in cartilaginous nodules (chondromas), which typically calcify into osteochondromas.¹ This condition may be primary or secondary to a degenerative process, such as osteoarthritis. SOC is often



Figure 1 Ultrasound Doppler scan of the left popliteal fossa.



Figure 2 Lateral X-ray of the left knee.



Figure 3 Anteroposterior X-ray of the left knee.

monoarticular, affecting the knee, and may be a secondary cause of Baker's cyst.² Malignant transformation into synovial chondrosarcoma is rare. The diagnosis is confirmed radiologically. CT/MRI may delineate SOC from differentials including chondrosarcoma, synovial haemangioma and pigmented villonodular synovitis.³ Management may be conservative, or involve arthroscopic or open resection of osteochondromas and synovectomy.

Learning points

- ▶ Synovial osteochondromatosis (SOC) is a clinicrodiological diagnosis that should be considered when encountering intra-articular or periarticular nodules on routine imaging. In younger patients, CT/MRI should be considered to delineate it from differential diagnoses including osteosarcoma.
- ▶ In addition to trauma and arthropathy, Baker's cyst may also be secondary to SOC. Baker's cyst may rupture and mimic deep vein thrombosis.



To cite: Sheikh K, Siau K.
BMJ Case Rep Published
online: [please include Day
Month Year] doi:10.1136/
bcr-2015-209756

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

- 2 English S, Perret D. Posterior knee pain. *Curr Rev Musculoskelet Med* 2010;3:3–10.
- 3 McKenzie G, Raby N, Rithcie D. A pictorial review of primary synovial osteochondromatosis. *Eur Radiol* 2008;18:2662–9.

REFERENCES

- 1 Coles MJ, Tara HH. Synovial chondromatosis: a case study and brief review. *Am J Orthop* 1997;26:37–40.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow