

A case of a large, painless retroperitoneal mass causing hydronephrosis and biliary obstruction

Joseph Attard,¹ Christine Azzopardi,¹ Kelvin Cortis,² Jo Etienne Abela³

¹Mater Dei Hospital Malta, Msida, Malta

²Department of Radiology, Mater Dei Hospital, Msida, Malta

³Department of Surgery, Mater Dei Hospital, Msida, Malta

Correspondence to

Dr Christine Azzopardi,
chrissyazz@yahoo.com

Accepted 2 October 2015

DESCRIPTION

A 56-year-old man presented with abdominal distention over a few months. He was otherwise well and routine blood tests were normal. Examination

of the abdomen revealed a firm swelling. CT scan was performed for further evaluation.

A large, cystic, non-enhancing mass on the right side of the abdomen was observed on CT scan. This was closely related to the right psoas muscle with secondary compression of the right ureter and resultant right-sided hydronephrosis (figure 1).

Subsequent MRI demonstrated a 19.5×17×21 cm, high T2 signal intensity lesion in the right retroperitoneum. On T1 fat saturated sequences, it appeared hypointense with thin internal mildly enhancing septations (figure 2). Right-sided hydronephrosis was observed (figure 3) together with compression of the distal common bile duct causing mild intra and extra-hepatic biliary dilation (figure 4).

The mass was removed surgically (figure 5). It was gelatinous in nature and was sent to the




Figure 1 Coronal reformat demonstrating the mass in the right side of the abdomen causing secondary hydronephrosis (red arrow).




Figure 2 Axial contrast-enhanced T1 fat saturated image demonstrates a large hypointense mass arising from the right psoas muscle with mildly enhancing internal septations.




Figure 3 The mass is hyperintense on coronal and sagittal T2-weighted MR sequences. A hydronephrotic right kidney is shown (red arrow).




Figure 4 Coronal T2-weighted MR sequence showing a dilated common bile duct (red arrow), secondary to a large T2 hyperintense mass arising from the right psoas muscle.



To cite: Attard J, Azzopardi C, Cortis K, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2015-212422




Figure 5 Surgical specimen.

laboratory for histopathological diagnosis, which revealed that the mass was a myxoid liposarcoma.

Myxoid liposarcoma is the second most common adult soft tissue sarcoma after well-differentiated liposarcomas. It may be difficult to establish the diagnosis with MRI due to the lack of fat signal intensity and, without administration of contrast, the tumour may mimic cystic pathology, which includes benign lesions such as a hydatid cyst.¹ Gadolinium-enhanced MRI is therefore an essential part of the management.² Myxoid liposarcomas have a high risk of local recurrence (50%) and a 20% rate of metastasis.³

Learning points

- ▶ Myxoid liposarcoma is an indolent slow-growing malignant tumour of mesenchymal origin.
- ▶ Myxoid liposarcoma may lead to diagnostic confusion on imaging—contrast-enhanced MRI is paramount in establishing the diagnosis.
- ▶ Sarcomatous tumours should be included in the differential of retroperitoneal masses.

Contributors CA and JA performed the literature review and drafted the article. KC made the radiological diagnosis and JEA performed the surgery.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Bermejo A, De Bustamante TD, Martinez A, *et al*. MR imaging in the evaluation of cystic-appearing soft-tissue masses of the extremities. *Radiographics* 2013;33:833–56.
- 2 Sung MS, Kang HS, Suh JS, *et al*. Myxoid liposarcoma: appearance at MR imaging with histologic correlation. *Radiographics* 2000; 20:1007–19.
- 3 Loubignac F, Bourtoul C, Chapel F. Myxoid liposarcoma: a rare soft-tissue tumor with a misleading benign appearance. *World J Surg Oncol* 2009;7:42. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2678127&tool=pmcentrez&rendertype=abstract>

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow