

Risk of operating on the wrong site: how to avoid a never event

Joy C Edlin, Robin Kanagasabay

Department of Cardiothoracic Surgery, St George's Hospital, London, UK

Correspondence to
Joy C Edlin, joy.edlin@nhs.net

Accepted 27 March 2018

DESCRIPTION

A 23-year-old man was referred to our unit for surgery for a right-sided recurrent primary spontaneous pneumothorax. On the eve of his surgery, he was consented and marked for the procedure. On examination of the patient, a play symbol tattoo was noted on the left side of the patient's torso and pointing towards the patient's left side, with the potential to cause confusion of the site of surgery. A black arrow in indelible ink was drawn on the patient's right shoulder to indicate the correct site for surgery (figure 1).

The first item on the National Patient Safety Agency's (NPSA) list of never events is 'Wrong site surgery'. As preventative measures, the agency advocates the use of standardised patient wristbands and the WHO Surgical Safety Checklist.¹ Introduced by the WHO in 2008, the checklist was launched as a tool to help improve safety of surgery and save lives. The tool helps the care team ensure the correct patient undergoes the correct procedure and includes checkpoints for site of surgery and supportive imaging to be displayed in the operating theatre.

A checklist is only as good as the team that uses it, and to encourage coherent team work the NPSA published 'Five Steps to Safer Surgery' in 2010 outlining five communication steps to make surgery safer. These include the steps: briefing, sign-in, time-out, sign-out and debriefing.²

Despite these interventions, 84 wrong site surgery events were reported in National Health Service

Patient's perspective

'My parents always told me my tattoos would serve as a lesson one day, I don't think they pictured this. Glad I could provide a source of bemusement and hopefully education.'

Learning points

- ▶ Even the best preparations can be derailed by patient tattoos.
- ▶ The importance of clear communication with and examination of the patient combined with a review of radiographic imaging.
- ▶ The team brief is the first communication step to encourage teamwork and avoid mistakes.
- ▶ The tattoo on the opposite side of the surgical site should be highlighted here.

hospitals in England between April and September 2017.³

Although the patient's tattoo gave rise to quite a stir in the anaesthetic room and operating theatre, it reminded the entire team of the importance of correct site marking, the five perioperative communication steps and adhering to the WHO Surgical Safety checklist.

The patient recovered well from his right Video-assisted thoracoscopic surgery (VATS) bullectomy and pleurectomy and was discharged from hospital 2 days later.

Contributors JCE wrote the article, spoke to and consented the patient. She also took and edited the photograph. RK has edited the article.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 NPSA. Wrong site surgery. <http://www.nrls.npsa.nhs.uk/resources/collections/never-events/core-list/wrong-site-surgery/#> (accessed 3 Nov 2017).
- 2 NPSA. 'How to guide'. *Five steps to safer surgery*, 2010.



Figure 1 Which is the correct surgical site? Photograph of male torso showing a surgical site marking on the patient's right shoulder and a play symbol tattoo, pointing towards the patient's left side, on the left side of the thorax.



To cite: Edlin JC, Kanagasabay R. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-223704

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow