

Hepatic abscess secondary to gastric perforation

Luísa Graça,^{1,2} Bárbara Viamonte,³ Cláudia Carvalho,^{1,4} António Sarmento^{1,2}

¹Infectious Diseases, Centro Hospitalar de São João EPE, Porto, Portugal

²Faculty of Medicine, University of Porto, Porto, Portugal

³Radiology, Centro Hospitalar de São João EPE, Porto, Portugal

⁴Universidade do Porto Instituto de Saúde Pública, Porto, Portugal

Correspondence to

Professor António Sarmento, internosdi@gmail.com

Accepted 27 May 2019

DESCRIPTION

A 60-year-old woman with lumbar spinal stenosis presented to the emergency department with 2-day history of shivering and worsened lumbar pain. Apart from fever physical examination was unremarkable. Routine blood tests revealed elevated inflammatory parameters but chest radiography and abdominal ultrasound showed no relevant findings. She was hospitalised for 3 days during which she remained afebrile. The hypothesis of spinal infection was excluded by lumbar MRI. Blood cultures were negative. She was discharged without specific treatment but 3 days later she was readmitted with shivering.

The patient remained stable besides intermittent fever. However, 3 days after readmission she developed right costovertebral angle tenderness and blood cultures yielded *Escherichia coli*. Considering these data the hypothesis of renal abscess was considered. Accordingly, ceftriaxone 2g once per day was started and an abdominal CT was performed, revealing a hepatic abscess (42×42×28 mm) associated to a foreign body that had migrated from the stomach (figure 1). *Streptococcus constellatus* was also isolated in blood cultures. The next day a fish bone was extracted by laparoscopic surgery. Given this finding the patient was able to recall an episode of intense epigastric pain after eating codfish a few months earlier. Both *E. coli* and *S. constellatus* were isolated from drained pus.

After surgery the patient became afebrile, inflammatory parameters declined to normal range and follow-up blood cultures were negative. A follow-up CT, 5 days later, revealed only residual hypodense areas. The patient completed 14 days of ceftriaxone postsurgery. On follow-up appointment, 3 months later, she remained asymptomatic.

The diagnosis of hepatic abscess requires a high index of suspicion as the classic triad of fever, jaundice and right upper abdominal pain is present in a minority of cases.¹ Frequently, patients present with fever and non-specific symptoms such as nausea, vomiting and malaise.¹

Regarding pyogenic abscesses, biliary tract disease is the most frequent cause, followed by

portal seeding in the setting of bowel disease. Other causes include arterial seeding secondary to hepatic artery thrombosis, chemoembolisation or bacteraemia; direct extension from a contiguous focus of infection and lastly, penetrating trauma.¹ Since stomach perforation by fish bone is the most frequent cause of hepatic abscess due to trauma, micro-organisms from normal oropharynx flora are the most common aetiological agents.^{2,3} Interestingly, patients rarely recall the episode of ingestion^{2,3} and therefore diagnosis is frequently made by imaging showing the foreign body, fistulous tract or thickened gastrointestinal wall in contact with the liver.^{3,4} CT is the preferred modality due to its higher accuracy to detect foreign bodies and small posterior abscesses.³ In most cases foreign body removal is necessary since the success rate of antibiotics alone is 9.5%.⁴ Although laparotomy is frequently performed, minimally invasive techniques are increasingly used in this setting^{3,4} and there are a few reports of successful foreign body removal percutaneously^{5,6} or laparoscopically.⁷ Our patient was successfully treated with laparoscopic drainage and foreign body removal followed by 14 days of directed antibiotic therapy.

Learning points

- ▶ Hepatic abscess diagnosis requires a high index of suspicion. Patients may not have any other symptom besides fever.
- ▶ Migration of a foreign body from the gastrointestinal tract is a rare cause of hepatic abscess.
- ▶ Abscess drainage and foreign body extraction in addition to antibiotics is frequently required.

Contributors LG and BV drafted the paper; CC and AS revised it critically for important intellectual content. All authors approved the final version to be published.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Longworth S, Han J. Pyogenic liver abscess. *Clin Liver Dis* 2015;6:51–4.
- Santos SA, Alberto SC, Cruz E, et al. Hepatic abscess induced by foreign body: case report and literature review. *World J Gastroenterol* 2007;13:1466–70.
- Chong LW, Sun CK, Wu CC, CcW, et al. Successful treatment of liver abscess secondary to foreign body penetration of the alimentary tract: a case report and literature review. *World J Gastroenterol* 2014;20:3703–11.




Figure 1 CT showing a hepatic abscess (green arrow) associated to a foreign body (yellow arrow) that migrated from the stomach (red arrows).



© BMJ Publishing Group Limited 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Graça L, Viamonte B, Carvalho C, et al. *BMJ Case Rep* 2019;12:e230452. doi:10.1136/bcr-2019-230452

- 4 Leggieri N, Marques-Vidal P, Cerwenka H, *et al*. Migrated foreign body liver abscess: illustrative case report, systematic review, and proposed diagnostic algorithm. *Medicine* 2010;89:2.85–95.
- 5 Horii K, Yamazaki O, Matsuyama M, *et al*. Successful treatment of a hepatic abscess that formed secondary to fish bone penetration by percutaneous transhepatic removal of the foreign body: report of a case. *Surg Today* 1999;29:922–6.
- 6 Lau CW, Wong KM, Gogna A. Image-guided percutaneous transhepatic removal of fish bone from liver abscess. *J Radiol Case Rep* 2017;11:2.1–7.
- 7 Tan CH, Chang SY, Cheah YL. Laparoscopic removal of intrahepatic foreign body: A novel technique for management of an unusual cause of liver abscess-fish bone migration. *J Laparoendosc Adv Surg Tech A* 2016;26:47–50.

Copyright 2019 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow