Deep infiltrating endometriosis of the uterus involving the urinary bladder

Samarth Agarwal, Manoj Kumar, Ashish Sharma, Satyanarayan Sankhwar

Urology, King George's Medical University, Lucknow, Uttar Pradesh, India

Correspondence to
Dr Samarth Agarwal,
rebellite@gmail.com

Accepted 31 October 2018

DESCRIPTION

A 35-year-old female patient, married with two children, presented to us with complaints of haematuria for the last 1 week which was sudden in onset, painless and intermittent. She had lower abdominal pain 6 months ago when she was evaluated and diagnosed as a case of uterine endometriosis for which treatment had already been initiated. She received multiple sessions of monthly injectable leuprolide 22.5 mg and was currently on oral tablet dienogest (oral progestin) 2 mg daily for the last 6 months.

On physical examination, a mass was palpable in the infraumbilical region, which was firm in consistency with irregular margins and did not move with respiration. Her complete blood haemogram and renal function tests were normal.

On cystoscopic evaluation a solid mass around 3×2 cm in size was discovered on the left lateral wall of the urinary bladder with bilateral ureteric orifices visualised with clear efflux of urine. The biopsy showed presence of endometrial glands and stroma in the detrusor muscle. Contrast-enhanced CT of the abdomen revealed a bulky uterus with a soft tissue uterine mass of 9×5.4×5 cm in size involving the anterolateral wall of the uterine body and invading the left lateral wall of the urinary bladder with endoluminal extension of the mass, as shown in figures 1 and 2. Fat planes with other organs were maintained. A diagnosis of deep Infiltrating uterine endometriosis with secondary bladder endometriosis was made.

A consultation was taken from gynaecology. As the patient was already on treatment and the disease had progressed, a decision to perform open surgery was taken. After thorough counselling and proper consent, the patient underwent total hysterectomy with bilateral oophorectomy along with partial cystectomy of her pelvic endometriosis. Preoperatively, the left ureteric

Figure 2 Coronal CT section showing uterine endometriosis involving the left lateral wall of the urinary bladder with endoluminal infiltration.

orifice was stented with 5Fr ureteric catheter. Intraoperatively, the left adnexa was also found adherent to the uterine body. However, the left ureter was free. A cuff of bladder around 4×4 cm in size containing the endometrial tissue was then removed. The urinary bladder was primarily closed in two layers. A per urethral catheter along with an abdominal drain was also placed.

Her drain was removed after 48 hours and the remaining postoperative period was uneventful. At follow-up of 3 months, a repeat CT scan was done which did not reveal any recurrence.

Deep infiltrating endometriosis of the uterus by definition involves more than 5 mm of the peritoneum and has a prevalence of 1% in menstruating women. This condition involves the urinary bladder in 18%–55% of cases.¹

The urinary bladder is involved in 70%–85% of all cases of urinary tract endometriosis, followed by the ureter. It presents with lower urinary tract symptoms such as dysuria, frequency or haematuria.²

Combined hormonal contraceptives and progestogens are the first-line therapy for bladder endometriosis. They are safe, efficacious and easily tolerable. But according to the evidence available, they are not curative and only aid in temporarily suppressing the condition. Apart from this, bladder endometriosis can also be treated surgically. Both transurethral resection (TUR) surgery and segmental bladder resection have been described. In our case, we performed a partial cystectomy for two reasons: (1) hormonal therapy had failed; and (2) a TUR surgery in such an infiltrating exouterine mass would have inadvertently resulted in bladder perforation. It has shown to give excellent long-term results.^{3,4}

Figure 1 Transverse CT section showing bladder endometriosis involving the urinary bladder.

© BMJ Publishing Group Limited 2018. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Agarwal S, Kumar M, Sharma A, *et al.* *BMJ Case Rep* 2018;**11**:e226927. doi:10.1136/bcr-2018-226927

Learning points

- ▶ Endometriosis is not an uncommon anomaly and can have myriad of presentations with involvement of a number of organs such as the uterus, urinary bladder, ureters and so on.
- ▶ Hormonal therapy (oral contraceptives and progestogens) is considered as the first line of therapy for treatment of bladder endometriosis.
- ▶ If the disease progresses or the presentation is late, then surgery, either in the form of transurethral resection or open segmental bladder resection, can provide excellent long-term outcomes.

Contributors SA: concept, design, supervision, processing, writing the manuscript and critical analysis. MK: concept, design, supervision, processing. AS: supervision, processing, writing the manuscript and critical analysis. SS: concept, supervision.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Koninckx PR, Ussia A, Adamyan L, *et al*. Deep endometriosis: definition, diagnosis, and treatment. *Fertil Steril* 2012;98:564–71.
- 2 Knabben L, Imboden S, Fellmann B, *et al*. Urinary tract endometriosis in patients with deep infiltrating endometriosis: prevalence, symptoms, management, and proposal for a new clinical classification. *Fertil Steril* 2015;103:147–52.
- 3 Leone Roberti Maggiore U, Ferrero S, Candiani M, *et al*. Bladder endometriosis: a systematic review of pathogenesis, diagnosis, treatment, impact on fertility, and risk of malignant transformation. *Eur Urol* 2017;71:790–807.
- 4 Fedele L, Bianchi S, Zanonato G, *et al*. Long-term follow-up after conservative surgery for bladder endometriosis. *Fertil Steril* 2005;83:1729–33.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>. BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow