Earlobe pulsation: a sign of tricuspid regurgitation

Kiyoshi Shikino, Masatomi Ikusaka

General Medicine, Chiba
University Hospital, Chiba, Japan

Correspondence to
Dr Kiyoshi Shikino,
kshikino@gmail.com

Accepted 22 November 2018

DESCRIPTION

A 72-year-old woman had been experiencing shortness of breath for 2 weeks. Her jugular venous pressure was elevated, with prominent systolic pulsations and wagging earlobes ([figure 1](#) and [video 1](#)). The prominent pulsations were eliminated with gentle pressure at the base of the neck. Auscultation revealed a loud pulmonary component of the second heart sound and a holosystolic murmur at the left lower sternal border that increased with inspiration. Transthoracic echocardiography confirmed severe tricuspid regurgitation and a dilated inferior vena cava without respiratory fluctuation. The estimated pulmonary artery systolic pressure was >50 mm Hg. Right-sided congestive heart failure was diagnosed and treated with oral diuretics. Within 1 week, the symptoms improved ([figure 2](#) and [video 2](#)).

Figure 1 Elevated jugular venous pressure with earlobe pulsation.

Video 1 Elevated jugular venous pressure with earlobe pulsation.

Video 2 Earlobe pulsation disappeared after treatment.

Jugular venous pulsations ascend with congestive heart failure and may reach the earlobes, causing them to pulse.¹ In severe tricuspid regurgitation, one may see movements of the earlobes coincident with systole, as the C-V wave causes them to 'waggle' with each pulsation.¹ This C-V wave of severe tricuspid regurgitation resembles the large

© BMJ Publishing Group Limited 2018. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Shikino K, Ikusaka M. *BMJ Case Rep* 2018;**11**:e227861. doi:10.1136/bcr-2018-227861

Figure 2 Earlobe pulsation disappeared after treatment.

carotid-pulse wave of severe aortic regurgitation and is often confused with it.²

Learning points

- ▶ Earlobe pulsation is indicative severe tricuspid regurgitation.
- ▶ The C-V wave of severe tricuspid regurgitation resembles the large carotid-pulse wave of severe aortic regurgitation.

Contributors KS managed patient. KS wrote the draft. MI revised this article.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Chizner MA. The diagnosis of heart disease by clinical assessment alone. *Dis Mon* 2002;48:7–98.
- 2 Byrd MD. Lateral systolic pulsation of the earlobe: a sign of tricuspid regurgitation. *Am J Cardiol* 1984;54:244.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/> BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow