Bidirectional ventricular tachycardia due to hypokalaemia

Inês Santos, João Alves Teixeira, Catarina Costa, Luis Vale

Medicina 2.1, Hospital de Santo António dos Capuchos, Centro Hospitalar Lisboa Central, Lisboa, Portugal

Correspondence to

Dr Catarina Costa,
costa.catarina@gmail.com

Accepted 25 November 2018

DESCRIPTION

Bidirectional ventricular tachycardia (BDVT) is a regular ventricular tachyarrhythmia (VT) with two different QRS morphologies alternating at a rate typically between 140 and 180 bpm.¹ There are not many known related causes and the most common include digoxin toxicity, catecholaminergic polymorphic VT, myocarditis and myocardial infarction.^{2,3}

We report a case of a 81-year-old woman, with a known history of diabetes mellitus and hypertension, admitted at the emergency

department for prostration, diarrhoea and vomiting. Biochemistry tests on admission revealed severe ionic deficit, with a potassium level of 1.7 mmol/L. No serum digoxin levels were measured because there was no history of therapy with this drug. The 12-lead ECG (figure 1) revealed a BDVT pattern with a heart rate of 153 bpm. Endovenous potassium replacement was initiated with an immediate ECG pattern normalisation (figure 2). The authors present a case of BDVT, a rare arrhythmia with a cause not previously described.

Figure 1 Bidirectional ventricular tachycardia.

Figure 2 Sinus rhythm.

© BMJ Publishing Group Limited 2018. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Santos I, Alves Teixeira J, Costa C, et al. *BMJ Case Rep* 2018;**11**:e228195. doi:10.1136/bcr-2018-228195

Learning points

- ▶ Bidirectional ventricular tachycardia (BDVT) is a rare form of ventricular arrhythmia with a limited number of known causes described in the literature.
- ▶ As described in this case, hypokalaemia was assumed as the cause of BDVT.
- ▶ When confronted with this ECG pattern, hypokalaemia should be part of the differential diagnosis.

Contributors IS: planning, reporting, conception, interpretation of data. JAT: design, analysis. CC: conduct, acquisition of data. LV: analysis.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Richter S, Brugada P. Bidirectional ventricular tachycardia. *J Am Coll Cardiol* 2009;54:1189.
- 2 Valent S, Kelly P. Images in clinical medicine. Digoxin-induced bidirectional ventricular tachycardia. *N Engl J Med* 1997;336:550.
- 3 Park YH, Kim J. Bidirectional ventricular tachycardia in a patient with acute myocardial infarction and aortic stenosis. *Int J Cardiol* 2013;162:e41–e42.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow