OPEN ACCESS

CASE REPORT

Resolution of asthmatic symptoms following successful endoscopic resection of tracheal mucoepidermoid carcinoma

Abdul Hanif Khan,¹ Mohamed Faisal,² Razana Mohd Ali,³
Jamalul Azizi Abdul Rahaman⁴

¹Department of Medicine,
Universiti Putra Malaysia Fakulti
Perubatan dan Sains Kesihatan,
Serdang, Selangor, Malaysia

²Department of Respiratory,
Universiti Kebangsaan Malaysia
Medical Centre, Cheras, Kuala
Lumpur, Malaysia

³Department of Pathology,
Universiti Putra Malaysia Fakulti
Perubatan dan Sains Kesihatan,
Serdang, Selangor, Malaysia

⁴Department of Pulmonology,
Hospital Serdang, Kajang,
Selangor, Malaysia

Correspondence to

Dr Mohamed Faisal,
arabinose@hotmail.com

Accepted 15 November 2018

SUMMARY

Mucoepidermoid carcinoma (MEC) is a rare tumour of the trachea accounting for up to 0.2% of reported primary lung malignancy. We report a case of a 54-year-old man, ex-smoker, whose presentation mimicked adult onset asthma with cough and wheezing, which did not respond to conventional treatment. He had occasional haemoptysis and weight loss in which CT scan performed for malignancy screening showed a protruding mass in the distal trachea causing endobronchial obstruction. Bronchoscopic intervention was performed to relieve the obstruction that resulted in resolution of asthmatic symptoms. Histological diagnosis confirmed MEC. This case emphasised the importance of a high index of suspicion in an unusual presentation of a common disease and the pivotal role of bronchoscopic intervention in malignant central airway obstruction.

BACKGROUND

Mucoepidermoid carcinoma (MEC) is a rare tumour which originates from the mucus and serous glands of upper airway.¹ The incidence being reported as 0.1%–0.2% of primary lung cancer and accounted for 1%–5% of tracheobronchial adenoma.¹ Clinical presentation can be non-specific with cough, dyspnoea and wheeze, which could be misdiagnosed or mimics asthma as per our case presentation.

CASE PRESENTATION

A 54-year-old man, an ex-smoker (stopped 10 years ago), was referred to a tertiary centre for poorly controlled asthma, which was diagnosed 2 months prior to presentation. Further history revealed he had chronic cough with occasional haemoptysis, wheezing and significant weight loss of 10 kg of 3-month duration. No family history of malignancy was found. Physical examinations were unremarkable apart from the presence of inspiratory rhonchi with prolonged expiratory phase.

INVESTIGATIONS

Tuberculosis work-up was negative and chest X-ray (CXR) (figure 1) was normal. CT thorax (figure 2) was performed in view of red flag symptoms of malignancy, revealed a lobulated heterogeneous-enhancing soft tissue lesion at T3/T4 level measuring 1.2 × 1.1 × 1.3 cm with no clear plane from trachea

wall and no infiltration into the surrounding structures. In view of the CT findings, a flexible bronchoscopy was done and revealed a broad-based fungating mass obstructing 90% of the tracheal lumen; 8 cm from the vocal cord and 2 cm from main carina. No biopsy was attempted at this point. Spirometry with flow loop volume was initially planned but unfortunately was not performed in view of clinical deterioration.

DIFFERENTIAL DIAGNOSIS

Primary tracheal carcinoma.

Primary/metastatic endobronchial carcinoma.

Tracheobronchial tuberculosis.

Tracheobronchial fungal infection.

TREATMENT

Patient was admitted to ward for an elective rigid bronchoscopy. However, while in the ward, he developed worsening upper airway obstruction. Emergency rigid bronchoscopy was performed followed by electrocautery snaring to remove bulk of the tumour (figure 3A). Residual tumour was removed via rigid forceps (figure 3B) followed by application of argon plasma coagulation at the base of the lesion (figure 3C). However, complete excision of the tumour was unattainable.

Histological report revealed malignant cells of clear cell, intermediate cell, epidermoid cell and

Figure 1 Chest X-ray showing normal findings.

© BMJ Publishing Group
Limited 2019. Re-use
permitted under CC BY-NC. No
commercial re-use. See rights
and permissions. Published
by BMJ.

To cite:

Hanif Khan A, Faisal M,
Mohd Ali R, et al. *BMJ Case
Rep* 2019;**12**:e226202.
doi:10.1136/bcr-2018-
226202

Figure 2 CT thorax mediastinal window (A) and lung window (B) showing tracheal obstruction by a lobulated heterogeneous enhancing soft tissue lesion at T3/T4 level measuring 1.2×1.1×1.3 cm.

mucous cell type, arranged mainly in solid sheets and lobules with multicystic component. The cells are moderately pleomorphic with low proliferative index. A diagnosis of MEC of intermediate grade by Modified Healey grading was made (figure 4A,B).

OUTCOME AND FOLLOW-UP

Post procedure, patient was admitted in intensive care unit and was extubated after 2 days. He had an uneventful recovery and was discharged after completion of 1 week of antibiotic for postobstructive pneumonia. The ‘asthmatic’ symptoms resolved following our intervention. Patient was subsequently referred to oncology due to incomplete excision during bronchoscopy for endobronchial brachytherapy; however, he refused treatment and subsequently lost to follow-up.

DISCUSSION

Primary tracheal tumours are rare where in adults are overwhelmingly malignant as opposed to be benign nature in paediatric population.² MEC accounts for only 0.1%–0.2% of primary tracheal tumours and was first reported in 1952.¹ It is asymptomatic in 25% of patients as the tumour is small or located behind the carina and would only be incidentally found on CXR.³ However, the presence of endobronchial exophytic mass may present with obstructive symptoms (cough, wheeze, stridor) and haemoptysis.^{3,4} The presence of wheezing may lead to misdiagnosis of asthma as in our patient. Previous reported cases have shown that MEC can mimics asthmatic symptoms even in children and may also be misdiagnosed as invasive thyroid cancer.^{5,6}

As in any patients with tracheal mass, spirometry with flow loop volume may show variable extrathoracic obstruction which may improve following intervention. Unfortunately in our patient spirometry was not feasible as his condition deteriorated rapidly needing emergency bronchoscopic intervention.

As for imaging, CT scan is a valuable tool in detecting endobronchial tumours as its detection on CXR can be challenging. It is also useful in evaluating the extent of tumour involvement beyond airway, regional lymphadenopathy and pulmonary

Figure 4 Solid sheets of tumour with intervening thin fibrous septae forming lobules with multicystic component (A) using H&E stain (×100). The tumour cells are composed of clear, intermediate and mucous cell type (B) using H&E stain (×200).

nodules, which are important in determining stage and prognosis of disease.

The gold standard of diagnosing MEC would be bronchoscopy with adequate biopsy as preoperative misdiagnosis as squamous cell carcinoma, adenocarcinoma, bronchial adenoma or unknown mass was reported before.³ Immunostaining with cytokeratin 7 (CK7) antibody and thyroid transcription factor 1 has been used to distinguished MEC with primary lung adenocarcinoma where CK7 is usually positive in the former.⁷

Prognosis of the disease depends on the grade of the tumour and possibility of complete resection.⁴ In a low and intermediate grade tumour, a complete resection may not require subsequent adjuvant chemotherapy or radiotherapy.³ However, a high-grade tumour carries an estimated 5-year survival of 31%.³ Song *et al*, in a retrospective study on 32 patients with MEC, found significantly better survival in patients with histologically low-grade tumours than in patients with high-grade tumours.⁸

Currently, there are no randomised controlled trials in determining the most efficacious treatment modalities of tracheal tumour; however, surgical resection coupled with or without radiotherapy is considered the treatment of choice.^{8,9} Surgical should be considered if the involved tracheal segment can be resected and reconstructed with primary anastomosis.¹⁰ However, its role may be limited due to anatomical limitations, especially if located near the carina; metastatic disease or poor medical conditions and their contributions mainly revolved around palliative management.^{8,10}

Bronchoscopic evaluation is the key step in the treatment of patient with malignant airway obstruction as it not only allows diagnostic confirmation and staging when considering surgery, but also provides significant palliative relief in unresectable lesion, relieve life-threatening obstruction and provide a pathway for other definitive intervention, for example, surgery, radiation or chemotherapy.⁸ Reports by Hosakawa *et al* and El-Sameed *et al* demonstrate the effectiveness of bronchoscopic interventions in managing central obstruction due to tracheal tumour.^{11,12}

Figure 3 Bronchoscopic resection of the tumour using electrocautery snaring (A), removal of residual tumour with rigid forceps (B) and application of argon plasma coagulation at the base of the lesion (C).

In conclusion, MEC is a rare tumour of the trachea, which may present with obstructive airway symptoms mimicking asthma. This case highlights the importance of high index of suspicion coupled with diligent use of imaging and bronchoscopy that could prevent the delay in diagnosing this condition. Although surgical resection may provide definitive management, bronchoscopic intervention plays a pivotal role especially in providing diagnosis, staging, immediate relief and even cure in malignant central airway obstruction.

Learning points

- ▶ Patients with asthmatic symptoms who failed to respond to medications warrants further investigations, for example, CT thorax or bronchoscopy to look for alternative diagnosis.
- ▶ Mucoepidermoid carcinoma is a rare tumour that may present with obstructive airway symptoms leading to misdiagnosis of asthma.
- ▶ Bronchoscopic intervention has a significant role in management of malignant airway obstruction.

Contributors AHYK, MFAH and RMA were involved in the conception of the case, report, review of literature and writing the manuscript. JAAR revised this critically and contributed to important intellectual content. All authors participated in the final revision of the manuscript and approved the final manuscript and are also accountable for the accuracy of the content.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Heitmiller RF, Mathisen DJ, Ferry JA, *et al.* Mucoepidermoid lung tumors. *Ann Thorac Surg* 1989;47:394–9.
- 2 Junker K. Pathology of tracheal tumors. *Thorac Surg Clin* 2014;24:7–11.
- 3 Vadasz P, Egervary M. Mucoepidermoid bronchial tumors: a review of 34 operated cases. *Eur J Cardiothorac Surg* 2000;17:566–9.
- 4 Chin CH, Huang CC, Lin MC, *et al.* Prognostic factors of tracheobronchial mucoepidermoid carcinoma—15 years experience. *Respirology* 2008;13:275–80.
- 5 Lin CH, Chao YH, Wu KH, *et al.* Primary mucoepidermoid carcinoma at the carina of trachea presenting with wheezing in an asthmatic child mimicking an attack of asthma: A case report. *Medicine* 2016;95:e5292.
- 6 Moon EJ, Kim BY, Chung JY, *et al.* Rare tracheal mucoepidermoid carcinoma with invasion of thyroid gland initially misdiagnosed as invasive thyroid cancer. *Int J Clin Exp Med* 2016.
- 7 Shilo K, Foss RD, Franks TJ, *et al.* Pulmonary mucoepidermoid carcinoma with prominent tumor-associated lymphoid proliferation. *Am J Surg Pathol* 2005;29:407–11.
- 8 Song Z, Liu Z, Wang J, *et al.* Primary tracheobronchial mucoepidermoid carcinoma—a retrospective study of 32 patients. *World J Surg Oncol* 2013;11:62.
- 9 Jeon K, Kim H, Yu CM, *et al.* Rigid bronchoscopic intervention in patients with respiratory failure caused by malignant central airway obstruction. *J Thorac Oncol* 2006;1:319–23.
- 10 Madariaga MLL, Gaissert HA. Overview of malignant tracheal tumors. *Ann Cardiothorac Surg* 2018;7:244–54.
- 11 Hosokawa Y, Nozaki T, Furuichi S, *et al.* Successful resection of endobronchial mucoepidermoid carcinoma by a high-frequency surgical snare. *Journal of Bronchology* 2007;14:295–6.
- 12 El-Sameed YA, Al Marzooqi SH. Primary mucoepidermoid carcinoma of the lung. *J Bronchology Interv Pulmonol* 2012;19:203–5.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow