Intravaginal torsion of undescended testes

Filipe La Fuente Carvalho ,¹ Ish Talati,² Aaron Krill³

¹Department of Urology, MedStar Georgetown University Hospital, Washington, District of Columbia, USA

²Georgetown University School of Medicine, Washington, District of Columbia, USA

³Division of Urology, Children's National Medical Center, Washington, District of Columbia, USA

Correspondence to

Dr Filipe La Fuente Carvalho; filipe.decarvalho@gunet.georgetown.edu

Accepted 13 October 2019

DESCRIPTION

A 5-month-old male infant status post routine neonatal circumcision with known left inguinal testis presented to the emergency room with swelling and redness in the left inguinal region for over 24 hours. Physical examination revealed an indurated inguinal bulge with associated erythema (figure 1). The right testis was normal, and the left hemiscrotum was empty. Scrotal ultrasound revealed an enlarged, heterogeneous testis without Doppler flow. He was emergently taken to the operating room for inguinal exploration which confirmed an intravaginal torsion, non-viable testis (figure 2) and closed inguinal canal. Left orchiectomy and right scrotal orchiopexy were performed without complication. The final pathology revealed an infarcted and haemorrhagic testicle and epididymis.

Undescended testes (UDTs) affect up to 8% of full term, but 33% of premature male neonates, as the testis enters the inguinal canal after 28 weeks' gestation.¹⁻³ Insulin-like 3 and its receptor, relaxin family peptide receptors (RXFP), may play a role in the transabdominal migration as their loss results in UDTs in animal models.⁴ Seventy per cent of UDTs are palpable in the upper scrotum or low inguinal canal. Of non-palpable UDTs, 55%, 35% and 15% are in the abdomen, inguinal canal or absent, respectively.⁵ A patient with solitary UDT and hypospadias requires evaluation for a disorder of sexual development, but in patients without penile anomalies, UDT treatment can be delayed until 6 months of age.⁶ The scrotal location supports

Figure 2 Gross pathology of the left inguinal testis showing necrosis.

testicular function such that UDTs are associated with subfertility and germ cell malignancies.¹⁻³

The aetiology of torsion in UDTs is unknown, but may be related to abnormal cremasteric muscle contractions or gubernacular attachments. Intravaginal torsion (torsion of the testis within the tunica vaginalis) is more common in infants over 30 days of age and those with UDTs, compared with extravaginal torsion (torsion of the testis and tunica vaginalis as a unit) which is more common in neonates.⁷⁻⁹ While the incidence of torsion in UDTs is unknown, it is higher than the incidence in orthotopic testes.^{1 2 10} As demonstrated in this case, findings include abdominal and groin pain with ipsilateral inguinal swelling and erythema. Infants may present with inconsolable crying⁶ and should be evaluated emergently with a differential diagnosis including incarcerated inguinal hernia, intussusception and viral gastroenteritis.¹²

Doppler ultrasound has a sensitivity of 77.8% and specificity of 90% for diagnosing testicular torsion. However, its accuracy in UDTs decreases due to difficulty evaluating the inguinal canal.¹ As such, in suggestive cases with acute pain, inguinal swelling and empty ipsilateral hemiscrotum, surgical exploration should not be delayed. Surgery within 6 hours of torsion is generally successful, however due to often delayed diagnosis, the success rate in UDT torsion is below 40%.^{2 11} While the specific risk of contralateral torsion in UDTs is unknown, contralateral orchiopexy is recommended due to the risk of metachronous torsion and subsequent anorchia, in addition to the low morbidity of the procedure.^{7 9 11}

Figure 1 Physical examination of the patient in the emergency room revealed an indurated left inguinal bulge with associated erythema.

© BMJ Publishing Group Limited 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Carvalho FLF, Talati I, Krill A. *BMJ Case Rep* 2019;**12**:e231244. doi:10.1136/bcr-2019-231244

Images in...

Our case illustrates the typical presentation of torsion of UDTs. As orchiopexy is routinely delayed until 6 months, emergency physicians, paediatricians and parents must be aware that torsion of the UDT is possible in the interim. Parents and paediatricians should be aware that changes in the inguinal region associated with UDT require emergent evaluation.

Learning points

- ▶ Acute inguinal pain or swelling in the setting of an empty ipsilateral hemiscrotum or history of undescended testis requires urgent surgical evaluation.
- ▶ Torsion of an undescended testis is associated with poor salvage rates due to delay in diagnosis.
- ▶ Parents and clinicians should be aware that torsion of an undescended testis can occur while awaiting spontaneous descent or surgical correction.

Contributors FLFC and AK had full access to all the data in the manuscript and take responsibility for the integrity of the data and the accuracy of the events. IT drafted the manuscript, provided technical and material support.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Filipe La Fuente Carvalho <http://orcid.org/0000-0003-2378-6512>

REFERENCES

- 1 Moore CP, Marr JK, Huang CJ. Cryptorchid testicular torsion. *Pediatr Emerg Care* 2011;27:121–3.
- 2 Pogorelić Z, Mrkljić I, Jurić I, et al. Testicular torsion in the inguinal canal in children. *J Pediatr Urol* 2013;9:793–7.
- 3 Barthold JS. Abnormalities of the Testis and Scrotum and their Surgical Management. In: McDougal WS, Kavoussi LR, Novick AC, eds. *Campbell Walsh urology*. 10th ed. Philadelphia, PA: WB Saunders, 2012: 642–5.
- 4 Kalfa N, Gaspari L, Ollivier M, et al. Molecular genetics of hypospadias and cryptorchidism recent developments. *Clin Genet* 2019;95:122–31.
- 5 Denes FT, Saito FJ, Silva FA, et al. Laparoscopic diagnosis and treatment of nonpalpable testis. *Int Braz J Urol* 2008;34:329–35.
- 6 Radmayr C, Dogan HS, Hoebeke P, et al. Management of undescended testes: European association of Urology/European Society for paediatric urology guidelines. *J Pediatr Urol* 2016;12:335–43.
- 7 Monteilh C, Calixte R, Burjonrappa S. Controversies in the management of neonatal testicular torsion: a meta-analysis. *J Pediatr Surg* 2019;54:815–9.
- 8 Mano R, Livne PM, Nevo A, et al. Testicular torsion in the first year of life--characteristics and treatment outcome. *Urology* 2013;82:1132–7.
- 9 Bordin G, Parolini F, Morandi A, et al. Intravaginal testicular torsion in newborns: to fix or not to fix the contralateral testis? *Pediatr Med Chir* 2013;35:225–7.
- 10 Zilberman D, Inbar Y, Heyman Z, et al. Torsion of the cryptorchid testis--can it be salvaged? *J Urol* 2006;175:2287–9.
- 11 Naouar S, Braiek S, El Kamel R. Testicular torsion in undescended testis: a persistent challenge. *Asian J Urol* 2017;4:111–5.

Copyright 2019 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/> BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow