OPEN ACCESS

Pericardial knock

Trenton E Burgess,¹ Ngoc N Le, Gary S Olds, Peter D Sullivan,² André Martin Mansoor²

¹School of Medicine, Oregon Health and Science University, Portland, Oregon, USA

²Department of Internal Medicine, Oregon Health and Science University, Portland, Oregon, USA

Correspondence to

Dr André Martin Mansoor;
mansooran@ohsu.edu

Accepted 10 December 2019

DESCRIPTION

A 69-year-old woman with a history of systemic lupus erythematosus complicated by recurrent episodes of acute pericarditis was admitted to the hospital with chronic, progressive dyspnoea. Physical examination was notable for a jugular venous pressure (JVP) of 18 cm H₂O, with paradoxical rise on inspiration (Kussmaul's sign). The y descent of the jugular venous waveform was observed to be sharp and deep (Friedreich's sign). Coinciding with the nadir of the y descent, an extra, early diastolic heart sound was heard over the apex. It occurred just after the second heart sound (S2) and was high-pitched, heard best with the diaphragm of the stethoscope. A digital stethoscope was used to record the heart sounds. Simultaneously, an antique phonocardiograph was used to produce visualisation of the sounds in the form of a phonocardiogram. Combining these results, a video was created for analytic and teaching purposes (video 1). The qualities of the extra sound, including pitch, timing

and location, were consistent with that of a pericardial knock. In a patient with a history of recurrent episodes of acute pericarditis, the constellation of elevated JVP, Kussmaul's sign, Friedreich's sign, and a pericardial knock led to the diagnosis of constrictive pericarditis. A pericardial knock can be difficult to distinguish from an S3 gallop; it is a high-pitched, diastolic sound heard 0.09 to 0.12 s after the aortic component of S2 (A2).^{1–3} It tends to be louder, higher-pitched, and slightly closer to A2 than the low-pitched S3 gallop, which occurs 0.1 to 0.2 s after A2.^{2,3} The sound is presumed to occur when stiff and thickened pericardium cause sudden arrest of ventricular filling during diastole.^{2,3}

Contributors AMM captured the audio and phonocardiographic tracing of the heart sounds. TEB created the synchronised video. TEB,>NNL, GSO, PDS, and AMM were involved in writing the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- 1 Wood P. Chronic constrictive pericarditis. *Am J Cardiol* 1961;7:48–61.
- 2 Marriott HJL. *Bedside cardiac diagnosis*. Philadelphia, PA: Lippincott Company, 1993.
- 3 Tavel ME. *Clinical phonocardiography and external pulse recording*. 2nd edn. Chicago, Illinois: Year Book Medical Publishers, Inc, 1967.

Learning points

- The pericardial knock is a high-pitched, early diastolic sound that occurs when unyielding pericardium results in sudden arrest of ventricular filling. It can be an important clue to the diagnosis of constrictive pericarditis.^{2,3}
- The pericardial knock occurs earlier in diastole (0.09–0.12 s after the aortic component of the second heart sound (A2)) compared with the S3 gallop (0.1–0.2 s after A2).^{1–3}
- Phonocardiography combined with modern digital recording devices can be used to facilitate the understanding and recognition of extra heart sounds.

© BMJ Publishing Group Limited 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Burgess TE, Le NN, Olds GS, et al. *BMJ Case Rep* 2019;12:e233546. doi:10.1136/bcr-2019-233546

Copyright 2019 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow

video 1 The combination of an audio recording and a phonocardiographic tracing of a pericardial knock in a patient with constrictive pericarditis, demonstrating its pitch and timing relative to the second heart sound (S2).