A case of chronic lead poisoning with herbal-based medication

Siddharth Jain,[✉] Ashish Gupta, Animesh Ray,[✉] Naval Kishore Vikram

Department of Medicine, All India Institute of Medical Sciences, New Delhi, India

Correspondence to
Dr Animesh Ray,
doctoranimeshray@gmail.com

Accepted 8 March 2019

DESCRIPTION

A 59-year-old man, who is a known case of type 2 diabetes mellitus and hypertension for 3 years, presented to us with complaints of colicky abdominal pain, decreased appetite and generalised weakness for 3 months. There was no history of vomiting, jaundice and fever. His general examination was unremarkable apart from pallor. Systemic examination was also unremarkable. Initial investigations revealed a haemoglobin level of 10.1 g/dL (red blood cell count of 3.33×10^6 cells/ μ L), and peripheral smear showed microcytic hypochromic anaemia with mild *basophilic stippling* (figure 1). Other investigations, including abdominal ultrasound, were essentially normal. On reviewing the case, he gave a history of taking herbal-based medication for diabetes from an indigenous medical practitioner for 1 year. Examination revealed a bluish pigmentation of the gum-tooth line or the *Burtonian line* (figure 2). There was no neurological deficit. Laboratory investigations showed high blood lead levels (91 μ g/dL). He was treated as a case of chronic lead poisoning with intramuscular British anti-Lewisite 250 mg 4 hourly for 14 days (disodium calcium edetate and succimer were not available at our centre). There were no complications or side effects with therapy. Post treatment, his blood lead levels reduced to 40 μ g/dL. The Burtonian line disappeared and other symptoms alleviated.

Lead poisoning often causes colicky abdominal pain, fatigue, anaemia and neurological symptoms like foot/wrist drop and seizures. Examination commonly reveals the Burtonian line and peripheral neuropathy. Basophilic stippling¹ with microcytic hypochromic anaemia and high blood lead levels are the main laboratory features. It is treated

Figure 2 Lower gum-tooth line showing bluish deposits of lead sulfide, called the Burtonian line (white arrow).

with chelation therapy² with oral dimercapto-succinic acid or intravenous/intramuscular disodium calcium EDTA or British anti-Lewisite. Common sources of lead exposure include lead paint, lead-acid batteries, soil contamination near factories, lead soldering, cosmetics and herbal-based medications.³ Incidence rate has significantly reduced as compared with that of the past due to stringent policies, so it is commonly missed by clinicians these days. The suspected source of exposure in our patient was herbal-based medication. Though toxicological analysis was not performed, absence of any other source of poisoning and circumstantial evidence of herbal-based medicinal use, which have been widely reported to cause lead poisoning, lent support to our suspicion.

Learning points

- ▶ Indigenous medicinal preparations can lead to chronic lead poisoning producing characteristic symptoms, signs and laboratory features.
- ▶ Having a high index of suspicion and eliciting history of usage of purported drugs are important for diagnosis.
- ▶ Removal of source of poisoning and chelation therapy are crucial strategies for cure.

Acknowledgements Dr Pawan Gabra had contributed as an intern posted in our unit.

Contributors SJ and AG were involved in treating and daily management of the patient. AR devised the treatment protocol. NV was the chief supervisor and the head of the unit.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Figure 1 Peripheral smear of the patient showing a microcytic hypochromic picture with RBC showing basophilic stippling (black arrow). RBC, red blood cell.

© BMJ Publishing Group Limited 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Jain S, Gupta A, Ray A, et al. *BMJ Case Rep* 2019;**12**:e227954. doi:10.1136/bcr-2018-227954

REFERENCES

- 1 Cheson BD, Rom WN, Webber RC. Basophilic stippling of red blood cells: a nonspecific finding of multiple etiology. *Am J Ind Med* 1984;5:327–34.
- 2 Porru S, Alessio L. The use of chelating agents in occupational lead poisoning. *Occup Med* 1996;46:41–8.
- 3 Environmental Health and Medicine Education. Lead Toxicity: Where is Lead Found? *Agency for Toxic Substances & Disease* 2017.

Copyright 2019 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow