OPEN ACCESS

Left-sided superior vena cava complicates central line insertion

Joseph Alexander Gallaher,¹ Karthik Somasundaram²

¹Anaesthesia, Maidstone and Tunbridge Wells NHS Trust, Tunbridge Wells, UK
²ITU, St Peters Hospital, Surrey, UK

Correspondence to

Dr Joseph Alexander Gallaher, josephgallaher89@gmail.com

Accepted 5 May 2019

DESCRIPTION

A 39-year-old man with necrotising pancreatitis was admitted to the intensive care unit (ICU). On day 14, his right-sided central venous catheter (CVC) was replaced with a left-sided catheter. A confirmatory chest X-ray revealed that rather than charting its usual course across the left brachiocephalic vein into the superior vena cava (SVC), the catheter journeyed straight down into the left atrium (figure 1). A CT thorax (figure 2) revealed the patient had two SVCs, and the patient's left-sided SVC had been unintentionally catheterised.

A percentage range of 0.3%–0.5% of the population have two SVCs or a persistent left SVC (PLSVC). There is an increased (1.3%–4.5%) prevalence of PLSVC in patients with congenital heart disease.¹ It occurs due to failure of the left anterior cardinal vein to obliterate during fetal development and is an incidental finding in the vast majority of cases. Ninety-two per cent of PLSVCs drain into the coronary sinus, but the remaining 8% drain into the left atrium causing a right-left shunt which is usually insignificant.² However, as a result of the latter group, central catheters pose a risk of systemic embolisation so should be removed if the patient's cardiothoracic anatomy has not been formally assessed. If there is no shunt present, catheters can remain though with due care and attention to the nearby coronary sinus and the associated risk of arrhythmias from a misplaced catheter tip.

In this case, the exact position of drainage from the extra SVC was uncertain so the CVC was replaced with another right-sided catheter. The patient was eventually discharged from the ICU and went on to make a full recovery.

Figure 1 Anteroposterior chest X-ray (AP CXR) shows the left-sided CVC tracking into the left atrium. The right SVC is already catheterised. CVCs are indicated by the red arrows. CVC, central venous catheter; SVC, superior vena cava.

Figure 2 CT thorax shows two superior vena cavae as indicated by red arrows.

Learning points

- ▶ Consider a persistent left superior vena cava (PLSVC) when a left-sided central venous catheter appears in an unusual position in the left thorax.
- ▶ Left-sided superior vena cava catheters can remain in situ though only if formal imaging (eg, echocardiography) of the cardiothoracic anatomy confirms the absence of a shunt.
- ▶ PLSVC is often associated with congenital cardiac abnormalities.

Contributors JAG, a senior house officer working in intensive care, inserted the central line and found the unusual course on chest X-ray before enlisting the help of consultants to discover what had occurred. A CT thorax revealed the presence of two superior vena cavae (SVC) and the fact that the left-sided SVC had in fact been cannulated originally. KS was JAG's clinical supervisor at the time and supervised the writing of the report—suggesting changes where appropriate. We hope this case report educates medical and nursing staff with respect to the potential dangers of left-sided SVC cannulation.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Higgs AG, Paris S, Potter F. Discovery of left-sided superior vena cava during central venous catheterization. *Br J Anaesth* 1998;81:260–1.

© BMJ Publishing Group Limited 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Gallaher JA, Somasundaram K. *BMJ Case Rep* 2019;12:e228474. doi:10.1136/bcr-2018-228474

Copyright 2019 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow