Varicose veins causing tibial nerve compression in the tarsal tunnel

Rudi Falovic,¹ Mithun Nambiar,^{1,2} Pamela Boekel,¹ Joanna Lenaghan¹

¹Department of Orthopaedic Surgery, Maroondah Hospital, Ringwood East, Victoria, Australia

²Department of Surgery, University of Melbourne, Parkville, Victoria, Australia

Correspondence to

Dr Mithun Nambiar, mithunnambiar1@gmail.com

Accepted 27 April 2019

DESCRIPTION

A 66-year-old retired man presented with a 12-month history of atraumatic intermittent right foot pain and altered sensation of the second to fourth toes, associated with a mass over the postero-medial ankle.

On examination, Tinel's test was positive. The ankle was non-irritable, alignment was normal and there were no gross varicose veins.

Laboratory analysis demonstrated a normal C reactive protein. X-rays showed no soft-tissue calcification or bony changes. MRI demonstrated a non-enhancing 2×2.2×1.3 cm complex ganglion cyst positioned within the tarsal tunnel, with possible extension into the subtalar joint (figures 1 and 2). Ultrasound examination demonstrated venous varices in the right distal leg.

Intraoperatively, no obvious ganglion was identified near the tibial nerve. Varicosity of the posterior tibia vein was evident, extending into the tarsal tunnel. The patient was treated with decompression and neurolysis of the tibial nerve.

The patient was followed up 2 weeks after the operation and had improvement in paraesthesia symptoms.

Tarsal tunnel syndrome is caused by compression of the tibial nerve as it passes through the tarsal tunnel, underneath the flexor retinaculum at the level of the ankle joint.^{1,2} Symptoms can include pain and paraesthesia involving distributions of the tibial nerve. Extrinsic causes include trauma or deformity that can stretch or compress the tibial nerve. Intrinsic causes include space-occupying masses causing compression.² Differential diagnoses include venous structures, ganglion cysts or soft-tissue tumour such as a synovial sarcoma. Unlike a synovial sarcoma, which can appear iso/hyperintense with heterogeneity on MRI T1 imaging with contrast enhancement, the mass

Figure 2 T1 MRI axial and coronal views demonstrating a ganglion cyst positioned within the tarsal tunnel, arising from the posterior subtalar articular articulation.

Learning points

- ▶ Tarsal tunnel syndrome is caused by compression of the tibial nerve.
- ▶ MRI and ultrasound imaging are important modalities to investigate causes of tarsal tunnel syndrome.
- ▶ Venous structures can cause compression symptoms, and caution should be exercised during exploration of the tarsal tunnel and decompression of the nerve.
- ▶ In the presence of varicosities, release of the retinaculum alone may be sufficient to relieve compressive symptoms.

in our patient was hypointense on T1 imaging and hyperintense on T2 imaging without contrast enhancement.³

Contributors Literature review: RF, MN. Drafting manuscript: RF, MN, PB, JL.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Alshami AM, Babri AS, Souvlis T, *et al.* Biomechanical evaluation of two clinical tests for plantar heel pain: the dorsiflexion-eversion test for tarsal tunnel syndrome and the windlass test for plantar fasciitis. *Foot Ankle Int* 2007;28:499–505.
- 2 DiDomenico LA, Masternick EB. Anterior tarsal tunnel syndrome. *Clin Podiatr Med Surg* 2006;23:611–20.
- 3 Duran-Stanton AM, Bui-Mansfield LT. Magnetic resonance diagnosis of tarsal tunnel syndrome due to flexor digitorum accessorius longus and peroneocalcaneus internus muscles. *J Comput Assist Tomogr* 2010;34:270–2.

Figure 1 T2 MRI sagittal and coronal views suggestive of a ganglion cyst in the tarsal tunnel, arising from the subtalar joint.

© BMJ Publishing Group Limited 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Falovic R, Nambiar M, Boekel P, *et al.* *BMJ Case Rep* 2019;12:e230072. doi:10.1136/bcr-2019-230072

Copyright 2019 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow