Sudden onset dysphagia in a patient with dementia

Babatunde Oremule,¹ Elliot Heward,² Sadie Khwaja³

¹ENT, Stockport NHS Foundation Trust, Stockport, UK

²ENT, Warrington and Halton Hospitals NHS Foundation Trust, Warrington, UK

³ENT, Wythenshawe Hospital, Manchester, UK

Correspondence to

Mr Babatunde Oremule,
b.oremule@doctors.org.uk

Accepted 20 May 2019

DESCRIPTION

An 81-year-old female patient with vascular dementia attended the emergency department with sudden onset dysphagia and hearing loss. She was pleasantly confused at her baseline and therefore unable to give a history. On presentation, she was able to tolerate her own saliva but unable to tolerate sterile water, spitting it back up immediately. An upper oesophageal foreign body or food bolus was suspected and a lateral soft tissue neck X-ray was performed (figure 1). This showed a radio-opaque foreign body opposite the C5 vertebral body. Her relatives arrived and noted that she was not wearing her right in-the-ear hearing aid, and it could not be found at home. There was a high level of suspicion that she may have ingested it. She underwent a combined urgent rigid oesophagoscopy by the ear, nose and throat surgeon and flexible oesophagogastroscope by the upper gastrointestinal surgeon, as the hearing aid was slippery and needed a basket to safely retrieve it. The hearing aid was checked to confirm it was completely removed, with the button battery found to be still safely inside the device (figure 2). There were no visible visceral injuries seen on repeat endoscopy following removal of the hearing aid.

A button battery is a small, flat and cylindrical single cell battery, usually 5–25 mm in diameter. Button battery ingestion is a recognised public health concern around the world, particularly in children.^{1–3} If not removed promptly, they can lead to severe complications and death. Button batteries

Figure 2 In-the-ear hearing aid retrieved with button battery inside.

can be identified on X-rays due to the characteristic 'halo sign'.⁴ Oesophageal perforations were identified in 189 cases of fatal or severe battery ingestions with oesophageal lodgment (53 fatal, 136 severe; 95.2% in children ≤4 years).⁵ Implicated batteries were predominantly lithium (91.0%) and 92.0% were ≥20 mm diameter. Button batteries are used to power many hearing aids; hence, the concern in this case was that the button battery could slip out and cause fatal injury.

Figure 1 Lateral soft tissue neck X-ray.

Patient's perspective

We are relieved that the hearing aid was quickly and safely removed, now knowing the danger posed by swallowing button batteries (Daughter-in-law).

Learning points

- ▶ Collateral history is important in patients with dementia, clinicians should have a low threshold for performing investigations.
- ▶ Button battery ingestion is associated with severe morbidity and death, and they must be removed as a matter of emergency.

© BMJ Publishing Group Limited 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Oremule B, Heward E, Khwaja S. *BMJ Case Rep* 2019;**12**:e230300. doi:10.1136/bcr-2019-230300

The patient recovered well from her procedure and resumed a normal diet the following day. She was discharged 6 days after admission while waiting for a dementia reassessment and an improved social package of care. She was referred to audiology for a new behind-the-ear hearing aid.

Contributors BO and SK were involved in the management of the patient. BO wrote the manuscript. EH helped with literature review and reviewed the final manuscript. SK also reviewed the final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Next of kin consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Cairns R, Brown JA, Lachiredy K, *et al*. Button battery exposures in Australian children: a prospective observational study highlighting the role of poisons information centres. *Clin Toxicol* 2019;57:1–7.
- 2 Labadie M, O'Mahony E, Capaldo L, *et al*. Severity of button batteries ingestions: data from French Poison Control Centres between 1999 and 2015. *Eur J Emerg Med* 2018;25:e1–8.
- 3 Centers for Disease Control and Prevention (CDC). Injuries from batteries among children aged < 13 years—United States, 1995–2010. *MMWR. Morbidity and mortality weekly report*. 2012;61:661.
- 4 Bolton SM, Saker M, Bass LM. Button battery and magnet ingestions in the pediatric patient. *Curr Opin Pediatr* 2018;30:653–9.
- 5 Soto PH, Reid NE, Litovitz TL. Time to perforation for button batteries lodged in the esophagus. *Am J Emerg Med* 2019;37.

Copyright 2019 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow