Conjunctival lymphoma

Jenny Jing Li,¹ Weina Chen,² Navid Sadeghi^{1,3}

¹Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, Texas, USA

²Department of Pathology, University of Texas Southwestern Medical Center, Dallas, Texas, USA

³Division of Hematology and Oncology, Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, Texas, USA

Correspondence to

Dr Navid Sadeghi, navid.sadeghi@utsouthwestern.edu

Accepted 24 June 2019

DESCRIPTION

A man in his 60s with no history of autoimmune diseases or preceding infections presented with bilateral painless red eyes. For the last few weeks, the patient had noticed painless conjunctival redness in both of his eyes, left worse than right. The redness was associated with some tear production and blurry vision but no purulent discharge. Physical examination revealed diffuse, salmon-coloured, slightly raised lesions involving most of his left conjunctiva and a smaller lesion involving his lateral right conjunctiva (figure 1A). Diffuse lymphadenopathy was noted on nodal survey. Laboratory studies revealed leukocytosis ($14.46 \times 10^9/L$) with lymphocyte predominance (78%). Serum protein electrophoresis detected the presence of M-component (IgM Lambda, 0.36 g/dL). CT scans of the chest, abdomen and pelvis demonstrated lymphadenopathy involving the axilla, mediastinum, retroperitoneum and pelvis.

The patient underwent bilateral conjunctival biopsies. Pathology revealed infiltrations of small- to medium-sized lymphocytes (figure 1B) that were PAX5 (+) (figure 1C), CD3 (–) (figure 1D), CD5 (–) and CD10 (–) consistent with a diagnosis of CD5 (–)/CD10 (–) B lineage non-Hodgkin's lymphoma (NHL). Flow cytometry showed a population of variably sized (predominantly small-sized to medium-sized) cells with the following immunophenotype: CD5 (–), CD10 (–), CD19 (+), CD20 (+), CD38 (partial +), CD45 (+), kappa (–), lambda (+). Bone marrow biopsy demonstrated extensive infiltration with cells of similar morphology and immunophenotype. No chromosomal abnormalities were noted on karyotype. Molecular analysis of the myeloid

differentiation primary response 88 (MYD88) gene was negative for mutations. Based on the above results and extranodal involvement, this patient was diagnosed with stage IV marginal zone lymphoma with conjunctival involvement (conjunctival lymphoma).

Ocular adnexal lymphoma (OAL) is a rare manifestation of NHL and accounts for 1%–2% of all NHL and 5%–10% of all extranodal lymphomas.¹ About 25%–30% of OALs involve the conjunctiva.¹ Conjunctival lymphoma typically presents in the fifth to seventh decade of life as a painless salmon-pink patch in the bulbar conjunctiva, which can be misdiagnosed as conjunctivitis. Clinically, reactive lymphoid hyperplasia can also appear nearly identical to conjunctival lymphoma, so it is crucial to establish the correct diagnosis via biopsy and further studies.

Treatment of conjunctival lymphoma is determined by the stage and extent of disease. Patients with disease localised to the ocular adnexa are typically treated with external beam radiation therapy at the site of disease, which typically results in excellent disease control.^{2,3} Patients with systemic disease are typically treated similarly to those with nodal marginal zone lymphoma. First-line therapies include single-agent rituximab, rituximab and bendamustine, R-CHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone) and R-CVP (rituximab, cyclophosphamide, vincristine, prednisone).² Recurrence rates after treatment can be significant (30%–80%) and close observation is warranted.⁴ This patient was initially treated with single-agent rituximab resulting in a short-lived partial response. Patient was then treated with rituximab and bendamustine with subsequent resolution of adenopathy and conjunctival infiltration.

Figure 1 (A) Salmon-pink patches involving most of the conjunctiva. (B) Infiltrations of small- to medium-sized cells (H&E, 500×) on conjunctival biopsy. (C) Lymphoma cells expressing PAX5 (IHC, 100×). (D) Lymphoma cells lacking CD3 (IHC, 100×).

Learning points

- ▶ Bilateral eye redness is not always conjunctivitis; salmon-coloured patches should raise suspicion for conjunctival lymphoma.
- ▶ Clinical features of conjunctival lymphoma are non-specific and the lesions can appear nearly identical to other more benign processes, so it is crucial to obtain biopsies and further studies.
- ▶ Disease localised to the ocular adnexa can be treated with external beam radiation therapy, but patients with systemic disease may need chemotherapy.

Contributors JJJ and NS were involved in the conception, logistics/planning (obtained informed consent, image acquisition with medical images), drafting the article and revising it critically for important intellectual content. WC was involved in the acquisition

© BMJ Publishing Group Limited 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Li JJ, Chen W, Sadeghi N. *BMJ Case Rep* 2019;**12**:e229599. doi:10.1136/bcr-2019-229599

and interpretation of pathological images. JJL, WC and NS gave final approval of the version published.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Sjö LD. Ophthalmic lymphoma: epidemiology and pathogenesis. *Acta Ophthalmol* 2009;87 Thesis 1(Thesis 1):1–20.
- 2 National Comprehensive Cancer Network. B-cell Lymphomas (Version 2). 2018. https://www.nccn.org/professionals/physician_gls/pdf/b-cell.pdf
- 3 Harada K, Murakami N, Kitaguchi M, *et al.* Localized ocular adnexal mucosa-associated lymphoid tissue lymphoma treated with radiation therapy: a long-term outcome in 86 patients with 104 treated eyes. *Int J Radiat Oncol Biol Phys* 2014;88:650–4.
- 4 Kirkegaard MM, Rasmussen PK, Coupland SE, *et al.* Conjunctival lymphoma—an international multicenter retrospective study. *JAMA Ophthalmol* 2016;134:406–14.

Copyright 2019 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/> BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow