Pigmented iris cyst in vitreous chamber

Nimesh Patel,¹ Rajeev Reddy Pappuru,¹ Soumyava Basu,² Mudit Tyagi^{1,2}

¹Smt Kanuri Santhamma Center for Vitreoretinal Diseases, LV Prasad Eye Institute, Hyderabad, Telangana, India

²Uveitis and Ocular Immunology Services, LV Prasad Eye Institute, Hyderabad, India

Correspondence to

Dr Mudit Tyagi;
drnudittyagi@gmail.com

Accepted 22 October 2020

DESCRIPTION

A 45-year-old male patient, diagnosed elsewhere as retinal detachment, was seen in the retina services of our institute. There was no history of prior ocular trauma or surgery. His vision was 20/20 in his right eye and 20/125 in the left eye. There were no signs of any ocular inflammation and the intra-ocular pressure was within normal limit. A retinal evaluation revealed a free-floating pigmented cyst in vitreous cavity (figure 1, white arrow). A pars plana vitrectomy was done and the cyst was removed. The histopathology was suggestive of iris pigment cells. The pigmented cells on the cyst wall are more likely to originate from the ciliary pigment epithelium. The cysts initially are formed on the ciliary body and then get dislodged into the vitreous, giving rise to the abrupt symptom of a diminution of vision.

Vitreous cysts are rare ocular malformations, which can be both pigmented and non-pigmented cysts.¹ Non-pigmented congenital cysts are believed to be remnants of the hyaloid vascular system such as Bergmeister's papilla and Mittendorf's dot,² while pigmented congenital cyst has been reported to be choristomas of the hyaloid vascular system.³⁻⁴ Acquired cysts have been reported to occur after ocular trauma or as a complication of ocular inflammation and uveitis.⁴ They have also been associated with uveal colobomas, retinitis pigmentosa and retinal detachment surgeries.⁵⁻⁷ Shields had postulated that epithelial cysts arise between the pigmented epithelial layers of the iris and often break free and migrate into the vitreous chamber and these are called dislodged cysts. These dislodged cysts can come in visual axis and cause a sudden diminution of vision.⁸

Figure 1 Pigmented iris cyst in vitreous chamber.

Patient's perspective

I had a black shadow moving in my eye, and after the surgery, my vision is more clear and the shadow has gone.

Learning points

- ▶ Congenital vitreous cysts may be remnants of the hyaloid vascular system such as Bergmeister's papilla and Mittendorf's dot or may represent choristoma of the hyaloid vascular system.
- ▶ Acquired cysts may be associated with trauma, uveitis, uveal colobomas and retinal detachment surgeries.
- ▶ Pigmented epithelial may also arise between the pigmented epithelial layers of the iris and can often break free and migrate into the vitreous chamber.

Contributors NP and MT wrote the manuscript and edited the image. NP conceptualised the case report. SB and RRP managed the patient. MT acquired the image and helped in literature search. MT is the corresponding author and assumes responsibility for the case report.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Cruciani F, Santino G, Salandri AG. Monolateral idiopathic cyst of the vitreous. *Acta Ophthalmol Scand* 1999;77:601–3.
- 2 Francois J. Pre-papillary cyst developed from remnants of the hyaloid artery. *Br J Ophthalmol* 1950;34:365–8.
- 3 Nork TM, Millecchia LL. Treatment and histopathology of a congenital vitreous cyst. *Ophthalmology* 1998;105:825–30.
- 4 Tranos PG, Ferrante P, Pavesio C. Posterior vitreous cyst and intermediate uveitis. *Eye* 2010;24:1115–6.
- 5 Lavric A, Urbancic M. Floating vitreous cyst: two clinical cases. *Case Rep Ophthalmol* 2013;4:243–7.
- 6 Tuncer S, Bayramoglu S. Pigmented free-floating vitreous cyst in a patient with high myopia and uveal coloboma simulating choroidal melanoma. *Ophthalmic Surg Lasers Imaging* 2011.
- 7 Ludwig CA, Leng T. Idiopathic pigmented vitreous cyst. *Acta Ophthalmol* 2016;94:e83–4.
- 8 Shields JA. Primary cysts of the iris. *Trans Am Ophthalmol Soc* 1981;79:771.

© BMJ Publishing Group Limited 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Patel N, Pappuru RR, Basu S, et al. *BMJ Case Rep* 2020;13:e239431. doi:10.1136/bcr-2020-239431

Copyright 2020 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow