Iatrogenic endometriosis due to isthmic stenosis post-trachelectomy

Pavitra Nanayakkara,^{1,2} Alex Ades^{1,2,3}

¹Agora Specialist Centre, Epworth Hospital, Richmond, Victoria, Australia

²Department of Obstetrics and Gynaecology, Royal Women's Hospital, Parkville, Victoria, Australia

³Department of Obstetrics and Gynaecology, University of Melbourne, Melbourne, Victoria, Australia

Correspondence to

Dr Pavitra Nanayakkara;
pavitra.nanayakkara@
thewomens.org.au

Accepted 4 February 2020

DESCRIPTION

A nulliparous 42-year-old woman presented for a prophylactic pre-pregnancy laparoscopic transabdominal cerclage (suture around the cervicoisthmic junction to prevent pregnancy loss due to cervical insufficiency) after having undergone a trachelectomy for stage IA cervical cancer 10 years ago. The patient had a laparoscopy for investigation of pelvic pain a couple of years before the trachelectomy. Mild superficial endometriosis was reported and treated at the time. She had been having regular menstrual cycles until the trachelectomy and became permanently amenorrhoeic postsurgery. Blood tests done prior to the procedure showed normal ovarian and thyroid functions. She did not undergo chemotherapy or radiotherapy that could have contributed to reduction in ovarian activity and hence cessation of menses. She has recently started In-vitro fertilisation (IVF) treatment for infertility attributed to anatomical factors (stenosed cervix). All hormonal investigations came back within normal range, again confirming adequate ovarian function. There was no transvaginal access to the endometrial cavity due to cervical stenosis.

The laparoscopic transabdominal cerclage was placed successfully; however, significant endometriosis was evident during the laparoscopy, including several lesions over the bowel (see figures 1 and 2). We believe that the more extensive endometriosis seen at the time of the transabdominal cerclage developed some time after the trachelectomy and supports the aetiological theory of retrograde menstruation as a contributor to endometriosis. It is not possible to determine exactly when it happened within the 10 years that passed; however, the retrograde menstruation caused by the blocked cervix seems to have been a major contributor. Acquired isthmic stenosis causing backflow of menstrual blood through the fallopian tubes has been suggested in small studies

Figure 2 Images without labels.

and demonstrated on magnetic resonance imaging (MRI), theoretically increasing predisposition to endometriosis.^{1–5} This case provides a clinical presentation of subsequent endometriosis. We have not found evidence in the literature definitively demonstrating this until now.

At present, there seems to be no evidence as to whether iatrogenic endometriosis behaves or progresses differently from the non-iatrogenic form of the disease. Hence, the management of iatrogenic endometriosis should be as for the non-iatrogenic disease. Indications for treating the condition are based on the clinical presentation and associated symptoms, mainly dysmenorrhoea, dysuria, dyschezia, chronic pain and subfertility. The same medical treatment for control of symptoms and surgical treatment for excision of endometriotic lesions should be considered. The authors are investigating measures such as insertion of a levonorgestrel intrauterine device after trachelectomy to prevent endometriosis.

Figure 1 Labelled images.

© BMJ Publishing Group Limited 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Nanayakkara P, Ades A. *BMJ Case Rep* 2020;**13**:e233579. doi:10.1136/bcr-2019-233579

Learning points

- Endometriosis can occur as a result of retrograde menstruation secondary to outflow tract obstruction.
- It is important to consider the possibility of endometriosis in women who have undergone procedures that may result in cervical or isthmic stenosis; preoperative counselling of the risk of endometriosis is suggested.

Contributors PN photographed and labelled the images, wrote the first draft of the manuscript, amended the article in consultation with AA, submitted the case summary and made changes as per *BMJ Case Reports* recommendations. AA performed the operation, reviewed and edited the summary, and approved the final submission.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Barbieri RL. Stenosis of the external cervical os: an association with endometriosis in women with chronic pelvic pain. *Fertil Steril* 1998;70:571–3.
- 2 Barbieri RL, Callery M, Perez SE. Directionality of menstrual flow: cervical os diameter as a determinant of retrograde menstruation. *Fertil Steril* 1992;57:727–30.
- 3 Sahdev A, Jones J, Shepherd JH, *et al.* Mr imaging appearances of the female pelvis after trachelectomy. *Radiographics* 2005;25:41–52.
- 4 Li X, Li J, Ju X, *et al.* Menstrual pattern after abdominal radical trachelectomy. *Oncotarget* 2017;8:53146–53.
- 5 Selo-Ojeme DO, Ind T, Shepherd JH. Isthmic stenosis following radical trachelectomy. *J Obstet Gynaecol* 2002;22:327–8.

Copyright 2020 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/> BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow