Sand injuries associated with the landslide

Masako Hata,¹ Toshihisa Ichiba,¹ Yuji Okazaki²

¹Department of Emergency Medicine, Hiroshima City Hiroshima Citizens Hospital, Hiroshima, Japan

²Department of Internal Medicine, Akiota Hospital, Yamagata-gun, Hiroshima, Japan

Correspondence to
Dr Yuji Okazaki;
ur8nibhc1@gmail.com

Accepted 14 April 2020

DESCRIPTION

A 41-year-old woman was involved in a landslide caused by heavy rainfall in July 2018 in Hiroshima City, Hiroshima, Japan. She was trapped in her car and was drifted away by the landslide. She was soaked in earth and sand up to her face, but she managed to escape by herself from her car. She then visited our emergency department complaining of whole body bruising and abrasions. On examination, her blood pressure was 123/82 mm Hg, heart rate was 86 beats per minute and body temperature was 38.8 degrees. Her hip was severely contaminated by sand. After washing her hip injury, we performed whole body CT. CT findings of her trunk showed a skin defect and hyperdense material indicating sand around her left hip and penetration of sand and air into the muscle (figure 1A). Moreover, sand and air extended upward subcutaneously (figure 1B). CT findings also showed sand in the stomach but not in the lung. CT findings of the head showed fluid and sand in the maxillary sinus and sphenoidal sinus (figure 2A) and sand in the external auditory canal and fluid in the mastoid sinus (figure 2B). Antimicrobial therapy was started, and a wide range of debridement was performed for sand injury of the trunk, during which yellow-coloured purulent material and sand were removed. After healing of the trunk injury, she complained of persistent fever and sinus discomfort, and endoscopic irrigation was therefore performed for the maxillary sinus and sphenoidal sinus under general anaesthesia. She was discharged without any complications such as olfactory disturbance.

Sand injuries associated with a landslide are rare but are important and potentially life-threatening traumatic conditions. As in a tsunami disaster, a

Figure 2 (A) CT findings of the head showed fluid and sand in the maxillary sinus and sphenoidal sinus (arrowhead). (B) CT findings of the head showed sand in the external auditory canal (arrowhead) and fluid in the mastoid sinus (white arrowhead).

Learning points

- ▶ Sand injuries associated with a landslide are rare but are important and potentially life-threatening traumatic conditions.
- ▶ Sand can spread from the body surface into subcutaneous tissue and inside the body and into various luminal organs.
- ▶ Whole body CT and careful evaluation should be performed in patients who have been injured in a landslide.

landslide can cause various injuries over the whole body, and diagnosis might be delayed because of multiple and other severe injuries.^{1 2} Sand can spread from the body surface into subcutaneous tissue and inside the body and into various luminal organs. Therefore, whole body CT and careful evaluation should be performed in patients who have been injured in a landslide.

Contributors All authors contributed to the development of this manuscript. MH was responsible for literature search and writing of all manuscript. YO and TI were supervisors.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Kaewlai R, Srisuwan T, Prasitvoranant W, *et al*. Radiologic findings in tsunami trauma: experience with 225 patients injured in the 2004 tsunami. *Emerg Radiol* 2007;14:395–402.
- 2 Pettiti N. Images in clinical medicine. sand as a foreign body. *N Engl J Med* 1997;337:96.

Figure 1 (A) CT findings of the trunk showed a skin defect and hyperdense material indicating sand around her left hip (arrowhead) and penetration of air (white arrowhead) and sand into the muscle. (B) CT findings of the trunk showed sand (arrowhead) and air (white arrowhead) that extended upwards from the hip subcutaneously.

© BMJ Publishing Group Limited 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Hata M, Ichiba T, Okazaki Y. *BMJ Case Rep* 2020;13:e235254. doi:10.1136/bcr-2020-235254

Copyright 2020 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow