Submental intubation is a viable alternative to tracheostomy in facial trauma

Inês Queixinho Martins,¹ Mafalda Castro,¹ Pedro Ferraz,² Inês Simões¹

¹Anesthesiology, Centro Hospitalar e Universitário de Coimbra EPE, Coimbra, Portugal
²Stomatology, Centro Hospitalar e Universitário de Coimbra EPE, Coimbra, Portugal

Correspondence to
 Dr Inês Queixinho Martins;
 ines.iqm@gmail.com

Accepted 21 April 2020

DESCRIPTION

A 38-year-old man victim of a high impact motor-cycle accident with cranial and facial trauma (bilateral type III Le Fort) was proposed for an open reduction and osteosynthesis of his lesion. Due to the extensive existing fracture, the preferred airway approach would be through a tracheostomy or a submental intubation. Although tracheostomy is the most used technique, it is associated with more complications such as haemorrhage, surgical emphysema, laryngeal recurrent nerve injury, tracheal stenosis, tracheoesophageal fistula and dysphagia. Submental intubation has a lower complication rate and can be performed in patients with type II and III Le Fort fractures so, it was chosen, and a balanced general anaesthesia was associated.^{1,2}

After anaesthetic induction with fentanyl and propofol, succinylcholine was used as a muscle relaxant and orotracheal intubation was performed with a reinforced tube to prevent kinking from the angle of submental insertion. Afterwards, in cooperation with the surgical team, a 2-centimetre submental incision was performed. Initially curved forceps were introduced into the incision, passing through the subcutaneous layer, platysma, mylohyoid muscle and into the sublingual space. Afterwards, they were passed through the mucosal layers, lateral to the sublingual ducts, avoiding the sublingual gland. The superior part of the orotracheal tube was then inserted through the previously created incision and emerged in the submental region (figure 1).

Figure 1 Orotracheal tube emerging from the submental region.

Figure 2 Submental incision, 1 month after surgery.

Surgery progressed without any complications, the airway tube was placed in its initial position and the submental region was sutured successfully as well. The patient's wake-up proceeded calmly without any nausea or pain associated. In the immediate postoperative care, extubation was performed with the patient already awake. The patient was discharged 9 days after the surgery without any complication during his stay at the ward and returned for a follow-up appointment 1 month later, with a great recovery, with minimal scarring (figure 2).

Airway management in facial injuries is difficult and often presents a challenge as oral route makes the surgery impossible and nasal route is contraindicated due to the facial and cranial base fractures.³ Tracheostomy is the most conventional technique in these cases, but it is more invasive and can lead to multiple complications, thus lengthening hospital

Learning points

- ▶ Tracheostomy is associated with multiple complications (surgical emphysema, tracheal erosion, haemorrhage, dysphagia, problems with decannulation and excessive scarring at the exit site) that do not occur with submental intubation.
- ▶ Submental intubation is an easy technique performed by an anaesthesiologist through an orotracheal intubation and a simple surgical procedure.
- ▶ Although it is an old technique, submental intubation should not be disused and can be considered in select cases due to its reduced risk, scarring and consequent shorter hospital stay.

© BMJ Publishing Group Limited 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Martins IQ, Castro M, Ferraz P, et al. *BMJ Case Rep* 2020;**13**:e235537. doi:10.1136/bcr-2020-235537

stay. Submental intubation, on the other hand, is a simple technique, with a low complication risk because of the lack of large blood vessels and nerves in its tract.² In this case we present a successful submental intubation instead of the usual tracheostomy with a great postoperative outcome. This technique, created in 1986, although underused, should be considered more often in these patients' airway approach.¹

Contributors IQM wrote the article, took the pictures and explained the informed consent to the patient was the anaesthesiologist following the patient's case. MC helped writing the case, provided critical feedback and followed the patient's case as well PF helped taking the pictures, corrected the case and translated IS provided critical feedback on the case, was the senior anaesthesiologist and corrected the final version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Amin M, Dill-Russell P, Manisali M, *et al*. Facial fractures and submental tracheal intubation. *Anaesthesia* 2002;57:1195–9.
- 2 Chandu A, Smith AC, Gebert R. Submental intubation: an alternative to short-term tracheostomy. *Anaesth Intensive Care* 2000;28:193–5.
- 3 Das S, Das TP, Ghosh PS. Submental intubation: a journey over the last 25 years. *J Anaesthesiol Clin Pharmacol* 2012;28:291–303.

Copyright 2020 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow