Case report

Rhabdomyolysis as an initial presentation in a patient diagnosed with COVID-19

Benjamin Valente-Acosta ¹, Francisco Moreno-Sanchez,¹ Omar Fueyo-Rodriguez,¹ Andres Palomar-Lever²¹Internal Medicine and Infectious Diseases, Centro Medico ABC, Ciudad de México, Mexico City, Mexico
²Pulmonary Medicine, Centro Medico ABC, Mexico City, Mexico

Correspondence to

Dr Benjamin Valente-Acosta; benjamin.valente-acosta1@alumni.lshrm.ac.uk

BV-A and FM-S are joint first authors.

Accepted 12 June 2020

SUMMARY

The presence of rhabdomyolysis secondary to multiple infections has been reported, predominantly viral, but also bacterial and fungal. It is well known that COVID-19 can present a wide variety of complications during the course of infection; however, the presence of rhabdomyolysis as an initial condition has not been reported so far. We report a case of rhabdomyolysis as an initial presentation in a patient diagnosed with SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) infection.

BACKGROUND

Rhabdomyolysis is defined as a dissolution of skeletal muscle, with acute kidney injury (AKI) as the most important and comorbid complication, especially in severe cases. Patients with SARS-CoV-2 infection can have a wide variety of complications during the course of infection. Pulmonary, cardiovascular and thrombotic complications have been widely reported. The present case reports a patient diagnosed with SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) infection presenting with rhabdomyolysis.

CASE PRESENTATION

A 71-year-old man, who had a history of benign prostatic hyperplasia and of smoking 20 cigarettes a day for the past 30 years, presented to the emergency department with a 1-week history of dry coughing, mild dyspnoea and a fever of 38°C that did not resolve with paracetamol. Two weeks previously, he had returned to Mexico City from Miami. On the day of his admission, he felt greater dyspnoea and was suffering from severe myalgia and arthralgia, predominantly in his legs.

At his initial evaluation, the patient reported generalised weakness and malaise. He did not report any nausea, vomiting, diarrhoea, urinary or neurological symptoms. He had not taken any medication except paracetamol (750 mg) every 8 hours to control the fever. He had no known allergies.

A physical examination revealed a blood pressure of 120/68 mmHg, a pulse of 90 beats/min and a respiratory rate of 22 breaths/min. His temperature was 36.5°C and his oxygen saturation was 84% while breathing ambient air. He was alert and oriented, but his speech was slow and pausing. A cardiopulmonary examination revealed rales in both lung bases, but without signs of respiratory

distress. Otherwise, his physical examination was unremarkable.

INVESTIGATIONS

Blood tests showed a normal leucocyte count with lymphopaenia ($0.85 \times 10^9/L$), haemoglobin level (161 g/L) and mild thrombocytopaenia ($118\,000 \times 10^9/L$). His creatinine level was increased, at 1.68 mg/dL, and his C-reactive protein (CRP) and procalcitonin (PCT) were elevated, at 2.9 mg/dL and 2.89 ng/mL, respectively. His interleukin (IL)-6 was also increased (233 pg/mL). His muscle enzymes were markedly elevated (creatinine phosphokinase at 8720 U/L and myoglobin at 2079 ng/mL). His ferritin and lactic dehydrogenase levels were high as well (at 2603 ng/mL and 541 U/L, respectively). A urinalysis showed haemoglobin without erythrocytes in the sediment microscopy (table 1). A nasopharyngeal swab for SARS-CoV2 was positive. Multiplex PCR for respiratory viruses and HIV test were negative. A lung CT scan showed bilateral infiltrates with areas of consolidation and extensive ground-glass opacities.

TREATMENT

The patient was admitted to the COVID-19 ward and administered oxygen by nasal cannula. We started aggressive fluid and bicarbonate therapy as well as enoxaparin, azithromycin and ceftriaxone. The patient agreed to the use of compassionate drug therapy, so we started him on hydroxychloroquine and lopinavir/ritonavir. His condition deteriorated on the third day of hospitalisation and he required invasive mechanical ventilation. Over the next 2 days, his condition continued to deteriorate, with fever, hypotension and high ventilatory requirements. His CRP and IL-6 levels also increased (to 26.9 mg/dL and 275 pg/mL, respectively). His family agreed to the use of tocilizumab on a compassionate use scheme. Consequently, we gave him two doses of tocilizumab 400 mg intravenous on the fifth and sixth day after admission.

OUTCOME AND FOLLOW-UP

On the seventh day after admission, his condition started to improve and he showed a decrease in lactic dehydrogenase, ferritin, CRP and PCT levels. It was possible to extubate him on the 12th day after admission. His clinical condition continued to improve and after a negative SARS-CoV2 test, we

© BMJ Publishing Group Limited 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Valente-Acosta B, Moreno-Sanchez F, Fueyo-Rodriguez O, et al. *BMJ Case Rep* 2020;**13**:e236719. doi:10.1136/bcr-2020-236719

Unusual association of diseases/symptoms

Table 1 Laboratory data

Variable	On admission	5th day	8th day	16th day	Reference range
Haemoglobin (g/L)	161	126	105	115	145–185
White cell count ($\times 10^9/L$)	7.1	6.2	4.5	4.2	1.8–10.0
Lymphocytes	0.85	1.4	0.6	2.43	1.00–3.50
Platelet count ($\times 10^9/L$)	118 000	137 000	218 000	280 000	150 000–45 000
Creatinine (mg/dL)	1.68	0.99	0.70	0.84	0.74–1.35
Electrolytes					
Sodium (mmol/L)	137	137	147	141	135–145
Potassium (mmol/L)	4	3.9	4.9	4.2	3.5–5.1
Carbon dioxide (mmol/L)	19.9	22.2	32	23.1	22–29
D-dimer (ng/mL)	983	945	656		40–500
Lactic dehydrogenase (U/L)	541	672	294		122–22
Creatin kinase (U/L)	8720	3876	460	71	39–308
Myoglobin (ng/mL)	2079	208			28–72
C-reactive protein (mg/dL)	2.99	26.9	4.2	0.39	0.00–0.50
Procalcitonin (ng/mL)	2.89	4.7	1.45	0.17	0.00–0.50
Ferritin (ng/mL)	2603	4073	1938	1544	30–400
IL-6 (pg/mL)	233	275			0.0–7.0

IL, interleukin.

were able to finally discharge him to his home on the 16th day after admission.

DISCUSSION

This case illustrates that rhabdomyolysis could be related to SARS-CoV2 infection and could be a presenting problem in patients with COVID-19 severe pneumonia, rather than being only a late complication, as was previously reported.¹ In a large series of COVID-19 patients, Guan *et al* reported two cases of rhabdomyolysis in non-severe cases. However, the study does not clarify specificities about the patients.² Likewise, Suwanwongse *et al* described a case of a patient with non-severe COVID-19 pneumonia with rhabdomyolysis as a presenting feature and Gefen *et al* described the first paediatric patient with rhabdomyolysis and non-severe COVID-19 infection.^{3,4}

Rhabdomyolysis has been associated with viral infections and especially influenza.⁵ It has also been reported in association with SARS.⁶ Our patient presented with clinical and biochemical evidence of rhabdomyolysis before he was started on any drug or had been placed on paralytic therapy for mechanical ventilation, which are known causes of muscular injury.⁷

A recent report of renal histopathological features in post-mortem COVID-19 patients found pigmented cast in three cases; the authors stated that drug-relevant or hyperventilation-relevant rhabdomyolysis contributed, although they did not rule out a possible direct viral injury on muscle.⁸ However, SARS-CoV-2 has been isolated in multiple tissues as kidneys, liver, brain and heart, which could suggest that the virus could also infect striated muscle tissue.⁹ In our case, the patient also presented with AKI possibly associated with rhabdomyolysis because the urinalysis had haemoglobin without red blood cells in the sediment.

Although high PCT is associated with severe bacterial infections,¹⁰ it has also been related to other conditions as pancreatitis, burn injury, mechanical trauma and rhabdomyolysis.¹¹ Our patient presented with a high serum PCT without a proven bacterial infection. It is possible that the PCT level was associated with the COVID-19 infection severity and the rhabdomyolysis.¹² Luckily, our patient responded well, and although tocilizumab

use is anecdotic, the patient's temporal sequence could suggest a possible positive role.

Learning points

- Rhabdomyolysis can be the initial presentation of COVID-19.
- High creatine kinase level could be related to rhabdomyolysis and acute kidney injury, which requires an aggressive treatment to prevent further complications.
- Rhabdomyolysis and COVID-19 infection can be associated with an increased procalcitonin level without a bacterial infection.

Twitter Francisco Moreno-Sanchez @drpacomoreno1

Acknowledgements We acknowledge the residents and staff members of the ABC Medical Center, who took care of the patient.

Contributors BV-A and OF-R conceived the idea and design of the article. BV-A, FM-S, OF-R and AP-L contributed to the preparation and review of the initial manuscript. All authors approved the final version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

ORCID iD

Benjamin Valente-Acosta <http://orcid.org/0000-0001-9885-3594>

REFERENCES

- 1 Jin M, Tong Q. Rhabdomyolysis as potential late complication associated with COVID-19. *Emerg Infect Dis* 2020;26:1–3.
- 2 Guan W-J, Z-Y N, Hu Y, *et al*. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020.
- 3 Suwanwongse K, Shabarek N. Rhabdomyolysis as a presentation of 2019 novel coronavirus disease. *Cureus* 2020;12:e7561.

- 4 Gefen AM, Palumbo N, Nathan SK, *et al.* Pediatric COVID-19-associated rhabdomyolysis: a case report. *Pediatr Nephrol* 2020;23:10–13.
- 5 Runnstrom M, Ebied AM, Khoury AP, *et al.* Influenza-Induced rhabdomyolysis. *BMJ Case Rep* 2018;11:1–3.
- 6 Chen L-L, Hsu C-W, Tian Y-C, *et al.* Rhabdomyolysis associated with acute renal failure in patients with severe acute respiratory syndrome. *Int J Clin Pract* 2005;59:1162–6.
- 7 Hirano M, Ott BR, Raps EC, *et al.* Acute quadriplegic myopathy: a complication of treatment with steroids, nondepolarizing blocking agents, or both. *Neurology* 1992;42:2082–7.
- 8 Su H, Yang M, Wan C, *et al.* Renal histopathological analysis of 26 postmortem findings of patients with COVID-19 in China. *Kidney Int* 2020. doi:10.1016/j.kint.2020.04.003. [Epub ahead of print: 09 Apr 2020].
- 9 Puelles VG, Lütgehetmann M, Lindenmeyer MT, *et al.* Multiorgan and renal tropism of SARS-CoV-2. *N Engl J Med* 2020:NEJMc2011400.
- 10 Becker KL, Snider R, Nylén ES. Procalcitonin assay in systemic inflammation, infection, and sepsis: clinical utility and limitations. *Crit Care Med* 2008;36:941–52.
- 11 Meisner M. Update on procalcitonin measurements. *Ann Lab Med* 2014;34:263–73.
- 12 Zheng Z, Peng F, Xu B, *et al.* Risk factors of critical and mortal COVID-19 cases: a systematic literature review and meta-analysis. *J Infect* 2020.

Copyright 2020 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow