Germ cell tumours: a scary presentation

Carlos Andre Barroso Oliveira , Ana Catarina Costa , Vasco Mendes

Department of Radiology,
Hospital de Braga, Braga,
Portugal

Correspondence to
Dr Carlos Andre Barroso
Oliveira;
carlosaboliveira2014@gmail.
com

Accepted 12 June 2020

DESCRIPTION

A 32-year-old patient presented with dyspnoea, fatigue and haemoptysis in the emergency room.

Thoracic radiography was performed and demonstrated a large mass in the left lung. CT was also performed and demonstrated a large heterogeneous mass with cystic component measuring $24 \times 16 \times 14$ cm.

CT showed the anteroposterior diameter allowing depth of lesion approach and great vessels deviation (figure 1).

Coronal view (figure 2) and sagittal view (figure 3) allowed better delineation of compression and longitudinal view of the lesion as well as viable lung. These views, specially sagittal (figure 3), allow a more accurate view of bronchi compression.

Biopsy was performed and revealed a mixed germ cell tumour combined with embryonal carcinoma, yolk sac and teratoma.

Tumour markers such as LDH, AFP and β -hCG were elevated (LDH 700 U/mL, AFP 1.4 ng/mL and β -hCG 400 mIU/mL).

The patient was treated with a cisplatin-based chemotherapeutic regimen (four cycles of bleomycin, etoposide and cisplatin). Chemotherapy allowed marked reduction of the tumour. Next, the pneumonectomy was performed and histology confirmed a non-seminomatous mixed germ cell tumour (85% embryonal carcinoma, 10% yolk sac and 5% teratoma).

However, the patient is extremely debilitated physically, and the prognosis remains uncertain. During the last appointment, tumour markers were normal.

The mediastinum is centrally located in the thorax and divided into anterior, middle and posterior compartments.

Figure 1 Germ cell tumour. Portal-phase contrast-enhanced CT (axial view) shows an heterogeneous giant mass occupying most of the left lung measuring $24 \times 16 \times 14$ cm. Note the heterogeneous enhancement of the mass.

Figure 2 Germ cell tumour. Portal-phase contrast-enhanced CT (sagittal view) heterogeneous giant mass occupying most of the left lung measuring $24 \times 16 \times 14$ cm.

Germ cell tumours comprise 15%–20% of all anterior mediastinal masses, occurs in young adults in their second to fourth decade of life, with equal sex distribution. The majority are benign, with teratoma being the most frequently type seen.^{1 2}

Benign teratomas are classified into the following three groups: epidermoid cysts, dermoid cysts and teratomas. Approximately 60% of teratomas are asymptomatic.^{1–3} The probability of malignancy increases with immature type.

Seminomas are the most common malignant mediastinal germ cell tumour and are located in the anterior and superior mediastinum.^{1 2} Malignant lesions can present with pain caused by local invasion.^{3 4}

Figure 3 Germ cell tumour. Portal-phase contrast-enhanced CT (coronal view) shows an heterogeneous giant mass occupying most of the left lung and that causes right deviation of the mediastinum. The mass respects the large vessels margins without invading them.

© BMJ Publishing Group
Limited 2020. No commercial
re-use. See rights and
permissions. Published by BMJ.

To cite: Oliveira CAB,
Costa AC, Mendes V. *BMJ
Case Rep* 2020;**13**:e234522.
doi:10.1136/bcr-2020-
234522

Images in...

Non-seminomatous germ cell tumours are a relative rare group and present a more aggressive behaviour. Approximately 80% of patients developing these tumours are men. Serum β -hCG is elevated in approximately 60% of patients with non-seminomatous germ cell tumours. Elevated alpha-fetoprotein levels are correlated with the presence of embryonic or yolk sac component.^{1 2}

CT usually shows a heterogeneous mass with cystic component, often infiltrative with obliteration of fat planes.^{1 2 4}

Patient's perspective

I feel scared for the future. I am very young and afraid to die.

Learning points

- ▶ Contrast-enhanced CT and biopsy play a central role to exclude malignancy of a mediastinal mass.
- ▶ Never forget germ cell tumour when a giant mediastinal mass is found.
- ▶ The risk of malignancy increases in immature types of germ cell tumours.

Contributors CABO reported the case, selected the images and also talked with a special team for the English language and image review. ACC asked the patient for consent to publish, helped reporting the CT and image improvement and selection and also helped with planning. VM supervised all the work and defined the plan to the study as well as analysis of the data.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iDs

Carlos Andre Barroso Oliveira <http://orcid.org/0000-0002-7553-7422>

Ana Catarina Costa <http://orcid.org/0000-0002-1071-3608>

REFERENCES

- 1 Fritzsch FR, Kristiansen G, Frauenfelder T, *et al.* Large mixed germ cell tumor in a young patient presenting as an intrapulmonary mass. *Pathol Res Pract* 2009;205:572–8.
- 2 Motzer RJ, Agarwal N, Beard C, *et al.* Testicular cancer. *J Natl Compr Canc Netw* 2012;10:502–35.
- 3 Lee KS, Im JG, Han CH, *et al.* Malignant primary germ cell tumors of the mediastinum: CT features. *AJR Am J Roentgenol* 1989;153:947–51.
- 4 Ronchi A, Cozzolino I, Montella M, *et al.* Extragonadal germ cell tumors: not just a matter of location. A review about clinical, molecular and pathological features. *Cancer Med* 2019;8:6832–40.

Copyright 2020 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow