OPEN ACCESS

Tick-borne relapsing fever

Emily Scrivens Aviles,¹ Michael Oakes,² Meg Algranati,² André Martin Mansoor³

¹School of Medicine, Oregon Health and Science University, Portland, Oregon, United States

²Internal Medicine Residency Program, Oregon Health and Science University, Portland, Oregon, United States

³Department of Medicine, Oregon Health and Science University, Portland, Oregon, United States

Correspondence to

Dr André Martin Mansoor;
mansooran@ohsu.edu

Accepted 1 July 2020

DESCRIPTION

A previously healthy 58-year-old man presented to an emergency department with a 3-day history of intermittent drenching sweats with rigours, diarrhoea, fatigue and decreased oral intake. These symptoms began a week after the patient returned from a summer vacation to a rural community in Central Oregon. On examination, temperature was 37.3°C, heart rate was 74 beats per minute and blood pressure was 88/56 mm Hg. The patient appeared diaphoretic. Laboratory evaluation was notable for a peripheral platelet count of 26 K/cu mm, white blood cell count of 6.38 K/cu mm, serum creatinine of 2.11 mg/dL, aspartate aminotransferase of 101 U/L and alanine aminotransferase of 102 U/L. A peripheral blood smear revealed the presence of multiple corkscrew-shaped organisms (figure 1).

The constellation of symptoms and pattern of laboratory findings was consistent with tick-borne relapsing fever (TBRF), however the patient never noticed or removed a tick during his vacation.¹ The patient was started on oral doxycycline 100 mg two times per day for a planned 10-day course. Within 2 hours of the first dose of doxycycline, the patient became delirious, tachycardic and diaphoretic, with a fever of 40.2°C, consistent with a Jarisch-Herxheimer reaction, an inflammatory response to antibiotic treatment of a spirochaetal infection. The pathogenesis is hypothesised to be due to lipoproteins that are released with lysis of spirochaetes and the ensuing cytokine stimulation.² The patient subsequently improved with supportive therapies and antipyretics. Two-tiered serologic testing via immunoassay and IgM western blot later returned positive for *Borrelia hermsii*, confirming the diagnosis of TBRF. Other spirochaetal infections were considered, including *Treponema pallidum* and *Leptospira*. However, those organisms are too small to be visualised by light microscopy. The presence

Learning points

- ▶ Consider tick-borne relapsing fever in the differential diagnosis for patients who present with an undulating febrile illness, compatible travel history and characteristic laboratory pattern.
- ▶ When corkscrew-shaped organisms are visible, light microscopy is a useful aid in differentiating tick-borne relapsing fever from other spirochaetal infections.
- ▶ The Jarisch-Herxheimer reaction is characterised by the onset of fever, rigours and hypotension 1 to 4 hours after starting antibiotic therapy in patients with a spirochaetal infection.

of corkscrew-shaped organisms on peripheral blood smear is pathognomonic for *Borrelia* infection and can confirm the diagnosis before the return of serologic testing.³

TBRF in the Northwestern region of North America is most commonly due to *B. hermsii*, a spirochaete transmitted from the rodent-associated *Ornithodoros hermsi* tick through a painless bite, often occurring at night when it is known to feed.⁴ TBRF typically presents after a week-long incubation period with a pattern of undulating fevers for 3 days, followed by a week without symptoms.⁵ *B. hermsii* is a corkscrew-shaped bacteria, approximately 3 to 25 µm in length, easily visualised with light microscopy.⁵ Exposure to *B. hermsii* peaks in July and August when travellers come in contact with the *O. hermsi* tick while vacationing to summer cabins or rural vacation homes in endemic regions.⁴

Acknowledgements The authors would like to thank Peter Sullivan for providing the information that the presence of corkscrew-shaped organisms on peripheral blood smear is pathognomonic for *Borrelia* infection, as other spirochaetes are too small to be visualised.

Contributors ESA, MO, MA and AMM were each involved in the care of the patient and contributed equally to writing the manuscript. MO annotated the figure.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

Figure 1 A peripheral blood smear demonstrating the presence of multiple corkscrew organisms (arrows), pathognomonic for *Borrelia* infection.

© BMJ Publishing Group Limited 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Aviles ES, Oakes M, Algranati M, et al. *BMJ Case Rep* 2020;**13**:e237296. doi:10.1136/bcr-2020-237296

REFERENCES

- 1 Dworkin MS, Schwan TG, Anderson DE, *et al.* Tick-Borne relapsing fever. *Infect Dis Clin North Am* 2008;22:449–68.
- 2 Butler T. The Jarisch–Herxheimer reaction after antibiotic treatment of spirochetal infections: a review of recent cases and our understanding of pathogenesis. *Am J Trop Med Hyg* 2017;96:46–52.
- 3 Malison MD. Relapsing fever. *JAMA* 1979;241:2819–20.
- 4 Dworkin MS, Shoemaker PC, Fritz CL, *et al.* The epidemiology of tick-borne relapsing fever in the United States. *Am J Trop Med Hyg* 2002;66:753–8.
- 5 Centers for Disease Control and Prevention. Relapsing Fever [Internet]. CDC, 2015. Available: <https://www.cdc.gov/relapsing-fever/index.html> [Accessed 17 Sep 2019].

Copyright 2020 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow