Left atrial vegetation after pulmonary vein isolation

Sean Gaine,¹ John Joseph Coughlan ,² Richard Szirt,³ Sadat Ali Edroos ⁴

¹Cardiology, St James's Hospital, Dublin, Ireland

²Cardiology, University Hospital Limerick, Limerick, Ireland

³Department of Cardiology, St George Hospital, Kogarah, New South Wales, Australia

⁴St James's Hospital, Dublin, Ireland

Correspondence to

Dr Sean Gaine; sgaine@tcd.ie

Accepted 29 June 2020

DESCRIPTION

A 67-year-old man presented with acute left hemiparesis 6 weeks after pulmonary vein isolation (PVI) for atrial fibrillation. His medical history was notable for hypertension and a metallic aortic valve which was implanted 23 years previously for bicuspid aortic stenosis. He was warfarinised for his metallic valve and his warfarin had been suspended for 5 days prior to ablation.

Clinical examination demonstrated left upper and lower limb weakness. CT brain suggested right middle cerebral artery branch occlusion. His international normalised ratio was high at 4.6 ruling out thrombolysis. The hemiparesis resolved within hours, and he was admitted for observation. He deteriorated the next day with bilateral weakness of left lower and right upper limbs. Following seizure activity, the patient dropped his Glasgow Coma Scale and required intubation. A repeat CT showed a small area of left frontal lobe bleed, suggestive of haemorrhagic transformation of embolus, deemed for conservative management following neurosurgical review. The patient was febrile and blood cultures grew *Streptococcus mitis* and *Streptococcus parasanguinis*. The warfarin was initially reversed and his cerebral haemorrhage appeared stable across serial imaging. As such anticoagulation was reinstated with a therapeutic unfractionated heparin infusion in view of his metallic aortic valve.

A transoesophageal echocardiogram (TOE) was performed, demonstrating a functioning prosthetic aortic valve and moderate mitral regurgitation. There was no vegetation or abscess identified. A clinical diagnosis of infective endocarditis (IE) with septic cerebral emboli was made despite the absence of echocardiographic evidence. He developed complete heart block the next day, requiring placement of a temporary pacing wire. This was replaced with a semipermanent externalised ventricular demand (VVI) pacemaker with active fixation

Figure 2 Gated cardiac CT confirming the presence of the vegetation in proximity to the oesophagus. It did not find any evidence of an atrio-oesophageal fistula (black arrow).

to permit long-term antibiotics in the context of sepsis, prior to committing to a permanent device.

Two weeks later, the patient became clinically unstable, required increasing inotropic support. Given ongoing concern regarding potential IE, repeat TOE was performed which demonstrated a large vegetation arising from the posterior left atrial wall (figure 1). Echocardiographic appearances were otherwise similar, with no evidence of valvular abscess or vegetation. Gated cardiac CT confirmed the presence of the vegetation in proximity to the oesophagus but did not find any evidence of an atrio-oesophageal fistula (figure 2). In view of the persistently poor neurological progress, a multidisciplinary team reached a consensus decision for conservative management of IE, with ongoing antibiotics, tracheostomy wean and neurological rehabilitation.

Septic vegetation after PVI has been reported previously.¹ Almost half of patients undergoing PVI are found to have some form of oesophageal injury or inflammation after procedure.² The resultant endothelial damage to the atrial wall may represent a potential substrate for infection. Atrio-oesophageal fistula is a rare complication of PVI

Figure 1 Transoesophageal echocardiogram demonstrating a large vegetation arising from the posterior left atrial wall (white arrow).

Learning points

- ▶ Septic vegetation is a rare but significant complication of pulmonary vein isolation.
- ▶ It is hypothesised that the damaged atrial endothelium may serve as a potential nidus for infection.
- ▶ Atrio-oesophageal fistula is a rare complication of pulmonary vein isolation and must be excluded prior to oesophagogastroduodenoscopy as this may result in insufflation and embolisation of air.

© BMJ Publishing Group Limited 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Gaine S, Coughlan JJ, Szirt R, et al. *BMJ Case Rep* 2020;**13**:e235833. doi:10.1136/bcr-2020-235833

and must be excluded prior to oesophagogastrroduodenoscopy as this may result in insufflation and embolisation of air.³

The patient was ultimately discharged to hospice care where he subsequently died due to complications of his illness.

Twitter John Joseph Coughlan @jjcoughl and Sadat Ali Edroos @saedroos

Contributors All authors discussed the report and contributed to the final manuscript. SAE, consultant cardiologist, and RS were instrumental in planning the structure of the case and interpreting the imaging. JJC and SG were involved in gathering information and ensuring data accuracy. Final approval of the version to be published was obtained from all contributors. The draft was revised and evaluated critically for important intellectual content.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Next of kin consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iDs

John Joseph Coughlan <http://orcid.org/0000-0001-6086-3279>

Sadat Ali Edroos <http://orcid.org/0000-0002-3267-6350>

REFERENCES

- 1 Weis S, Piorkowski C, Arya A, *et al*. Septic vegetation at the left atrial appendage entrance after pulmonary vein ablation for atrial fibrillation. *Europace* 2008;10:215–7.
- 2 Schmidt M, Nölker G, Marschang H, *et al*. Incidence of oesophageal wall injury post-pulmonary vein antrum isolation for treatment of patients with atrial fibrillation. *Europace* 2008;10:205–9.
- 3 Garuba HA. Left atrial-oesophageal fistula following ablation for atrial fibrillation. *J Am Coll Cardiol* 2014;63:A569.

Copyright 2020 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow