OPEN ACCESS

Anterior clinoidal mucocoele causing the orbital apex syndrome

Katsunari Kiko,¹ Kazuhiro Omura ,^{2,3} Shigeru Oya¹

¹Neurosurgery, Kokuho Asahi Chuo Hospital, Asahi, Chiba, Japan

²Otorhinolaryngology, The Jikei University School of Medicine, Minatoku, Tokyo, Japan

³Otorhinolaryngology, Dokkyo Medical University Saitama Medical Center, Koshigaya, Saitama, Japan

Correspondence to

Dr Kazuhiro Omura;
kazuhiro.omura@jikei.ac.jp

Accepted 1 July 2020

DESCRIPTION

A 63-year-old woman presented with diplopia and left upper-lid ptosis. Ophthalmological assessment showed oculomotor nerve palsy and low vision in the left eye. Brain MRI showed a signal corresponding to a cystic lesion in the left anterior clinoid process (ACP) with a hyperintense signal on T2-weighted images, an intermediate signal on diffusion-weighted images, and no suppression of the T1-hyperintense signal on fat saturated images. There was no fluid correction in her paranasal sinus (figure 1).

In a little while, the patient's symptoms improved; thus, outpatient follow-up was conducted at our department. However, 6 months later, she presented with acute loss of vision for 1 week; brain MRI showed a cystic lesion with definite growth in the left ACP.

The patient underwent surgery, where a left pterional craniotomy was performed. The ACP was drilled away through an extradural approach, allowing for the exposure of the mucocoele. Thereafter, the mucocoele was resected, and the roof of the optic nerve canal was drilled, decompressing the optic nerve (figure 2). The postoperative pathological results suggested inflammation of the nasal mucosa.

The patient's visual acuity recovered remarkably at 1 month postoperation.

The rate of ACP pneumatisation is around 9.5%. Pneumatised patterns are classified according to the route of pneumatisation, that is via the optic strut, the anterior root, or both.¹ Awareness of these variations in pneumatisation can prevent the destruction of the mucous membrane and the occurrence

Figure 2 Operative field view (from the direction of the arrow). (A) After the left pterional craniotomy, the mucocoele was exposed (circle) through an extradural approach. (B) The mucous membrane is held with forceps. (C) Operative field after removing the mucous membrane. Dot circle indicates the anterior clinoid process space.

of surgical complications, such as cerebrospinal fluid leakage, following an anterior clinoidectomy.

Blockage of the drainage of the paranasal sinus for any reason can result in the accumulation of trapped secretions and mucocoele. Owing to the proximity of the ACP to the optic nerve, patients with mucocoele in this location often present with ocular symptoms.²

Prompt surgical treatment is usually advocated to achieve best recovery.³ Surgical options include open surgical resection or transnasal endoscopic approach. The latter may be preferable to the cases where sinus cysts expand to the ACP. Nonetheless, if the mucocoele is localised in the ACP, cystic resection by craniotomy may be safer, because the cystic area is over the optic nerve by transnasal endoscopic surgery. However, a history of sinusitis can aid in the diagnosis of a reversible form of optic

Figure 1 Preoperative CT/MRIs. (A) Axial T1-weighted images showing an anterior clinoidal mucocoele (arrow). (B) Axial T2-weighted images showing an anterior clinoidal mucocoele (arrow). (C) Coronal T2-weighted image showing that the left optic nerve (arrow head) exists more medially than the mucocoele (arrow). (D) Coronal CT suggesting that the mucocoele dissolves the anterior clinoidal process and optic strut (arrow). There are onodi cells bilaterally (square).

Learning points

- ▶ Awareness of anterior clinoid process (ACP) pneumatisation can prevent the occurrence of cerebrospinal fluid leakage as a result of an anterior clinoidectomy.
- ▶ Patients with anterior clinoid mucocoele presenting with visual loss who have a history of sinusitis receive only medical treatment with antibiotics and can thus avoid surgery.
- ▶ An endoscopic approach may be preferable to cases where the sinus cysts expand to the ACP. However, if the mucocoele localized in the ACP, the cystic resection by craniotomy or endoscopic trans-orbital approach may be safer, because the cystic area is over the optic nerve by transnasal endoscopic surgery.

© BMJ Publishing Group Limited 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Kiko K, Omura K, Oya S. *BMJ Case Rep* 2020;**13**:e236527. doi:10.1136/bcr-2020-236527

neuritis, for which proper medical treatment with steroids and antibiotics may be preferable to surgical intervention.²

Twitter Kazuhiro Omura @KAZ

Contributors KK drafted the manuscript. KO supervised the manuscript. SO edited the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which

permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Kazuhiro Omura <http://orcid.org/0000-0002-4337-4745>

REFERENCES

- 1 Mikami T, Minamida Y, Koyanagi I, *et al.* Anatomical variations in pneumatization of the anterior clinoid process. *J Neurosurg* 2007;106:170–4.
- 2 Deshmukh S, DeMonte F. Anterior clinoidal mucocele causing optic neuropathy: resolution with nonsurgical therapy. Case report. *J Neurosurg* 2007;106:1091–3.
- 3 Abozed M, Alsulaiti G, Almannai F, *et al.* Anterior clinoid mucocele causing optic neuropathy: a case report and review of literature. *eNeurologicalSci* 2017;7:57–9.

Copyright 2020 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow