Retromolar intubation for severe trismus: a less invasive alternative to tracheostomy and submental intubation

Brandon Michael Togioka ,¹ Vincent Pinkert,¹ Mathew Geltzeiler,² Elliza M Chen¹

¹Anesthesiology & Perioperative Medicine, Oregon Health & Science University, Portland, Oregon, USA

²Otolaryngology, Oregon Health & Science University, Portland, Oregon, USA

Correspondence to

Dr Brandon Michael Togioka;
togioka@ohsu.edu

Accepted 23 December 2020

DESCRIPTION

A 67-year-old woman with renal cell cancer metastatic to the paranasal sinuses presented for nasal endoscopy and palliative debridement. The extensive sinonasal mass was a contraindication to transnasal intubation. Tumour invasion into the lateral and medial pterygoid muscles as well as prior palliative sinus radiotherapy resulted in severe trismus with 5 mm interincisor distance—too small to accommodate an endotracheal tube ([figure 1](#)). Debridement was delayed for a year in the hope that jaw physical therapy would increase mouth opening and allow for oral intubation—intubation was deemed necessary due to the vascularity of the metastases. After a year, mouth opening remained restricted and awake tracheostomy was planned.

The anaesthesia team noticed the patient's facial skin was malleable. Skin retraction at the left angle of the mouth revealed a retromolar space large enough to fit an endotracheal tube. Tracheostomy was not anticipated to be difficult. The neck was long and supple. Laryngeal cartilage and sternal notch were visible. Cricoid cartilage was palpable.

A plan was made for fiberoptic intubation through the retromolar space with tracheostomy backup. The otolaryngologist injected 10 mL of 1% lidocaine containing 10 µg/mL epinephrine into the anterior neck. Then, under dexmedetomidine sedation at 0.8 µg/kg/hour, the airway was anaesthetised with 5 mL of 4% nebulised lidocaine, 1 inch of 5% lidocaine ointment to the posterior tongue and 3 mL of 4% lidocaine via transtracheal block. A 6.5 mm inner diameter flex-tip endotracheal tube was inserted behind the left maxillary molar. A 5.5 mm diameter flexible video endoscope guided the endotracheal tube into the pharynx and past the vocal cords ([figure 2](#)). Surgery was uneventful, blood loss was 100 mL and there were no problems with endotracheal tube compression or kinking. The patient was extubated awake and discharged home the next day.

One in four patients with head and neck cancer acquires trismus.¹ Tracheal intubation is often required for these patients due to the risk for airway bleeding. Trismus makes bag mask ventilation more difficult and severe trismus precludes

Figure 1 Measurement of interincisor distance with maximal mouth opening.

© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Togioka BM, Pinkert V, Geltzeiler M, et al. *BMJ Case Rep* 2021;**14**:e239817. doi:10.1136/bcr-2020-239817

Figure 2 Retraction of skin at the angle of the mouth to reveal endotracheal tube position in the left retromolar space.

oropharyngeal and laryngeal mask airway insertion. Standard practice for patients with severe trismus and contraindication to nasal intubation is to perform a submental intubation or an awake tracheostomy. These procedures are invasive and associated with a higher rate of infection, bleeding, fistula formation, tracheal injury, voice change and cosmetic deformity, compared with oral intubation.^{2,3}

The retromolar space is present in all individuals. It can be visualised by retracting the skin at the angle of mouth. The retromolar space is bound posteriorly by the ascending ramus of the mandible, superiorly by the maxillary tuberosity and anteriorly

by the back molars.⁴ While there are reports of 'retromolar intubations,' most descriptions are actually standard intubations with intraoperative shifting of the endotracheal tube to the retromolar space prior to maxillomandibular fixation.^{3,5-10} Our report differs from these in that we used the retromolar space to access the oropharynx. This case highlights an additional non-invasive point of access to the oropharyngeal cavity when midline oral and nasal passages are unavailable.

Acknowledgements The authors would like to thank Ms. Marie Kane for her assistance in editing and formatting.

Contributors The work was supervised by BMT. The patient was under the care of MG and EMC. The report was written by BMT, VP, MG and EMC. All authors give final approval for the version submitted and agree to be accountable for all aspects of the work.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Brandon Michael Togioka <http://orcid.org/0000-0002-8285-7811>

REFERENCES

- 1 van der Geer SJ, van Rijn PV, Kamstra JI, *et al.* Prevalence and prediction of trismus in patients with head and neck cancer: a cross-sectional study. *Head Neck* 2019;41:64–71.
- 2 Cipriano A, Mao ML, Hon HH, *et al.* An overview of complications associated with open and percutaneous tracheostomy procedures. *Int J Crit Illn Inj Sci* 2015;5:179–88.
- 3 Dutta A, Kumar V, Saha SS, *et al.* Retromolar tracheal tube positioning for patients undergoing faciomaxillary surgery. *Can J Anaesth* 2005;52:341.
- 4 Truong A, Truong D-T. Retromolar fiberoptic orotracheal intubation in a patient with severe trismus undergoing nasal surgery. *Can J Anaesth* 2011;58:460–3.
- 5 Arora S, Rattan V, Bhardwaj N. An evaluation of the retromolar space for oral tracheal tube placement for maxillofacial surgery in children. *Anesth Analg* 2006;103:1122–5.

Patient's perspective

I was worried going into surgery that I would have a tracheostomy because that is what my ear, nose and throat doctor told me. I was told I would have to have the tracheostomy surgery awake. This scared me and also worried me that it would affect my quality of life after surgery. I was very grateful to the anaesthesia team for avoiding the tracheostomy.

Learning points

- The retromolar space can be used to access the oropharyngeal cavity and intubate patients with severe trismus and nasal passage blockage.
- Retromolar intubation is less invasive and is associated with less morbidity than tracheostomy and submental intubation.
- The retromolar space, which can be visualised by retracting skin at the corner of the mouth, is bound posteriorly by the ascending ramus of the mandible, superiorly by the maxillary tuberosity and anteriorly by the back molars.

- 6 Habib A, Zanaty OM. Use of retromolar intubation in paediatric maxillofacial trauma. *Br J Anaesth* 2012;109:650–1.
- 7 Malhotra N. Retromolar intubation--a simple alternative to submental intubation. *Anaesthesia* 2006;61:515–6.
- 8 Martinez-Lage JL, Eslava JM, Cebrecos AI, et al. Retromolar intubation. *J Oral Maxillofac Surg* 1998;56:302–5.
- 9 Rao M, Shetty D, Goyal KA, et al. Retromolar intubation: a better alternative to submental intubation or tracheostomy for dental occlusion by intermaxillary fixation. *Saudi J Anaesth* 2015;9:222–4.
- 10 Gibbons AJ, Hope DA, Silvester KC. Oral endotracheal intubation in the management of midfacial fractures. *Br J Oral Maxillofac Surg* 2003;41:259–60.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow