Psychiatric non-epileptic seizure: diagnostic utility of density spectral array

Shuichiro Neshige

Department of Clinical Neuroscience and Therapeutics, Graduate School of Biomedical and Health Sciences, Hiroshima University, Hiroshima, Japan

Correspondence to

Dr Shuichiro Neshige;
s-neshige@hiroshima-u.ac.jp

Accepted 21 December 2020

DESCRIPTION

A 40-year-old woman with a history of insomnia admitted to the emergency room for generalised convulsion lasting for more than 30 min. The attack was characterised by bilateral asynchronous hyperkinetic prolonged jerking with loss of responsiveness. The serial intravenous diazepam (10 mg) had a limited effect. As the attack was accompanied by a decrease in the oxygen saturation due to the apnoea, she was intubated followed by deep sedation with barbiturates for almost 24 hours. Continuous electroencephalography (EEG) monitoring revealed a suppression-burst pattern during the sedation. However, after the discontinuation of sedation, the hyperkinetic attack reappeared immediately. The ictal EEG during this attack was obscured by the motion artefacts (figure 1). Given the unresponsiveness, she seemed to be impaired awareness. However, density spectral array (DSA)^{1,2} that was analysed from the data of O1 and O2 electrodes during the prolonged attack clearly showed preserved alpha-band activity (figure 2). This activity was consistent with the posterior dominant rhythm, that is, the presence of this preserved background alpha activity suggested that she was aware during the attack. Thus, she was diagnosed with psychiatric non-epileptic seizure (PNES).³ The attack of PNES gradually disappeared after a liaison psychiatry therapy with drugs for anxiety and insomnia. While EEG waves that were recorded during a hyperkinetic attack were often obscured by motion artefacts, DSA was occasionally useful to

Figure 1 Electroencephalography (EEG) during a hyperkinetic attack. EEG waves are shown on the monopolar montage with the time constant of 0.1 s and high-frequency filter of 120 Hz. Artefact of electromyogram is evident.

Figure 2 Density spectral array (DSA). The DSA is the data based on the time–frequency analysis calculated from the amplitude of O1 and O2 electrodes during the electroencephalogram (EEG) examination. Preserved alpha activity, 8–9 Hz, is visible (white arrow). Power spectrum change due to motion artefact is also shown (yellow arrow). The asterisk indicates the time window of EEG in figure 1.

Patient's perspective

I was relieved to hear the diagnosis. As the number of medicines has decreased, I feel better now.

Learning points

- ▶ Hyperkinetic prolonged attack is one of the characteristic semiology of psychiatric non-epileptic seizure (PNES).
- ▶ Conventional EEG evaluation is often limited by motion artefacts during the seizure.
- ▶ Density spectral array from bilateral occipital region occasionally reveals the posterior-dominant rhythm which may assist in the diagnosis of PNES.

clarify the background EEG activity which might assist in the diagnosis of PNES.

Contributors SN: Designed and conceptualised the study, analysed the data and drafted the manuscript including intellectual content.

Funding JSPS KAKENHI (grant number:20K16601).

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Neshige S. *BMJ Case Rep* 2021;**14**:e241015. doi:10.1136/bcr-2020-241015

ORCID iD

Shuichiro Neshige <http://orcid.org/0000-0002-5879-155X>

REFERENCES

- 1 Katoh H, Kinefuchi Y, Takiguchi M, *et al.* Density-modulated T's array, a new technique of processed electroencephalogram, for monitoring the effects of midazolam and nitrous oxide during spinal anesthesia. *J Anesth* 1994;8:434–40.
- 2 Neshige S, Matsumoto R, Inouchi M, *et al.* Absence of an autonomic sign assists in the diagnosis of extratemporal lobe epilepsy manifesting generalized convulsion with retained awareness. *Intern Med* 2019;58:1151–5.
- 3 LaFrance WC, Baker GA, Duncan R, *et al.* Minimum requirements for the diagnosis of psychogenic nonepileptic seizures: a staged approach: a report from the International League against epilepsy nonepileptic seizures task force. *Epilepsia* 2013;54:2005–18.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow