High-attenuation mucus in a patient with allergic bronchopulmonary aspergillosis

Sachiho Inada, Hiroshi Sugimoto , Tomomi Terashita, Kyosuke Nakata

Department of Respiratory Medicine, Konan Medical Center, Kobe, Hyogo, Japan

Correspondence to
Dr Hiroshi Sugimoto;
dr.sugimoto@gmail.com

Accepted 23 September 2021

DESCRIPTION

A 63-year-old Japanese woman presented with a 9-month history of productive cough. Although she had been treated with oral antibiotics (amoxicillin and levofloxacin), her symptoms worsened. She had no history of allergic diseases, including asthma, except for known allergies to eggs. She had never smoked.

Her vital signs and physical examination were unremarkable. Laboratory tests revealed white blood cell count of $12.26 \times 10^9/L$ (normal: $3.3\text{--}8.6 \times 10^9/L$) with absolute eosinophil count of $2.45 \times 10^9/L$ (normal: $0.10\text{--}0.30 \times 10^9/L$), serum total IgE levels of 235 IU/mL (normal: <173 IU/mL), *Aspergillus*-specific IgE levels of 5.49 UA/mL (normal: <0.35 UA/mL), positive for *Aspergillus*-specific precipitin and negative for serum antigen of *Aspergillus* spp.

Chest radiography showed consolidation in the left middle lung field. Chest CT revealed high-attenuation mucus (HAM), a typical finding in allergic bronchopulmonary aspergillosis (ABPA), and central bronchiectasis (figure 1). We performed a transbronchial biopsy of the mucus plugs in the left upper lobe. Pathological findings showed remarkable eosinophil infiltration with Charcot-Leyden crystals, and branching hyphae on Grocott stain (figure 2). In the culture of bronchial lavage fluid, *Aspergillus fumigatus* was detected.

Consequently, the patient fulfilled 8 of 10 components in the criteria for allergic bronchopulmonary mycosis (ABPM) by the Japan ABPM Research Programme.¹ Therefore, we confirmed the diagnosis of ABPA, and subsequently, prednisolone with tapering was administered. After the treatment, her symptoms, eosinophilia and

Figure 2 Grocott stain for biopsied specimen of mucus plugs shows branching hyphae (arrows).

consolidation on chest radiography were gradually improved.

ABPM is a disease caused by an allergic reaction to fungi in the lower respiratory tract, especially those caused by *Aspergillus* spp are called ABPA.² There are several clinical diagnostic criteria for ABPM, including Rosenberg-Patterson criteria proposed in 1977,³ the International Society for Human and Animal Mycology (ISHAM) criteria in 2013⁴ and the Japan ABPM Research Programme criteria in 2020.¹

The Rosenberg-Patterson criteria have a high specificity of 98.4% but a low sensitivity of 49.2%, while the ISHAM criteria have a sensitivity of 82.7% and a specificity of 86.8%. In contrast, the sensitivity and specificity of the Japan ABPM Research Programme criteria are 94.4% and 90.0%, respectively, indicating that ABPM can be diagnosed with high accuracy.¹

ABPM often shows mucoid impaction, and its differential diagnosis includes endobronchial lesions or foreign bodies, bronchial atresia and bronchiectasis due to various causes.⁵ HAM is defined as mucus plugs with higher densities than paravertebral muscles on high-resolution CT, and the CT density value of 70 Hounsfield units is an

Figure 1 Chest CT shows high-attenuation mucus (arrow).

© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Inada S, Sugimoto H, Terashita T, et al. *BMJ Case Rep* 2021;**14**:e246729. doi:10.1136/bcr-2021-246729

Learning points

- ▶ Allergic bronchopulmonary mycosis can be diagnosed by Japan Allergic Bronchopulmonary Mycosis Research Programme criteria with higher accuracy than other criteria.
- ▶ High-attenuation mucus is an important finding in the early diagnosis of allergic bronchopulmonary mycosis.

adequate cut-off value.⁶ It has been assumed that iron, manganese and calcium produced by fungi in the mucus may cause higher densities.⁶ The sensitivity and specificity of HAM for ABPA are reported as 39.7% and 100%, respectively.⁷ HAM is an important finding in the early diagnosis of ABPM.

Acknowledgements We thank Editage (www.editage.com) for English language editing.

Contributors SI and HS drafted the manuscript. TT and KN revised the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s)

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Hiroshi Sugimoto <http://orcid.org/0000-0002-2053-8858>

REFERENCES

- 1 Asano K, Hebisawa A, Ishiguro T, *et al.* New clinical diagnostic criteria for allergic bronchopulmonary aspergillosis/mycosis and its validation. *J Allergy Clin Immunol* 2021;147:1261–8.
- 2 Knutsen AP, Bush RK, Demain JG, *et al.* Fungi and allergic lower respiratory tract diseases. *J Allergy Clin Immunol* 2012;129:280–91.
- 3 Rosenberg M, Patterson R, Mintzer R, *et al.* Clinical and immunologic criteria for the diagnosis of allergic bronchopulmonary aspergillosis. *Ann Intern Med* 1977;86:405–14.
- 4 Agarwal R, Chakrabarti A, Shah A, *et al.* Allergic bronchopulmonary aspergillosis: review of literature and proposal of new diagnostic and classification criteria. *Clin Exp Allergy* 2013;43:850–73.
- 5 Agarwal R, Aggarwal AN, Gupta D. High-attenuation mucus in allergic bronchopulmonary aspergillosis: another cause of diffuse high-attenuation pulmonary abnormality. *AJR Am J Roentgenol* 2006;186:904.
- 6 Phuyal S, Garg MK, Agarwal R, *et al.* High-Attenuation mucus impaction in patients with allergic bronchopulmonary aspergillosis: objective criteria on high-resolution computed tomography and correlation with serologic parameters. *Curr Probl Diagn Radiol* 2016;45:168–73.
- 7 Agarwal R, Maskey D, Aggarwal AN, *et al.* Diagnostic performance of various tests and criteria employed in allergic bronchopulmonary aspergillosis: a latent class analysis. *PLoS One* 2013;8:e61105.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow