

Progressive myocardial dysfunction following COVID-19

Mirosława Maria Gorecka ¹, Sharmaine Thirunavukarasu,¹ Eylem Levelt,^{1,2} John P Greenwood ^{1,2}

¹Leeds Institute for Cardiovascular and Metabolic Medicine, University of Leeds, Leeds, UK

²Cardiology, Leeds Teaching Hospitals NHS Trust, Leeds, UK

Correspondence to

Professor John P Greenwood; J.Greenwood@leeds.ac.uk

Accepted 29 October 2021

DESCRIPTION

A 66-year-old man with background of type 2 diabetes mellitus, hypertension and functionally non-significant coronary artery disease (CT coronary angiogram 2 years prior demonstrated moderate ostial atheroma in the first diagonal branch and in the dominant right coronary artery, however, a subsequent stress echocardiogram did not reveal any inducible ischaemia) presented to the hospital with non-productive cough and fever, and was subsequently confirmed to have COVID-19. He denied dyspnoea and chest pain. Laboratory testing revealed elevated troponin T (peak 296 ng/L; Upper Limit of Normal (ULN) <57 ng/L), elevated D-dimer (peak 504 ng/mL; ULN <230 ng/mL) and acute kidney injury stage III. He was treated conservatively with intravenous fluid replacement and made an uneventful recovery.

He underwent cardiovascular magnetic resonance (CMR) imaging as part of a UK, multi-centre, Urgent Public Health research study (ISRCTN58667920).¹ CMR revealed mild impairment in left ventricular ejection fraction (LVEF 45%). Stress perfusion CMR revealed no inducible ischaemia, whereas late gadolinium enhancement images showed patchy mid-wall hyperenhancement consistent with COVID-19 myocarditis (figure 1A). T1-mapping confirmed myocardial injury in a non-ischaemic pattern (figure 1B) and global myocardial T2 was also elevated consistent with acute inflammation (figure 1C).

At 6 months, the patient felt well. Repeat investigations, however, revealed persistent elevation of troponin T (597 ng/L) and brain natriuretic peptide (BNP) (494 ng/L; ULN<331 ng/L). Repeat CMR showed significant deterioration




Figure 1 Baseline CMR scan. (A) Short-axis late gadolinium enhancement image demonstrates patchy mid-wall hyperenhancement consistent with COVID-19 myocarditis (see arrow); (B) T1-mapping image shows myocardial injury in a non-ischaemic pattern (arrow); (C) Image of a T2 map demonstrates global elevation of T2 consistent with acute inflammation. CMR, cardiovascular MR.




Figure 2 Follow-up scan at 6 months. (A) LGE image shows persisting mid-wall hyperenhancement (arrows); (B) T1-mapping image demonstrates elevated myocardial T1 (arrow); (C) T2 map shows normalisation of T2. LGE, late gadolinium enhancement.

in LVEF to 28%, persisting mid-wall hyperenhancement (figure 2A) and elevated myocardial T1 (figure 2B). Myocardial T2 had normalised (figure 2C) and stress perfusion CMR was again unremarkable.

Despite the absence of cardiovascular symptoms and an uneventful recovery from COVID-19, CMR demonstrated progressive myocardial dysfunction, which warranted introduction of heart failure therapies and immunosuppression.² The patient was treated with a beta-blocker and an angiotensin-II receptor blocker in the community. The ultimate goal was to introduce an angiotensin receptor neprilysin inhibitor (ARNI), mineralocorticoid receptor antagonist and a sodium-glucose cotransporter-2 (SGLT2) inhibitor, alongside immunosuppression; however, the patient has since failed to attend his hospital appointments.

Although the differential diagnosis in such a case is quite broad, in view of the functionally non-significant (non-obstructive) coronary artery disease, lack of any specific cardiac symptoms and an elevated, but relatively static troponin T rise, COVID-19-associated myocardial injury was felt to be the most likely clinical diagnosis (biopsy data were not available). As the patient failed to attend further hospital appointments, the final case disposition is uncertain.

We believe this case highlights the importance of follow-up imaging in cases of COVID-19 myocarditis, as ongoing and progressive



© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Gorecka MM, Thirunavukarasu S, Levelt E, et al. *BMJ Case Rep* 2021;**14**:e246291. doi:10.1136/bcr-2021-246291

Learning points

- ▶ COVID-19 can cause progressive myocardial dysfunction.
- ▶ Follow-up imaging is required to establish the diagnosis and to guide appropriate therapies.

myocardial injury may occur in the absence of any obvious symptoms.

Contributors MMG was responsible for performing the CMR scan, collection of data and write up of the case. ST was responsible for performing the CMR scan, collection of data and final review of manuscript. EL was responsible for literature review and final review of the case. JPG was responsible for patient's care in CMR, final review and approval of final version of the case.

Funding This study was funded by NIHR-UKRI COVID-19 Rapid Response Rolling Call (COV0254) (COV0254).

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s)

Provenance and peer review Not commissioned; externally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful,

non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

ORCID iDs

Mirosława Maria Gorecka <http://orcid.org/0000-0002-5688-8980>

John P Greenwood <http://orcid.org/0000-0002-2861-0914>

REFERENCES

- 1 Gorecka M, McCann GP, Berry C, *et al*. Demographic, multi-morbidity and genetic impact on myocardial involvement and its recovery from COVID-19: protocol design of COVID-HEART-a UK, multicentre, observational study. *J Cardiovasc Magn Reson* 2021;23:77.
- 2 Nicol M, Cacoub L, Baudet M, *et al*. Delayed acute myocarditis and COVID-19-related multisystem inflammatory syndrome. *ESC Heart Fail* 2020;7:4371–6.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow