Graves' disease in children: an enlarged goitre causes severe tracheal stenosis

Shota Hiroshima,¹ Yushi Ueki,² Keisuke Yamazaki,² Keisuke Nagasaki ¹

¹Division of Pediatrics, Department of Homeostatic Regulation and Development, Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan

²Department of Otolaryngology Head and Neck Surgery, Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan

Correspondence to

Dr Keisuke Nagasaki;
nagasaki@med.niigata-u.ac.jp

Accepted 7 October 2021

DESCRIPTION

A teenage high school student underwent an X-ray of the chest during a routine physical examination at the time of admission to school, which revealed marked tracheal stenosis (figure 1A). The boy was diagnosed with Graves' disease a year before due to goitre and easy fatigability and was under treatment with an antithyroid drug. He remained in an euthyroid state with 15–20 mg/day of methimazole. Since the disease onset, he had been reporting discomfort during swallowing owing to the enlarged goitre. Ultrasonography of the thyroid performed 6 months earlier showed a highly diffused goitre with a heterogeneous echogenic pattern (40 mm width, 40 mm thickness, 60 mm height in the right lobe and 35 mm width, 40 mm thickness, and 60 mm height in the left lobe). No apparent tracheal stenosis was noted. Furthermore, with the increase in duration of physical education after entering the high school, the boy became aware of dyspnoea during exercise. The CT scan of the neck showed that the minimum tracheal diameter was 3 mm (figure 1B), indicating severe tracheal stenosis. Therefore, total thyroidectomy (thyroid weight: 300 g) was performed with an oral tracheal intubation under conscious control. Histopathological examination revealed no neoplastic lesion. The tracheal stenosis did not improve immediately after

thyroidectomy; therefore, intraoperative tracheostomy was performed. The cannula was removed on postoperative day 5, and tracheal stenosis was not detected on day 14 (figure 1C). Consequently, the respiratory distress during exercise disappeared.

Tracheal stenosis occurs rarely due to an enlarged goitre even in a benign goitre associated with Graves' disease.¹ Further, airway stabilisation is a significant problem in the case of goitre with tracheal stenosis. Previous reports suggest that extracorporeal membrane oxygenation was used at the time of thyroid surgery in some cases,² while in others, a tracheal stent was placed preoperatively.³ An enlarged goitre may cause dyspnoea due to tracheal stenosis even in adolescents with Graves' disease whose thyroid function is normally controlled. After the symptoms of tracheal compression become clinically evident, the occurrence of complete airway occlusion may be sudden and unpredictable. Hence, when goitre-induced tracheal stenosis is evident, an early thyroidectomy is recommended.

Learning points

- ▶ An enlarged goitre may cause dyspnoea due to tracheal stenosis even in adolescents with Graves' disease whose thyroid function is normally controlled.
- ▶ Early thyroidectomy is recommended in patients with severe tracheal stenosis due to goitre associated with Graves' disease.

Figure 1 (A) A routine X-ray of the chest during physical examination for admission to a school revealed severe tracheal stenosis. (B) Horizontal image of CT of the cervical. Tracheal diameter is 3 mm with marked tracheal stenosis. (C) X-ray of the chest 2 weeks after total thyroidectomy. Tracheal stenosis has improved.

Contributors KN and SH conceptualised and designed the study, drafted the initial manuscript, and reviewed and revised the manuscript. YU and KY collected the data and reviewed and revised the manuscript. All the authors approved the final version of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Consent obtained from parent(s)/guardian(s)

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Keisuke Nagasaki <http://orcid.org/0000-0002-5882-661X>

REFERENCES

- 1 Alfonso A, Christoudis G, Amaruddin Q, et al. Tracheal or esophageal compression due to benign thyroid disease. *Am J Surg* 1981;142:350–4.
- 2 Shao Y, Shen M, Ding Z, et al. Extracorporeal membrane oxygenation-assisted resection of goiter causing severe extrinsic airway compression. *Ann Thorac Surg* 2009;88:659–61.
- 3 Piao M, Yuan Y, Wang Y, et al. Successful management of trachea stenosis with massive substernal goiter via thacheobronchial stent. *J Cardiothorac Surg* 2013;8:212.

© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Hiroshima S, Ueki Y, Yamazaki K, et al. *BMJ Case Rep* 2021;14:e246906. doi:10.1136/bcr-2021-246906

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow