OPEN ACCESS

Subinvolution of the placental site as the cause of hysterectomy in young woman

Olga Triantafyllidou,¹ Stavroula Kastora ,^{2,3} Irini Messini,⁴ Dimitrios Kalampokis⁵

¹Reproductive Medicine Unit, Maternity Hospital Lito, Athens, Greece

²Department of Medicine, University of Aberdeen College of Life Sciences and Medicine, Aberdeen, UK

³Department of General Surgery, Aberdeen Royal Infirmary, Aberdeen, UK

⁴Department of Pathology, Maternity Hospital Lito, Athens, Greece

⁵Department of Obstetrics, Maternity Hospital Lito, Athens, Greece

Correspondence to

Dr Stavroula Kastora; stavroula.kastora@abdn.ac.uk

Accepted 9 January 2021

SUMMARY

Subinvolution of placental sites (SPSs) is a rare but severe cause of secondary postpartum haemorrhage (PPH). SPS is characterised by the abnormal persistence of large, dilated, superficially modified spiral arteries in the absence of retained products of conception. It is an important cause of morbidity and mortality of young women. In this study, we present a case of secondary PPH in a young woman after uncomplicated caesarean delivery who was deemed clinically unstable, and finally, underwent emergent total abdominal hysterectomy. We reviewed the literature with an emphasis on the pathophysiology of this situation. Treatment of patients with SPS includes conservative medical therapy, hysterectomy and fertility-sparing percutaneous embolotherapy.

BACKGROUND

Postpartum haemorrhage (PPH) is a major problem worldwide and it still represents a leading cause of maternal deaths.¹ The incidence of PPH is approximately 5%–20% of all deliveries, with the great percentage to be attributed to the developing countries.^{1,2} It is commonly divided into two groups: primary and secondary PPH. Primary postpartum bleeding occurs within the first 24 hours after delivery, whereas secondary PPH occurs between 24 hours and 6 weeks post partum and it is not particularly associated with the way of delivery (vaginal or caesarean section). The causes of primary PPH include uterine atony, retained placenta, congenital of acquired disorders of coagulation and trauma. The most common cause of primary PPH worldwide is still uterine atony. Secondary PPH accounts only for 1% of all labours and its aetiology includes severe endometritis, retained products of conception, gestational trophoblastic disease, uterine artery pseudoaneurysm and subinvolution of the placental site (SPS).

SPS is defined as delayed or inadequate physiological closure and sloughing of the superficial modified spiral arteries at the placental site (failed process of normal involution).³ It was reported in 1910 for the first time in the literature by Küster, and only in 1945 Rutherford and Hertig described clinical and pathological findings of ‘Non-involution of the Placental Bed’.^{4,5} It is difficult to estimate the precise frequency of SPS in all secondary PPH because the histopathological examination of aspiration products after uterine surgical evacuation is essential to the diagnosis. The clinical management of SPS demands immediate action with little time, due to severity of haemorrhage. In this report, we

present an additional case of severe SPS in young woman, who underwent hysterectomy, as well as the review of the literature.

CASE PRESENTATION

A 34-year-old nulliparous primigravida Caucasian woman with a normal single pregnancy after ovarian stimulation and In Vitro Fertilisation (IVF) treatment due to poor ovarian reserve was admitted at 38+3 weeks of pregnancy with signs of labour. In spite of good uterine contractions, the cervical dilatation after 7 hours was only 2 cm and caesarean section was performed as a result of the failure to progress in first stage. The postpartum period was uneventful and patient was discharged with recommendations on fourth postoperative day.

INVESTIGATIONS

Three weeks later, she was admitted with severe vaginal bleeding. Transvaginal ultrasound (US) examination revealed possible organised blood clots inside the uterus. There was not echogenic material suspicious for retained placenta within the endometrial cavity using the gray-scale US findings and colour Doppler US imaging was negative for the presence of internal vascular flow. Moreover, the patient did not have any signs or symptoms of inflammation (endometritis), fever, rigours, pelvic or abdominal pain. Complete blood count was normal [haemoglobin (Hb): 12.1 g/dL and platelets: $325 \times 10^9/L$, white cell count $8.500 \times 10^9/L$, CRP 0.7 mg/L]. Serum β -subunit of hCG gonadotropin (β -hCG) was negative and there were no signs of coagulopathy (prothrombin time, activated partial thromboplastin time, International Normalised Ratio (INR) and plasma fibrinogen levels were within normal levels. The patient was resuscitated with warm fluids and blood was sent for cross match (four units of red blood cells and four units of fresh frozen plasma). Uterine curettage was performed and 150 mL of curetted material was negative for retained placental products. Despite curettage of the endometrium, rectal misoprostol and intravenous administration of ergometrine maleate (Mitrotan, GAP), the bleeding did not cease and intrauterine tamponade was performed. Twelve hours after curettage, the patient was not stabilised, fainted and kept bleeding with gradual decline of Hb. She was tachycardic with a blood pressure of 100/65 mm Hg and her haemoglobin reached 6 g/dL. In combination with close monitoring of haemodynamics and urine output, patient's coagulation profile was repeatedly measured (prothrombin time/activated

© BMJ Publishing Group Limited 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Triantafyllidou O, Kastora S, Messini I, et al. *BMJ Case Rep* 2021;**14**:e238945. doi:10.1136/bcr-2020-238945

Figure 1 Subinvolution of uteroplacental arteries. Dilated, clustered, myometrial arteries with partial obliteration by thrombi (haematoxylin-eosin staining, magnification: $\times 400$).

partial thromboplastin time increased until 12.6 s /41.4 s, respectively, and fibrinogen level was 1.7 g/L).

TREATMENT

The patient was given four units of packed red blood cells and four units of fresh frozen plasma, however abdominal US re-examination revealed an enlarged uterus with the accumulation of blood around the intrauterine tampon. Pelvic CT and MRI imaging with contrast did not reveal a pathological cause of bleeding. Given that the patient was deemed clinically unstable, an emergency total abdominal hysterectomy was performed without further imaging such as angiography. Histopathological examination of uterus identified large thrombus in the endometrial cavity and the typical findings of subinvolved placental bed vessels. Microscopic sections of myometrium revealed large, dilated superficial myometrial blood vessels with thickened walls at the placental implantation site indicative of SPS (figure 1). There was no evidence of retained products of conception or inflammation of the decidua neither in the curettage material nor in the removed uterus. The diagnosis of SPS was made.

OUTCOME AND FOLLOW-UP

The neonate was delivered safely without complications and was admitted in the neonate ward for a day of observations given the maternal complications. Ovaries were spared during the total abdominal hysterectomy in view of preserving maternal ovarian fertility with the further possibility of surrogacy.

DISCUSSION

PPH is a serious obstetric problem, one of the leading causes of maternal mortality and morbidity. Secondary PPH is less frequently addressed, partly, due to the majority of patients who are treated medically on an outpatient basis, without hospitalisation. SPS is an important cause of secondary PPH and it should be considered in patient with severe PPH, due to its association with life-threatening haemorrhage. SPS is diagnosed by excluding other causes of persistent secondary postpartum haemorrhagic vaginal discharge. These include endometritis, retained products of placenta, gestational trophoblastic disease and coagulopathy. The diagnosis of SPS can be confirmed only histologically as it happened in our case. Excluding all above-mentioned diagnosis by clinical, US examination, laboratory tests and histology of curetted products, the diagnosis of PPH remained unknown until the final histopathological examination of uterus. However,

Petrovitch *et al*⁶ in their study mentioned that the suspicion of this situation may done through the sonographic characteristics of the pathological vessels. Placental site subinvolution may be suspected in US examination in the infrequent case where hypoechoic tortuous vessels are visible along the inner third of the myometrium.⁶ Pulsed wave doppler US then might indicate increased peak systolic velocity (PSV; >0.83 m/s; normal 0.22 m/s 3 days post partum, falling to 0.10 m/s after 6 weeks). The increased vascularity areas may correlate with the placental implantation site documented by predelivery. However, these findings are not always pathognomonic and may overlap with congenital or acquired arteriovenous malformations or retained products of conception.

Large uteroplacental arteries that were modified for pregnancy by the extravillous trophoblasts remain patent past the immediate postpartum period, failing to undergo the normal process of involution. The incidence of PPH from subinvolution is most common in the second week post partum.⁷ The precise incidence of severe secondary PPH was estimated by Dossou *et al*,⁸ who described the distribution of the different causes of severe secondary PPH after 26 023 deliveries. The incidence of severe secondary PPH was very low, only 0.23% (n: 60/26 023). Subinvolution of the placental bed was noted in 13.3% of the patients, while the placental retention was the most common cause of haemorrhages (30.0%). In contrast with clinical doctors, veterinarians are more familiar with this condition, as SPS is the major cause of vaginal discharge in canines, occurring in 10%–20% of postpartum bitches from 6 weeks post partum onwards.⁹

In terms of pathophysiology, normal physiological changes in spiral arteries happened during the pregnancy. From early gestation (6–9 weeks), extravillous cytotrophoblasts invade the decidual part of the maternal spiral arteries, replacing the endothelium and it is completed in second trimester when the endothelial lining of the artery is completely replaced. This process causes vascular remodelling with destruction of the musculoelastic medial tissues of the arteries which are replaced by fibrinoid material.¹⁰ These changes play an important role during pregnancy providing an adequate uteroplacental arterial flow. It is well known that altered remodelling of the spiral arteries cause preeclampsia. From this point of view, the SPS vessels can represent the opposite end of a spectrum of abnormal interactions found in preeclampsia.¹¹

In the third trimester, physiologic mechanism of involution of altered uteroplacental vessels begins, as the endovascular cytotrophoblast is replaced by maternal-derived endothelial cells.⁷ Uteroplacental arteries undergo thrombosis and shrinkage of the placental site. Contraction of the uterine smooth muscle also contributes to mechanical shrinkage and involution of these vessels and the uterus revert to the nonpregnant state.¹² In subinvolution of the placental implantation site, the normal remodelling of the uteroplacental arteries during the third trimester is either delayed or inadequate, resulting in the persistence of low-resistance dilated vessels with increased flow.³ In SPS, the uterus is grossly enlarged. Histologically, subinvolved placental bed vessels in the superficial myometrium are large, patent and dilated, with intravascular thrombosis (figure 1). Their walls in microscopic section are distorted and thickened due to deposition of hyaline with partial absence of the endothelial lining.^{3 13}

The exact pathogenesis of SPS is still unknown. There are some factors that may play a role in this process. Sasagawa *et al*,¹⁴ was the first who suggest that immunological patterns are essential for the mechanism of normal involution of uteroplacental arteries, but this study did not explain the exact pathophysiology of subinvolution of these arteries. Later, Andrew *et al*¹⁵ found

the absence of deposition of immunoglobulins (IgG, IgA, IgM) and complement proteins (C1q, C3d, C4) in the walls of subinvolved vessels. Another theory of pathophysiology of subinvolution consist of the increased expression of bcl-2 oncoprotein, which is associated with inhibition of apoptosis and prolonged cell survival in subinvolved placental bed vessels.¹⁶ Bcl-2 oncoprotein is normally not detected in the uteroplacental arteries during the third trimester of pregnancy, however, can be identified in involution and, more strongly, in subinvolution of the placental bed vessels. The review of the literature also revealed the involvement of some nuclear factors identified in controlling human trophoblast invasion, such as Signal Transducer and Activator of Transcription factors (STATs), Peroxisome Proliferator-Activated Receptor gamma (PPAR- γ), homeobox genes or Wntless and Int-1 (WNT)-dependent transcription factors.¹⁷ These factors may contribute to the pathogenesis of pregnancy diseases with abnormal placentation or failed trophoblast differentiation. Epithelial markers such as Troma 1 and CAM 5.2 almost always highlight the trophoblasts while placental alkaline phosphatase, human placental lactogen and epithelial membrane antigen are not pathognomonic.¹⁸

SPS is an important and clinically elusive, diagnosis, as this mechanism responsible for an idiopathic and not iatrogenic cause of postpartum uterine bleeding.¹⁸ Farley and Kohlmeier¹⁹ described the death of a 19-year-old woman in USA due to SPS. Eight days after delivery, her husband brought her to the emergency room pulseless and without respirations. The diagnosis of SPS usually is under-recognised and made by excluding other causes of persistent secondary PPH. The sonographic features of subinvolution of the placental site, through visualising the low-resistance vessels present along the inner third of the myometrium can assist in the diagnosis of SPS.⁶ Pulsed wave Doppler sonography may confirm an increased PSV with a low-resistance waveform which, given that high flow sonographic findings are commonly a result of true arteriovenous vascular malformations (AVMs), that may be acquired (eg, iatrogenic trauma leading to scar tissue and AVM formation) or congenital (eg, developmental vascular defects leading to a bed of abnormal arterio-venous communications).^{20–25} While SPS diagnosis is only solidified via histological analysis, a high degree of suspicion from the clinical history such as abnormal uterine bleeding or recurrent loss of pregnancy may prompt the clinician to request imaging. Contrast-enhanced MRI scans could confirm the marked vascular dilatation of the uterine wall, helping us to suspect SPS before the hysterectomy.^{23 24} Angiography is considered the gold standard in AVM and possibly SPS diagnosis. SPS symptomatology may increase under hormonal changes such as menstruation or pregnancy, which may raise further suspicion.²⁵

The management of subinvolution of the placental site with bleeding in symptomatic patients remains controversial. It is an emergent situation and full work-up should be performed to exclude any other possible cause of secondary PPH, as well as an immediate stabilisation of the patient before the development of disseminated intravascular coagulopathy. In most cases, curettage is performed to exclude retained placental products even if US examination fails to reveal visible abnormality. Conservative approach with uterine tamponade using gauze or various balloons (Bakri) and uterotonic treatment is acceptable. However, surgical treatments such as ligation of the uterine vessels and hysterectomy are considered as standard therapies.^{26 27} According to the literature, percutaneous embolisation of the uterine artery is considered as an alternative therapy of SPS before hysterectomy.^{4 26–28} This procedure provides fertility preservation, but requires special equipment and ready team of

medical experts. Therefore, even these days, unfortunately, not in all hospitals arterial embolization can be performed.

In conclusion, SPS is one of the rare but important cause of secondary PPH and as other AVMs, it may coexist with other causes of PPH such as retained Products Of Conception (POC).²⁹ This condition is likely under recognised by gynaecologist due to the absence of specific diagnostic criteria, as in our case. Only excluding other forms of delayed PPH and after failed curettage and conservative therapy, the diagnosis of SPS should be included in the differential diagnosis. MRI scan with contrast can indicate the focal areas of uterine hypervascularity. However, the histological confirmation of dilated, clustered, myometrial arteries, partially occluded by thrombi is done only after the urgent hysterectomy.

Learning points

- The incidence of postpartum haemorrhage from subinvolution is most common in the second week post partum. Subinvolution of the placental bed was noted in 13.3% of 26.023 patients, while the placental retention was the most common cause of haemorrhages (30.0%).
- The exact molecular mechanism of the pathogenesis of subinvolution of placental site is still unknown but theories include increased local expression of bcl-2 oncoprotein, absence of deposition of immunoglobulins and complement proteins in the walls of subinvolved vessels, STATs, PPAR- γ , homeobox genes and WNT-dependent transcription factor deregulation.
- Unfortunately, subinvolution still remains a histological diagnosis and no imaging or other markers are available to identify patients' prior major complications arise.

Contributors Concept, analysis, format, revision, editing, literature search and drafted the manuscript: IM, DK, OT, SK; Literature search, extraction analysis and drafted the manuscript: OT, SK.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Stavroula Kastora <http://orcid.org/0000-0003-4470-5267>

REFERENCES

- 1 Carroli G, Cuesta C, Abalos E, *et al*. Epidemiology of postpartum haemorrhage: a systematic review. *Best Pract Res Clin Obstet Gynaecol* 2008;22:999–1012.
- 2 Khan KS, Wojdyla D, Say L, *et al*. WHO analysis of causes of maternal death: a systematic review. *Lancet* 2006;367:1066–74.
- 3 Akgun H, Gonlusen G, Cartwright J, *et al*. Are gadolinium-based contrast media nephrotoxic? a renal biopsy study. *Arch Pathol Lab Med* 2006;130:1354–7.
- 4 Küster H. Ueber Spätblutungen im Wochenbett. Neue Gesichtspunkte zur Beurteilung ihrer Aetiologie nach mikroskopischen Untersuchungen. *Z Geburtsh Gynaek* 1910;67:430–4.
- 5 Rutherford RN, Hertig AT. Noninvolution of the placental site. *Am J Obstet Gynecol* 1945;49:378–84.
- 6 Petrovitch I, Beatty M, Jeffrey RB, *et al*. Subinvolution of the placental site. *J Ultrasound Med* 2009;28:1115–9.
- 7 Andrew AC, Bulmer JN, Wells M, *et al*. Subinvolution of the uteroplacental arteries in the human placental bed. *Histopathology* 1989;15:395–405.

- 8 Dossou M, Debost-Legrand A, Déchelotte P, *et al.* Severe secondary postpartum hemorrhage: a historical cohort. *Birth* 2015;42:149–55.
- 9 Ma K. Canine Postpartum Disorders. In: Jd B, Dc T, eds. *Kirk's Current Veterinary Therapy Xiv*. 14th edn. St Louis, Mo: Saunders, 2000: 999–1002.
- 10 Gerretsen G, Huisjes HJ, Hardonk MJ, *et al.* Trophoblast alterations in the placental bed in relation to physiological changes in spiral arteries. *Br J Obstet Gynaecol* 1983;90:34–9.
- 11 Meekins JW, Pijnenborg R, Hanssens M, *et al.* A study of placental bed spiral arteries and trophoblast invasion in normal and severe pre-eclamptic pregnancies. *BJOG* 1994;101:669–74.
- 12 Anderson WR, Davis J. Placental site involution. *Am J Obstet Gynecol* 1968;102:23–33.
- 13 Wells M, Bulmer JN. The human placental bed: histology, immunohistochemistry and pathology. *Histopathology* 1988;13:483–98.
- 14 Sasagawa M, Watanabe S, Ohmomo Y, *et al.* Reactivity of two monoclonal antibodies (Troma 1 and CaM 5.2) on human tissue sections: analysis of their usefulness as a histological trophoblast marker in normal pregnancy and trophoblastic disease. *Int J Gynecol Pathol* 1986;5:345–56.
- 15 Andrew A, Bulmer JN, Morrison L, *et al.* Subinvolution of the uteroplacental arteries: an immunohistochemical study. *Int J Gynecol Pathol* 1993;12:28–33.
- 16 Khong TY, Abdul Rahman H, Rahman A. Bcl-2 expression delays postpartum involution of pregnancy-induced vascular changes in the human placental bed. *Int J Gynecol Pathol* 1997;16:138–42.
- 17 Knöfler M. Critical growth factors and signalling pathways controlling human trophoblast invasion. *Int J Dev Biol* 2010;54:269–80.
- 18 Kavalir R, Arko D, Fokter Dovnik N, *et al.* Subinvolution of placental bed vessels: case report and review of the literature. *Wien Klin Wochenschr* 2012;124:725–30.
- 19 Farley NJ, Kohlmeier RE. A death due to subinvolution of the uteroplacental arteries. *J Forensic Sci* 2011;56:803–5.
- 20 Hoffman MK, Meilstrup JW, Shackelford DP, *et al.* Arteriovenous malformations of the uterus: an uncommon cause of vaginal bleeding. *Obstet Gynecol Surv* 1997;52:736–40.
- 21 Vogelzang RL, Nemcek AA, Skrtic Z, *et al.* Uterine arteriovenous malformations: primary treatment with therapeutic embolization. *J Vasc Interv Radiol* 1991;2:517–22.
- 22 Huang MW, Muradali D, Thurston WA, *et al.* Uterine arteriovenous malformations: gray-scale and Doppler US features with MR imaging correlation. *Radiology* 1998;206:115–23.
- 23 Godoy LL, Torres US, D'Ippolito G. Subinvolution of the placental site associated with focal retained products of conception and placenta accreta mimicking uterine arteriovenous malformation on CT and MRI: a lesson to be learned. *Radiol Bras* 2018;51:135–6.
- 24 Sellmyer MA, Desser TS, Maturen KE, *et al.* Physiologic, histologic, and imaging features of retained products of conception. *Radiographics* 2013;33:781–96.
- 25 Aziz N, Lenzi TA, Jeffrey RB, *et al.* Postpartum uterine arteriovenous fistula. *Obstet Gynecol* 2004;103:1076–8.
- 26 Maleux G, Timmerman D, Heye S, *et al.* Acquired uterine vascular malformations: radiological and clinical outcome after transcatheter embolotherapy. *Eur Radiol* 2006;16:299–306.
- 27 Cheng Y-Y, Hwang J-I, Hung S-W, *et al.* Angiographic embolization for emergent and prophylactic management of obstetric hemorrhage: a four-year experience. *J Chin Med Assoc* 2003;66:727.
- 28 Chauleur C, Fanget C, Tourne G, *et al.* Serious primary post-partum hemorrhage, arterial embolization and future fertility: a retrospective study of 46 cases. *Hum Reprod* 2008;23:1553–9.
- 29 I raha Y, Okada M, Toguchi M, *et al.* Multimodality imaging in secondary postpartum or postabortion hemorrhage: retained products of conception and related conditions. *Jpn J Radiol* 2018;36:12–22.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow