OPEN ACCESS

False elevations in urinary metanephrines: under-recognised pitfall with 24-hour urinary volume collection

Terry Shin,^{1,2} Thanh Duc Hoang,^{2,3} Mary Thomas Plunkett,¹ Mohamed K M Shakir^{2,3}

¹Endocrinology, Womack Army Medical Center, Fort Bragg, North Carolina, USA

²Uniformed Services University of the Health Sciences, Bethesda, MD, USA

³Endocrinology, Walter Reed National Military Medical Center, Bethesda, Maryland, USA

Correspondence to

Dr Terry Shin;
terrytshin@gmail.com

Accepted 13 January 2021

SUMMARY

One pitfall in 24-hour urine collection is the input of incorrect urinary volume by the reference laboratory. This may lead to an incorrect diagnosis of pheochromocytoma or paraganglioma. A 48-year-old African-American woman was seen in the clinic for an elevated 24-hour urinary metanephrine screen during workup for secondary hypertension. Urine volume was found to be incorrectly inputted by the lab as 9750 mL rather than 975 mL. The urinary metanephrines were then recalculated and the 24-hour urinary metanephrines resulted within normal limits. This case highlights this unique and potentially under-recognised error in testing with 24-hour urine volume collection.

BACKGROUND

Since 1956, when Armstrong *et al* reported the presence of catecholamine products in the urine, testing methodologies have gradually improved in detecting pheochromocytoma and paraganglioma (PPGL).¹ The measurement of fractionated 24-hour urinary metanephrines is now one of the standard first line tests to detect PPGL.² Detection accuracy and precision of urinary metanephrines have improved further with implementation of mass spectrometry.³ Even with improved accuracy in analytical results, the 24-hour urine collection in the real world can be fraught with errors. Specifically, errors in urine results can occur in the pre-analytical phase, defined as the errors that occur from physician order to the point of laboratory testing.⁴

These errors in laboratory medicine can occur despite dedicated technicians with high-quality processes in place to avoid such errors. The frequency of laboratory errors is reported at a rate of 0.012%–0.6%.⁵ Since 80%–90% of the clinical decision making is based on labs, this small percentage can substantially affect patient care.⁵ Clinician's should be cognizant of this possibility and this case highlights the errors that can occur with 24-hour urine collection and approaches to avoid these diagnostic pitfalls.

CASE PRESENTATION

A 48-year-old African-American woman was referred to the endocrine clinic for treatment of PPGL. Endocrine workup for thyroid disorder and hyperaldosteronism was ruled out, but she was found to have marked elevation in 24-hour urine metanephrines. She was diagnosed with PPGL and underwent further evaluation with CT scan of the

adrenals that did not identify any adrenal nodules or abnormalities.

She was referred to the endocrine clinic for further evaluation. Her workup was notable for two episodes of hypertensive urgency where she required treatment in the emergency room. She denied any symptoms of pheochromocytoma. She was on metoprolol, valsartan and chlorthalidone for blood pressure control. No other significant medical conditions reported. She denied any family history of PPGL, early heart disease or stroke. Physical examination was overall unremarkable. Her laboratory data reported marked elevation in urinary metanephrines at 995 µg/24 hours (normal 45–290 µg/24 hours) and normetanephrine at 3393 µg/24 hours (normal 80–500 µg/24 hours). However, both 24-hour urinary and plasma metanephrines were repeated and resulted within normal limits.

OUTCOME AND FOLLOW-UP

Due to the significant discrepancy in results, her prior urinary samples were scrutinised. The reference laboratory was contacted, and it was discovered that the measurements were calculated with a urinary volume of 9750 mL. After querying the patient, she reported a urinary volume of approximately 1 L. It was deduced that the actual urinary volume was 975 mL and not 9750 mL. The urinary metanephrines was recalculated and the 24-hour urinary metanephrines resulted within normal limits at 99 µg/24 hours for metanephrines and 339 µg/24 hours for normetanephrines. The patient was reassured of results and blood pressure was optimised with adjustment in antihypertensive medications. No further medical issues were reported.

DISCUSSION

The measurement of fractionated 24-hour urinary or plasma metanephrines (metanephrines and normetanephrines) is the standard first-line screen for PPGL. There is controversy on which is considered the best initial screening test.² Plasma metanephrines have a higher sensitivity (96%–100%) but it is less specific than 24-hour metanephrines (77%–89%).⁶ Twenty-four-hour urine metanephrines have a sensitivity of 97% and a specificity of 85%–91.1%.^{6,7} Measurement of 24-hour metanephrines combined with 24 hour urine catecholamines increases the specificity to 98%.⁷ Sawka *et al* at Mayo Clinic has proposed measurement of the two methods based on pretest probability of PPGL.⁶

© BMJ Publishing Group Limited 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Shin T, Hoang TD, Plunkett MT, *et al*. *BMJ Case Rep* 2021;**14**:e241147. doi:10.1136/bcr-2020-241147

Case report

In patients with high suspicion for PPGL, the recommendation is plasma metanephrines.⁶ In patients with low pretest probability of PPGL, the recommendation is 24-hour urine metanephrines. Though this is the standard recommended practice, institutions often use plasma metanephrines as an initial screening test because 24-hour urine collection can be fraught with errors and burdensome to patients.²

Preamalytical errors in collection of 24-hour urine is quite frequent, with studies reporting a collection accuracy of 50%–51% $p=0.017$.^{8–10} Undercollection or overcollection of 24-hour urine is the most common error in urinary collection accuracy.⁸ Excessively high urine volumes will lead to false elevation, and low urine volume collection will conversely lead to falsely low results. Despite detailed instructions, one-third of the patients changed their fluid intake habits to increase urinary volume.¹¹ Another complicating factor is the possibility of using the wrong urine preservative for the 24 hours urine collection. Interestingly, in our patient's case, the error occurred with actual input of laboratory data. With urine collection, the volume must be manually entered by the laboratory technician to calculate the 24-hour urine laboratory result. The frequency of errors that occur from incorrect technician input is unknown, but this specific type of error is believed to be a unique cause of an abnormal 24-hour urine result. Given that the laboratory errors from a human factor is high as 80%, it is possible that this error may occur more frequently than recognised.¹² To avoid this possibility, the clinician should always review and confirm that the urine volume is appropriate for the 24-hour collection. By using basic stoichiometry, we were able to calculate the actual patient's urinary metanephrines based on the aliquot and corrected urinary volume.

Due to issues associated with 24-hour urine collection, guidelines for PPGL recommend measurement of 24-hour urinary excretion creatinine to verify completeness. Collecting creatinine simultaneously with metanephrines can help confirm adequacy. Daily creatinine excretion is approximated at 20–25 mg/kg of lean body weight in males and 15–20 mg/kg of lean body weight in females up to 50 years of age.¹³ There is a gradual decline in creatinine beyond 50 years of age due to loss of muscle mass. Other formulas are available that take race, sex and serum phosphate into account in estimating creatinine.¹⁴ Another study has validated use of a creatinine ratio formula to correct for overcollection and undercollection of 24-hour urine results.¹⁵ Though measured 24-hour creatinine cannot be assumed to guarantee accurate measurement, it can help as a surrogate in assessing collection accuracy.¹⁴

A 24-hour urine collection will likely remain the standard for PPGL screening, but there have been alternative collection strategies proposed in detecting urinary metanephrines. Since catecholamines have a diurnal excretion, morning urine void was shown to be effective with similar diagnostic sensitivity and specificity as 24-hour urine collection.^{16 17} Other studies have proposed the potential of spot urine metanephrine to creatinine ratios via direct electrochemical measurement.^{16 18 19}

In summary, a 24-hour urine collection is an important screening test to diagnose PPGL. Clinicians must be able to identify potential errors in a 24-hour calculation that occurs with urine collection. Identifying these errors will reduce diagnostic uncertainty and inappropriate treatment. This not only applies to detection of PPGL in urinary metanephrine collection, but also applicable to all 24-hour urine collection samples for other medical conditions.

Contributors All listed authors (TS, TDH, MTP and MKMS) were involved with authoring and reviewing this case report.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Disclaimer The authors are employees of the United States Government. This work was prepared as part of their official duties. Title 17 U.S.C. §105 provides that "Copyright protection under this title is not available for any work of the United States Government." Title 17 U.S.C. § 101 defines a U.S. Government work as a work prepared by a military service member or employee of the U.S. Government as part of that person's official duties. The identification of specific products or scientific instrumentation is considered an integral part of the scientific endeavor and does not constitute endorsement or implied endorsement on the part of the author, DoD, or any component agency. The views expressed in this article are those of the author and do not reflect the official policy of the Department of Defense or the U.S. Government.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Armstrong MD, Shaw KN, Wall PE. The phenolic acids of human urine; paper chromatography of phenolic acids. *J Biol Chem* 1956;218(1):293–303.
- Lenders JWM, Duh Q-Y, Eisenhofer G, et al. Pheochromocytoma and paraganglioma: an endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2014;99:1915–42.
- Eisenhofer G, Peitzsch M, Kaden D, et al. Reference intervals for plasma concentrations of adrenal steroids measured by LC-MS/MS: impact of gender, age, oral contraceptives, body mass index and blood pressure status. *Clin Chim Acta* 2017;470:115–24.
- Abdollahi A, Saffar H, Saffar H. Types and frequency of errors during different phases of testing at a clinical medical laboratory of a teaching hospital in Tehran, Iran. *N Am J Med Sci* 2014;6:224–8.
- O'Kane M. The reporting, classification and grading of quality failures in the medical laboratory. *Clin Chim Acta* 2009;404:28–31.
- Sawka AM, Jaeschke R, Singh RJ, et al. A comparison of biochemical tests for pheochromocytoma: measurement of fractionated plasma metanephrines compared with the combination of 24-hour urinary metanephrines and catecholamines. *J Clin Endocrinol Metab* 2003;88:553–8.
- Perry CG, Sawka AM, Singh R, et al. The diagnostic efficacy of urinary fractionated metanephrines measured by tandem mass spectrometry in detection of pheochromocytoma. *Clin Endocrinol* 2007;66:703–8.
- Boyd C, Wood K, Whitaker D, et al. Accuracy in 24-hour urine collection at a tertiary center. *Rev Urol* 2018;20:119–24.
- Sawyer MD, Dietrich MS, Pickens RB, et al. Adequate or not? A comparison of 24-hour urine studies for renal stone prevention by creatinine to weight ratio. *J Endourol* 2013;27:366–9.
- McGuire BB, Bhanji Y, Sharma V, et al. Predicting patients with inadequate 24- or 48-hour urine collections at time of metabolic stone evaluation. *J Endourol* 2015;29:730–5.
- Miller M, Simundić A-M. Low level of adherence to instructions for 24-hour urine collection among hospital outpatients. *Biochem Med* 2013;23:316–20.
- Szecs PB, Ødum L. Error tracking in a clinical biochemistry laboratory. *Clin Chem Lab Med* 2009;47:1253–7.

Learning points

- ▶ A 24-hour urinary metanephrines may be falsely abnormal due to incorrect input of urinary volume.
- ▶ The clinician should review and confirm the urinary volume in all patients that complete a 24-hour urine volume collection to ensure results are accurate.
- ▶ The clinician may reduce uncertainty of appropriate 24-hour urine collection by pairing the collection with 24-hour urinary creatinine.

- 13 Walser M. Creatinine excretion as a measure of protein nutrition in adults of varying age. *JPEN J Parenter Enteral Nutr* 1987;11:735–8.
- 14 Ix JH, Wassel CL, Stevens LA, *et al.* Equations to estimate creatinine excretion rate: the CKD epidemiology collaboration. *Clin J Am Soc Nephrol* 2011;6:184–91.
- 15 Mann SJ, Gerber LM. Addressing the problem of inaccuracy of measured 24-hour urine collections due to incomplete collection. *J Clin Hypertens* 2019;21:1626–34.
- 16 Peitzsch M, Kaden D, Pamporaki C, *et al.* Overnight/first-morning urine free metanephrines and methoxytyramine for diagnosis of pheochromocytoma and paraganglioma: is this an option? *Eur J Endocrinol* 2020;182:499–509.
- 17 Peaston RT, Lennard TW, Lai LC. Overnight excretion of urinary catecholamines and metabolites in the detection of pheochromocytoma. *J Clin Endocrinol Metab* 1996;81:1378–84.
- 18 Shi Z-H, Zhang X-Q, Zhen Q-N, *et al.* Direct electrochemical measurement of metanephrines in spot urine samples for the diagnosis of phaeochromocytomas. *Sci Rep* 2017;7:8041.
- 19 Zuo M, Zhen Q, Zhang X, *et al.* High specificity of spot urinary free metanephrines in diagnosis and prognosis of pheochromocytomas and paragangliomas by HPLC with electrochemical detection. *Clin Chim Acta* 2018;478:82–9.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow