OPEN ACCESS

Postoperative lumbar paraspinal compartment syndrome

 Niloufar Saadat ,¹ Kourosh Rezania ²

¹Department of Radiology,
University of Chicago, Chicago,
Illinois, USA

²Department of Neurology,
University of Chicago, Chicago,
Illinois, USA

Correspondence to
Dr Kourosh Rezania;
krezania@uchicago.edu

Accepted 12 October 2020

SUMMARY

Lower lumbar paraspinal muscles constitute a compartment as they are surrounded by distinct fascial and bony boundaries. Lumbar paraspinal compartment syndrome is a rare entity, often caused by intense exercise, but also can be a postoperative complication. We present a 60-year-old man with low back pain, numbness in the left lower back and radicular pain in the left lower extremity, which started after a surgery that involved prolonged positioning on the left side 7 years before, and persisted to the day of evaluation. There was an immediate transient rise in the creatine kinase after surgery. Electromyography showed a left lower lumbar–sacral plexopathy and a lumbar spine MRI revealed fatty infiltration of the lower lumbar–sacral paraspinal muscles. The emergence of radicular lower limb pain was likely due to the compression of the proximal portion of lumbar–sacral plexus during the acute stage of rhabdomyolysis.

BACKGROUND

Lumbar paraspinal compartment syndrome is a rare cause of rhabdomyolysis and of chronic low back pain. A universal feature is high creatine kinase (CK) (generally >5000 IU/L) and rhabdomyolysis.¹ Characteristic symptomatology includes local paraspinal muscle pain exacerbated by passive movements, that is, lateral truncal tilting, forward flexion and straight leg raising, and alleviated with the extension of the back; and tenderness and firm texture/board-like rigidity of the affected paraspinal musculature on palpation.^{1–3} Timely diagnosis and appropriate surgical treatment will usually result in favourable outcomes.

CASE PRESENTATION

Our patient is a 60-year-old man with a history of diabetes and morbid obesity who presented with chronic low back pain, left lower back numbness, radicular pain, numbness in the left leg and lateral foot that started after surgery on the right ankle, for debridement of a wound and skin grafting 7 years before, the symptoms persisted to the day of evaluation. During the surgery, he was positioned on the left side. He developed rhabdomyolysis in the immediate postoperative period with a rise of serum CK to ~5000 U/L from a preoperative level of 188 U/L on postoperative day 1, increasing to a peak of ~20 000 U/L on postoperative day 3, which trended back to normal over 2 weeks. Initial lumbar spine MRI, by report, did not show significant abnormalities except for spinal osteoarthritis. He was treated with gabapentin, tramadol and

underwent epidural steroid injections for his refractory back and lumbar radiculopathy pain, which only provided partial relief of symptoms. Physical examination revealed decreased sensation in the left lower flank, lower leg and lateral foot and absent left ankle reflex.

INVESTIGATIONS

Nerve conduction studies showed mildly low amplitude of the left sural sensory action potential, normal amplitude and mild slowing of the left peroneal motor nerve action potential (recorded at extensor digitorum brevis), and normal tibial motor responses as well as F wave latencies.

Needle electromyography showed neurogenic units and recruitment in the left gastrocnemius and reduced insertional activity with fibrillation potentials and positive waves in the left lower lumbar paraspinal muscles. The study was consistent with a left lower lumbar–sacral radiculopathy and a concomitant distal axonal neuropathy versus a lumbar–sacral plexopathy. MRI revealed severe atrophy and adipose infiltration of left-sided lower lumbar–sacral paraspinal muscles extending from the L3 to L4 level inferiorly through the sacral levels (large arrows in [figure 1A, B](#)).

DIFFERENTIAL DIAGNOSIS

Lumbar paraspinal compartment syndrome is a very rare cause of low back and radicular lower limb pain. More common conditions include different pathological processes caused by degenerative arthritis of the lumbar spine, including spondylolisthesis, intervertebral disk bulging and herniation, and facet joint arthropathy. Neoplastic and inflammatory infiltration of the nerve roots and plexi are other less common causes of radiculopathy and plexopathy. Another consideration in our case who had diabetes and radiculopathy symptoms in the left lower limb would have been diabetic lumbosacral radiculoplexus neuropathy, a monophasic entity that can present in even well-controlled cases of type 2 diabetes.⁴

OUTCOME AND FOLLOW-UP

The patient has been followed-up for a period of 4 years after his initial neurological evaluation. His back and radicular lower limb pain have partially responded to oral gabapentin. Duloxetine was temporarily tried and stopped due to a lack of efficacy. He has also undergone left L3 and L4 transforaminal epidural steroid injections on three occasions during the last 3 years of follow-up ([figure 2](#)).

© BMJ Publishing Group Limited 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Saadat N, Rezania K. *BMJ Case Rep* 2021;**14**:e236040. doi:10.1136/bcr-2020-236040

Figure 1 Sagittal T1-weighted image (A) and axial T2-weighted image (B) demonstrate adipose infiltration of the lumbar-sacral paraspinal muscles (large arrow), intact fascial boundaries of the lumbar paraspinal compartment (arrowheads) and proximity of the affected paraspinal muscles to the very proximal part of lumbar-sacral plexus, that is, ventral rami of the nerve roots (small arrow).

DISCUSSION

The chronic lower back and radicular pain in the left lower limb in our patient started after prolonged positioning on his left side 7 years before evaluation, and there was elevated CK in the postoperative period that trended to normal in 2 weeks (figure 2). Fatty infiltration of the left lower lumbar paraspinal compartment syndrome as the cause of postoperative transient rhabdomyolysis and subsequent emergence of the chronic lower back pain.

Lower lumbar and sacral paraspinal muscles constitute a compartment as they are surrounded by well-developed fascial sheaths (arrowheads in figure 1A, B) and bony structures. The lumbar paraspinal compartment is contained by lamellae of the fascial sheet posteriorly that merge with the spinous process and interspinous ligaments medially, the fascia of abdominal muscles laterally, iliac crest and sacrum caudally and the transverse spine processes ventrally.^{5 6}

At least 30 previous cases of lumbar paraspinal compartment syndrome have been previously reported (table 1); with an age range of 16–67, predominantly male individuals (M/F: 28/3). This is a condition that predominantly affects athletic male individuals in the second to fourth decades, with strenuous exertion usually involving weightlifting being the cause in the majority of cases.^{2 5 7–12} Other less common culprits include direct trauma due to falls^{13 14} and use of vasoconstrictors cocaine, ephedrine and pseudoephedrine.^{2 3 11 15} Only five (including our) cases are reported following surgical procedures.^{16–18} Diagnosis of lumbar paraspinal compartment syndrome should be confirmed

by direct measurement of intramuscular pressure in the acute stage.^{3 5 10 12 13 19} Changes in the MRI and CT scan in the acute stage consist of swelling of the paraspinal muscles and abnormal signal intensity in T2-weighted MRI images.^{2 10 13 18} The aforementioned changes can be subtle and be missed in the early stage.¹⁸ When undiagnosed, myonecrosis and concomitant nerve injury may result in chronic back pain and persistent numbness in the lumbar/sacral dermatomes.^{1 2 7 17 18} Lower back numbness and paraesthesia are caused by the involvement of dorsal rami of the nerve roots and cutaneous branches of the cluneal nerves.¹⁵

Radicular pain, sensory symptoms and absent reflexes in the lower limbs have previously been reported following lumbar paraspinal compartment syndrome.^{15 18 20} In contrast, patients may present with loin or groin pain, suggesting renal colic.^{21 22} The emergence of radicular lower limb pain in our patient can be explained by the compression of the anterior divisions (ventral rami) of the lower lumbar-sacral nerve roots, because of oedema in the paraspinal muscles associated with the acute stage of rhabdomyolysis. Although the anterior divisions course ventrally after exiting from the neural foramina, they are separated from the erector spinae only by fascial planes at certain points during their course (small arrow in figure 1B⁶). In contrast, another possible explanation of development of left lower limb plexopathy in our case would have been traction injury to the lower lumbar-sacral plexus due to positioning, unrelated to rhabdomyolysis.

Similar to the compartment syndromes involving the extremities, treatment of acute lumbar paraspinal compartment syndrome consists of surgical decompression (fasciotomy), which may improve the long term outcome.^{1 8–10 13} Sixteen of the 29 previously reported cases underwent fasciotomies, 14 of whom had complete recovery of back pain days to weeks after the procedure (table 1). In contrast, the recovery was more gradual, over a period of 4–6 months in patients who had conservative treatment (management of pain, intravenous hydration and monitoring of kidney function). The favourable outcome when defined as complete recovery or mild residual symptoms was reported in only 6 of 14 of the conservatively treated patients, whereas another 6 patients, including our patient, had significant chronic low back pain when followed-up for 2 months to 8 years after the incident (table 1). Optimal surgery timelines have not been established with regard to lumbar paraspinal compartment syndrome given the rarity of this condition¹³; decompression fasciotomies have been done 2–3 days after the onset of symptoms with good long-term outcomes.^{8 11 23}

Figure 2 Timeline of events and treatments, timing of surgery, which preceded rhabdomyolysis, and of the lumbar MRI study are presented by arrows. Timing of treatment with gabapentin and duloxetine are demonstrated with grey and striped bars, respectively. Epidural injections are shown by arrowheads. CK, creatine kinase.

Table 1 Previously reported cases of lumbar paraspinal compartment syndrome

Reference	Age/sex	Setting	Fasci.	Outcome/duration of follow-up
Present case	53/M	Ankle surgery (prolonged positioning), obesity	–	Chronic back pain, radicular pain in the lower limb/7 years
⁵	24/M	Exercise (skiing)	–	Intermittent residual back pain/3 months
⁷	27/M	Exercise (skiing)	–	Mild residual back pain/4 months
¹⁶	55/M	Abdominal aortic aneurysm surgery	–	Slight residual back pain/3 months
¹⁰	21/M	Weightlifting	+	Significant improvement of back pain/7 days
¹⁸ Case 1	57/M	Aortoiliac bypass surgery	–	Chronic low back pain/10 months
¹⁸ Case 2	34/M	Gastric bypass surgery	–	Sensory loss in the peri-spinal region/1 year
¹³	43/F	Trauma, fall	+	Significant improvement in back pain/6 months
⁹	35/M	Exercise (skiing)	+	Complete recovery/3 months
⁸	25/M	Exercise (surfboarding)	+	Complete recovery/2 months
¹⁵	29/F	Weightlifting	–	Resolution of back pain/6 months
¹⁹	16/M	Weightlifting	+	Complete resolution of back pain/2 years
¹⁷	67/M	abdominal aortic surgery	–	Residual mild back pain/7 months
³	45/M	Cocaine	+	Complete recovery/2 weeks
²	30/M	Bodybuilding, anabolic steroids	–	Significant residual back pain/2 months
¹¹	32/M	creatine, pseudoephedrine	+	Complete recovery/14 days
²⁴	23/M	Weightlifting	–	Residual back pain only with exercise/4 months
¹²	20/M	Strenuous exercise	+	Complete recovery/1 months
²⁵	16/M	Strenuous exercise	+	Complete recovery/3 months
²³	30/M	Weightlifting	+	Complete recovery/3 months
²⁶	23/M	Exercise	+	Complete recovery/1 year
¹⁴	43/F	Direct trauma	+	Post-operative recovery/NA
²²	25/M	Weightlifting	–	Residual back pain/13 days
²⁷	23/M	Weightlifting	+	Complete recovery/1 year
²⁸	25/M	Weightlifting	–	Persistent back pain/3 days
²⁰	25/M	Exercise (cross-fit)	–	Refractory back pain/5 days
²¹	24/M	Weightlifting	–	Persistent back pain/9 days

Continued

Table 1 Continued

Reference	Age/sex	Setting	Fasci.	Outcome/duration of follow-up
²⁹ Case1	30/M	Weightlifting	+	Complete recovery/6 months
²⁹ Case 2	31/M	Exercise	+	Complete recovery/6 months
³⁰	17/M	Exercise (callisthenics)	+	Complete recovery/4 months

F, female; Fasci, fasciotomy; M, male.

Learning points

- Lumbar paraspinal compartment syndrome is a rare cause of chronic lower back pain and radicular pain in the lower limb.
- It is most commonly reported in younger male individuals after weightlifting exercises.
- It is also a postoperative complication, such as prolonged positioning on the side.
- This condition has to be considered in patients with rhabdomyolysis and back pain after heavy exercise or surgical procedures.
- Surgical intervention in the acute stage may prevent chronic back pain.

Acknowledgements We acknowledge the patient who graciously agreed with and consented to this publication and Dr Javad Hekmat-Panah who reviewed and commented on the manuscript.

Contributors The manuscript was written and edited by NS and KR. KR revised the manuscript for important intellectual content.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Niloufar Saadat <http://orcid.org/0000-0001-8152-599X>
Kourosh Rezanian <http://orcid.org/0000-0003-3986-5709>

REFERENCES

- Nathan ST, Roberts CS, Deliberato D. Lumbar paraspinal compartment syndrome. *Int Orthop* 2012;36:1221–7.
- Wik L, Patterson JM, Oswald AE. Exertional paraspinal muscle rhabdomyolysis and compartment syndrome: a cause of back pain not to be missed. *Clin Rheumatol* 2010;29:803–5.
- Rogers ME, Lowe JA, Vanlandingham SC. Acute erector spinae compartment syndrome: case report and review of diagnostic criteria. *Injury* 2014;45:813–5.
- Dyck PJB, Windebank AJ. Diabetic and nondiabetic lumbosacral radiculoplexus neuropathies: new insights into pathophysiology and treatment. *Muscle Nerve* 2002;25:477–91.
- Carr D, Gilbertson L, Frymoyer J, et al. Lumbar paraspinal compartment syndrome. A case report with physiologic and anatomic studies. *Spine* 1985;10:816–20.
- Willard FH, Vleeming A, Schuenke MD, et al. The thoracolumbar fascia: anatomy, function and clinical considerations. *J Anat* 2012;221:507–36.
- DiFazio FA, Barth RA, Frymoyer JW. Acute lumbar paraspinal compartment syndrome. A case report. *J Bone Joint Surg Am* 1991;73:1101–3.

- 8 Kitajima I, Tachibana S, Hirota Y, *et al.* Acute paraspinal muscle compartment syndrome treated with surgical decompression: a case report. *Am J Sports Med* 2002;30:283–5.
- 9 Khan RJK, Fick DP, Guier CA, *et al.* Acute paraspinal compartment syndrome. A case report. *J Bone Joint Surg Am* 2005;87:1126–8.
- 10 Golovko DM, Knox JB. An uncommon cause of low back pain. *Am J Med* 2017;130:e551–2.
- 11 Minnema BJ, Neligan PC, Quraishi NA, *et al.* A case of occult compartment syndrome and nonresolving rhabdomyolysis. *J Gen Intern Med* 2008;23:871–4.
- 12 Paryavi E, Jobin CM, Ludwig SC, *et al.* Acute exertional lumbar paraspinal compartment syndrome. *Spine* 2010;35:E1529–33.
- 13 Harper KD, Phillips D, Lopez JM, *et al.* Acute traumatic thoracolumbar paraspinal compartment syndrome: case report. *J Neurosurg Spine* 2018;30:140–5.
- 14 Wasserman DD, Isenberg DL. A case of acute traumatic lumbar paraspinal compartment syndrome. *J Emerg Med* 2018;55:544–6.
- 15 Lauschke LGM G. Acute paraspinal compartment syndrome related to use of proprietary weight loss product, by a patient with sodium channelopathy. *J Spine* 2016;57.
- 16 Ferreira J, Galle C, Aminian A, *et al.* Lumbar paraspinal rhabdomyolysis and compartment syndrome after abdominal aortic aneurysm repair. *J Vasc Surg* 2003;37:198–201.
- 17 Osamura N, Takahashi K, Endo M, *et al.* Lumbar paraspinal myonecrosis after abdominal vascular surgery: a case report. *Spine* 2000;25:1852–4.
- 18 Haig AJ, Hartigan AG, Quint D. Low back pain after nonspinal surgery: the characteristics of presumed lumbar paraspinal compartment syndrome. *Pm R* 2009;1:383–8.
- 19 Maeckelbergh L, Moke L. An acute paraspinal compartment syndrome: anatomic description of the compartment and surgical technique. *Clin Spine Surg* 2019;32:E277–81.
- 20 Chavez JM, Gonzalez PG. Suspected lumbar compartment syndrome: a rare cause of low back pain after strenuous exercise. *Spine J* 2013;13:1409–10.
- 21 Hoyle A, Tang V, Baker A, *et al.* Acute paraspinal compartment syndrome as a rare cause of loin pain. *Ann R Coll Surg Engl* 2015;97:e11–12.
- 22 Allerton C, Gawthorpe IC. Acute paraspinal compartment syndrome as an unusual cause of severe low back pain. *Emerg Med Australas* 2012;24:457–9.
- 23 Mattiassich G, Larcher L, Leitinger M, *et al.* Paravertebral compartment syndrome after training causing severe back pain in an amateur rugby player: report of a rare case and review of the literature. *BMC Musculoskelet Disord* 2013;14:259.
- 24 Karam MD, Amendola A, Mendoza-Lattes S. Case report: successful treatment of acute exertional paraspinal compartment syndrome with hyperbaric oxygen therapy. *Iowa Orthop J* 2010;30:188–90.
- 25 Vanbrabant P, Moke L, Meersseman W, *et al.* Excruciating low back pain after strenuous exertion: beware of lumbar paraspinal compartment syndrome. *J Emerg Med* 2015;49:641–3.
- 26 Zyskowski M, Schwarz A, Huber-Wagner S. Paraspinal compartment Syndrome-an insidious cause of acute back pain. *Dtsch Arztebl Int* 2019;116:118.
- 27 Alexander W, Low N, Pratt G. Acute lumbar paraspinal compartment syndrome: a systematic review. *ANZ J Surg* 2018;88:854–9.
- 28 Calvert N, Bhalla T, Twerenbold R. Acute exertional paraspinal compartment syndrome. *ANZ J Surg* 2012;82:564–5.
- 29 Rha EY, Kim DH, Yoo G. Acute exertional lumbar paraspinal compartment syndrome treated with fasciotomy and dermatotraction: case report. *J Plast Reconstr Aesthet Surg* 2014;67:425–6.
- 30 Schreiber VM, Ward WT. Exercise-Induced pediatric lumbar paravertebral compartment syndrome: a case report. *J Pediatr Orthop* 2015;35:e49–51.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow