Hair-thread Tourniquet Syndrome causing uvular strangulation in a child

Jessica Leighton ¹, Jason Powell,^{1,2} Claire McLarnon¹

¹Otolaryngology, Newcastle Upon Tyne Hospitals NHS Foundation Trust, Newcastle Upon Tyne, UK

²Translational and Clinical Research Institute, Newcastle University, Newcastle upon Tyne, UK

Correspondence to

Dr Jessica Leighton;
jess.leighton@newcastle.ac.uk

Accepted 25 March 2021

DESCRIPTION

A normally fit and well 3-year-old girl awoke distressed. Her parents identified a long hair protruding from her mouth. She was comforted and given some food and drink. Afterwards they could no longer see the hair and she had no ongoing pain or issues initially. Over the subsequent days, she had increasing difficulty swallowing solid foods due to pain on swallowing and the parents later noticed a discolouration of the tip of her uvula. The family attended a walk in centre, the general practitioners and subsequently the children's emergency department 3 days after the initial incident. She was reviewed by the ear, nose and throat team who found that the distal tip of her uvula was white and that a hair could be seen trailing from it inferiorly. History and examination were otherwise unremarkable.

The child remained well and was taken to theatre for examination under general anaesthetic. On examination in theatre, the hair was found to be strangulating the distal uvula and trailing into the oesophagus ([figure 1](#)). It was not possible to untangle the hair, and as the tip was non-viable, the decision was taken to remove the uvula tip with bipolar cautery to reduce discomfort ([figure 2](#)).

Figure 1 Intraoperative image of strangulated uvula with hair visible.

Figure 2 Intraoperative image after bipolar resection of the distal uvula and hair removal.

The child was well postoperatively and discharged home the same day. Telephone follow-up with the parents at approximately 1 month post procedure confirmed no ongoing symptoms or complications.

Hair tourniquet syndrome (HTS) is a rare condition, more commonly seen affecting infants' external appendages, such as digits and genitalia.¹ Oral HTS has been described rarely, affecting sites,

Patient's perspective

My daughter uncharacteristically awoke one morning crying. Unable to comfort her, I identified a long hair protruding from her mouth, which caused her severe distress if touched. Unable to adequately examine her mouth, I assumed the hair was caught around a tooth. Fluids and a banana soothed her distress and allowed me to check her mouth where I could no longer see any sign of a hair. Over the following 2 days, she was well but mentioned discomfort when eating. I regularly tried to examine her throat but was not confident that the hair was wrapped around her uvula, partly as it was difficult to examine and her hair is blonde but also due to my assumption that such an event was highly improbable. However, when I started to witness a discolouration in the tip of her uvula, I sought medical assistance. Although the uvula was discoloured, a minor injuries unit dismissed the likelihood of Hair Tourniquet Syndrome, but following this up with our general practitioner led to a hospital admission and surgery. Following discharge, my daughter had no ill effects in either the short term or long term, awaking the following day without the need for pain relief or a change in diet. I assume her naturally long curly hair attributed to this incident.

Learning points

- ▶ Hair tourniquet syndrome (HTS) is a rare cause of strangulation of appendages.
- ▶ Swallowing difficulty with a history of a hair in the mouth should prompt consideration of oral HTS.
- ▶ Management of oral HTS will depend on the appendage involved and viability of the strangulated tissue. Under anaesthesia, options include either removal of the hair to allow for tissue reperfusion or autoamputation, or surgical debridement of the tissue if it is clearly non-viable.

© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Leighton J, Powell J, McLarnon C. *BMJ Case Rep* 2021;**14**:e242350. doi:10.1136/bcr-2021-242350

including the teeth, tongue and uvula.¹ A systematic literature review demonstrated only three published cases of uvular HTS in the past 40 years, all in infants.^{1–3} In these previously documented cases, the hair was removed and the uvula allowed to reperfuse or autoamputate. Manipulating a knotted hair in the oral cavity is challenging, and there are no documented cases of the hair being removed without sedation/general anaesthesia.

The uvula is a posterior extension of the soft palate which prevents nasal regurgitation on swallowing and resonates during speech.⁴ Uvular necrosis is most commonly iatrogenic, following intubation, endoscopy or suction.⁵ Most documented cases, predominantly in adults, resolve conservatively, with necrotic tissue sloughing off in 7–14 days.⁵

Contributors Patient seen by all three authors. JP took photographs. JL and JP wrote up. CM was the responsible clinician and supervised the write-up.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Jessica Leighton <http://orcid.org/0000-0002-6027-8026>

REFERENCES

- 1 Flores JR. Hair tourniquet syndrome in the dental patient. *Anesth Prog* 2014;61:111–2.
- 2 Krishna S, Paul RI. Hair tourniquet of the uvula. *J Emerg Med* 2003;24:325–6.
- 3 McNeal RM, Cruickshank JC. Strangulation of the uvula by hair wrapping. *Clin Pediatr* 1987;26:599–600.
- 4 Cascarini L, Schilling C, Gurney B, et al. *Oxford Handbook of oral and maxillofacial surgery*. Oxf. Handb. Oral Maxillofac. Surg. Oxford University Press.
- 5 Reid JW, Samy A, Jeremic G, et al. Postoperative Uvular necrosis: a case series and literature review. *Laryngoscope* 2020;130:880–5.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow