

A rare case of COVID-19 infection with laryngeal involvement

Masato Asaoka,¹ Shotaro Chubachi,¹ Yoshitake Yamada,² Koichi Fukunaga¹

¹Division of Pulmonary Medicine, Department of Medicine, Keio University School of Medicine, Shinjuku-ku, Tokyo, Japan

²Department of Radiology, Keio University School of Medicine, Shinjuku-ku, Tokyo, Japan

Correspondence to

Dr Shotaro Chubachi;
bachibachi472000@z6.keio.jp

Accepted 11 May 2021

DESCRIPTION

A 44-year-old man with obesity, but no other comorbidities, presented with a 3-day onset of fever and headache. On examination, he was not in respiratory distress, his body temperature was 38.0°C, body mass index was 35.8 kg/m² and oxyhaemoglobin saturation was 98% on room air. Novel COVID-19 was confirmed on nasopharyngeal swab PCR testing, and he was admitted to our hospital. On day 5 of admission (8 days after symptom onset), he experienced rapidly progressing acute hypoxemic respiratory failure and was treated with dexamethasone (6.6 mg intravenously) and remdesivir (200 mg loading dose intravenously on day 1, followed by 100 mg/day for 4 days). On day 7 of admission, he reported of new-onset hoarseness and difficulty breathing at rest. CT revealed bilateral peripheral/basal-predominant ground-glass opacities with linear opacities and laryngeal oedema (figure 1A,B). His symptoms were considered to be due to laryngeal oedema rather than pneumonia, and he was treated with a hydrocortisone drip (100 mg). On the next day, his symptoms became less severe, and dexamethasone (6.6 mg intravenously) administration was continued for 10 days. On day 14 of admission, all his symptoms were completely improved, and follow-up CT revealed an improvement of both lung involvement and laryngeal oedema (figure 1C,D). On day 19 of admission, the patient was discharged.




Figure 1 (A) Chest CT image on day 7 of admission showing bilateral peripheral/basal-predominant ground-glass opacities with linear opacities. (B) Neck CT images on day 7 of admission showing bilateral laryngeal oedema (arrows). (C) Chest CT images on day 14 of admission showing improvement in COVID-19 pneumonia. (D) Neck CT images on day 14 of admission showing improvement in laryngeal oedema (arrowheads).

Learning points

- ▶ laryngeal oedema may be a critical complication of COVID-19 infection, and CT can be an alternative to laryngeal fiberoscopy for the detection of laryngeal oedema.
- ▶ Systemic steroids may be effective against not only COVID-19 pneumonia but also the associated laryngeal oedema.

DISCUSSION

We report a rare case of COVID-19 infection with laryngeal involvement. Our main findings are as follows: first, laryngeal oedema may be a critical complication of COVID-19 infection, and CT can be an alternative to laryngeal fiberoscopy for the detection of laryngeal oedema. Second, systemic steroids may be effective against not only COVID-19 pneumonia but also the associated laryngeal oedema. Laryngeal oedema is an acute inflammation of the larynx that presents with difficulty breathing, hoarseness and stridor. Although most patients with COVID-19 experience upper airway symptoms, laryngeal oedema should be considered because it may cause sudden obstruction of the upper airway requiring rapid airway management. It is mainly caused by bacterial infection, injury from smoke and heat inhalation and as a complication of intubation.^{1 2} There have been several case reports of laryngeal oedema in COVID-19.³ However, all patients were diagnosed with postextubation laryngeal oedema. Our patient did not receive any intervention that could have caused laryngeal injury, such as intubation or suctioning. In addition, he had no smoking history or evidence of bacterial infection. Consequently, we considered that severe acute respiratory syndrome coronavirus 2 directly infected the larynx. As the mechanism of laryngeal involvement in COVID-19 is unclear, further studies are warranted.

In conclusion, we encountered a unique case of COVID-19 infection with laryngeal involvement. In this case, laryngeal involvement developed 7 days after hospitalisation. This case highlights that COVID-19 infection may cause life-threatening upper airway symptoms.

Contributors MA and SC attended to the patient, and MA and SC wrote the manuscript. YY and KF reviewed and approved the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.



© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Asaoka M, Chubachi S, Yamada Y, et al. *BMJ Case Rep* 2021;**14**:e242426. doi:10.1136/bcr-2021-242426

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

REFERENCES

- 1 Reid A, Ha JF. Inhalational injury and the larynx: a review. *Burns* 2019;45:1266–74.
- 2 Benjamin B. Prolonged intubation injuries of the larynx: endoscopic diagnosis, classification, and treatment. *Ann Otol Rhinol Laryngol* 2018;127:492–507.
- 3 Moran JV, Godil SA, Goldner B, *et al.* Post-Extubation stridor complicating COVID-19-Associated acute respiratory distress syndrome: a case series. *Cureus* 2020;12:e10492.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow