Straatsma syndrome: unilateral myelinated retinal nerve fibre layer, high myopia, strabismus and amblyopia

Mayank Jain ¹, Jalli Monica Sharon,¹ Rupa Anjanamurthy,¹
Hiruni Kaushalya Wijesinghe ²

¹Paediatric Ophthalmology,
Aravind Eye Hospital, Madurai,
Tamil Nadu, India

²Glaucoma Services, Aravind Eye
Hospital, Madurai, Tamil Nadu,
India

Correspondence to

Dr Mayank Jain;
sandymayank@gmail.com

Accepted 3 June 2021

Figure 2 (A) OCT of the right eye through the macula demonstrating hyper-reflective (arrowheads) retinal nerve fibre layer in the parafoveal area. (B) Left eye OCT showing normal retinal layers. OCT, optical coherence tomography; OD, right eye; OS, left eye.

DESCRIPTION

A 12-year-old boy presented with report of decreased vision in the right eye for the last year. On examination, his best-corrected visual acuity was 1.78 logMAR in the right eye and 0 logMAR in the left eye. Cycloplegic retinoscopy revealed a refractive error of a -15 dioptre sphere with a -3.5 dioptre cylinder at 180° in the right eye. The patient had right exotropia of 10 prism diopters on modified Krimsky's test with full extraocular motility. He had no stereopsis and demonstrated

Patient's perspective

I was deeply saddened by the fact that we came to know about my child's condition late enough to be treated effectively. Wearing contact lenses also did not help him much. Had I known about this disease earlier, my child would stand a better chance of regaining his eyesight. I hope there is a way to know about this earlier.

Figure 1 (A) Fundus photo of the right eye showing extensive myelination of the retinal nerve fibre layer. (B) Normal fundus of the left eye.

© BMJ Publishing Group
Limited 2021. No commercial
re-use. See rights and
permissions. Published by BMJ.

To cite: Jain M, Sharon JM,
Anjanamurthy R, et al. *BMJ
Case Rep* 2021;**14**:e244362.
doi:10.1136/bcr-2021-
244362

Learning points

- Myelination of retinal nerve fibre layer is a benign condition and does not affect vision, but may be associated with high myopia and strabismus.
- Early recognition of this condition in children by paediatricians and ophthalmologists can prevent dense amblyopia later in life.
- Treatment consists of full optical correction and initiation of antiamblyopia therapy.

right eye suppression on sensory examination. Dilated fundus examination was performed, which showed the presence of a myelinated retinal nerve fibre layer in the right eye along the superior and inferior temporal arcades covering the whole posterior pole with a dull foveal reflex (figure 1A). The left eye fundus was normal (figure 1B). The axial length of the right eye was 27.60 mm and that of the left eye was 22.07 mm. Optical coherence tomography was performed, which depicted parafoveal hyper-reflective retinal nerve fibre layer in the right eye and left eye was normal (figure 2). An optical correction was prescribed in the form of a contact lens in the right eye and occlusion therapy was given in the left eye. There was no improvement in the visual acuity after 6 months of amblyopia treatment and the patient was explained about the possible need for strabismus surgery in the future. Straatsma syndrome is defined as myelinated nerve fibre with myopia and amblyopia.¹ Myelination of

the optic nerve begins from the lamina cribrosa and the lamina cribrosa acts as a barrier at term. Defects in the lamina cribrosa or ectopic oligodendrocyte progenitor cells lead to the myelinated retinal nerve fibre layer.² Amblyopia, if detected early in life, can be treated with good visual outcomes.^{3 4}

Contributors MJ diagnosed the condition, investigated and advised treatment. JMS was involved in collection of images with valid consent. MJ and JMS were involved in drafting the manuscript of the case report. MJ, RA and HKW revised the manuscript and maintained the standards of the manuscript as per BMJ protocols. The final manuscript was approved by all the four authors. All authors are responsible for the integrity and accuracy of the data provided.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer-reviewed.

ORCID iDs

Mayank Jain <http://orcid.org/0000-0002-8314-4725>

Hiruni Kaushalya Wijesinghe <http://orcid.org/0000-0001-6234-8537>

REFERENCES

- 1 Straatsma BR, Heckenlively JR, Foos RY, *et al.* Myelinated retinal nerve fibers associated with ipsilateral myopia, amblyopia, and strabismus. *Am J Ophthalmol* 1979;88:506–10.
- 2 Ramkumar HL, Verma R, Ferreyra HA, *et al.* Myelinated retinal nerve fiber layer (RNFL): a comprehensive review. *Int Ophthalmol Clin* 2018;58:147–56.
- 3 Yalcin E, Balci O, Akingol Z. Association of extensive myelinated nerve fibers and high degree myopia: case report. *Indian J Ophthalmol* 2013;61:606–7.
- 4 Vide-Escada A, Prior Filipe H. Unusual Straatsma Syndrome - How dogmatic is a bad prognosis? *Am J Ophthalmol Case Rep* 2017;8:71–3.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow