Extremely viscous stool in a newborn leading to an early diagnosis of a life-long disease

Vilhelmiina Parikka ¹, Mari Fihlman,² Armatias Raitio ³

¹Pediatrics and Adolescent Medicine, Turku University Hospital, Turku, Finland

²Department of Anaesthesiology and Intensive Care, Turku University Hospital, Turku, Finland

³Department of Paediatric Surgery, Turku University Hospital, Turku, Finland

Correspondence to
Dr Vilhelmiina Parikka;
vilpar@utu.fi

Accepted 8 June 2021

DESCRIPTION

A female infant, born at 37+3 gestational weeks after an uncomplicated pregnancy, was referred to level III neonatal intensive care unit at the age of 24 hours due to a suspected bowel obstruction,

with a history of not having passed meconium and some non-bilious emesis. Physical examination showed considerable abdominal distension with visible bowel loops. Abdominal X-ray identified dilated bowel loops, with no evidence of perforation ([figure 1](#)).

Explorative laparotomy revealed significantly dilated loops of the ileum, with otherwise normal anatomy. The distal ileum was packed with extremely thick and sticky meconium, which could barely be removed through an enterotomy ([figure 2](#), [video 1](#)). The bowel was emptied by milking and irrigation with saline, after which an ileostomy was performed. Meconium ileus and viscous meconium were suggestive of cystic fibrosis. Laboratory tests revealed a low faecal elastase level (<15 µg/L) and a high chloride level (133 mmol/L) in a sweat test. Two

Figure 1 Long thorax X-ray showing distended bowel loops.

Figure 2 The distal ileum was packed with extremely thick meconium, which could barely be milked out through an enterotomy (see [video 1](#)).

Video 1 Thick meconium was difficult to milk out through an enterotomy.

© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Parikka V, Fihlman M, Raitio A. *BMJ Case Rep* 2021;**14**:e243828. doi:10.1136/bcr-2021-243828

heterozygous mutations, F508del and CFTRdele2,3 (21kb), were identified in genetic testing, confirming the diagnosis of cystic fibrosis at the age of 19 days. Early diagnosis enabled an early introduction of pancreatic enzymes, as well as enhanced nutritional support. The bowel recovered well, and the stoma was closed after 7 weeks.

Meconium ileus is manifested in one-fifth of the patients diagnosed with cystic fibrosis and is often the first manifestation of the disease.¹ Without screening, the median age at diagnosis can be more than 12 months.² Early diagnosis is

associated with a better prognosis.³ It is imperative that all infants with meconium ileus are tested for cystic fibrosis.

Twitter Arimatias Raitio @ArimatiasRaitio

Contributors VP drafted the manuscript; VP, MF and AR edited the text and images. All authors approved the final version of the manuscript to be published.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iDs

Vilhelmiina Parikka <http://orcid.org/0000-0002-9281-8154>

Arimatias Raitio <http://orcid.org/0000-0001-9114-2204>

REFERENCES

- 1 Sathe M, Houwen R. Meconium ileus in cystic fibrosis. *J Cyst Fibros* 2017;16 Suppl 2:S32–9.
- 2 Pedersen MG, Højte C, Olesen HV, *et al.* Late diagnosis and poor nutrition in cystic fibrosis diagnosed before implementation of newborn screening. *Acta Paediatr* 2019;108:2241–5.
- 3 Rosenfeld M, Sontag MK, Ren CL. Cystic fibrosis diagnosis and newborn screening. *Pediatr Clin North Am* 2016;63:599–615.

Learning points

- Infants with cystic fibrosis have exceptionally thick and sticky meconium, which can lead to bowel obstruction called meconium ileus.
- Most of the infants with meconium ileus have cystic fibrosis, making it imperative to test all infants with meconium ileus for cystic fibrosis.
- Early diagnosis of cystic fibrosis allows introduction of multidisciplinary care and instigation of life-long management and follow-up, and is associated with a better prognosis.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow