Hyphae in external auditory canal

Jeyasakthy Saniasiaya , Prepageran Narayanan

Department of
Otorhinolaryngology, Faculty
of Medicine, University of
Malaya, Kuala Lumpur, Wilayah
Persekutuan, Malaysia

Correspondence to
Dr Jeyasakthy Saniasiaya;
shakthy_18@yahoo.com

Accepted 16 July 2021

DESCRIPTION

A previously healthy 22-year-old man presented with a 1-week history of intermittent left aural pruritus. Besides that, there was no otalgia, otorrhoea, reduced hearing or vertigo. No history of allergy or recent trauma was noted. Rigid nasendoscopy and facial nerve examination were normal. Otoscopic examination revealed blackish-white spores occupying the left external auditory canal with an intact tympanic membrane (figure 1). Right otoscopic examination was unremarkable. Diagnosis of otomycosis was made based on clinical findings and by swab sent for culture and sensitivity which showed *Aspergillus niger*. Mechanical debridement was performed under microscopic guidance using suction, and the patient was prescribed antifungal ear drops, which demonstrated resolution of symptoms.

Otomycosis, also known as fungal otitis externa, is fungal infections involving the external auditory canal and occasionally middle ear with high prevalence noted among the tropical and humid subtropical regions. Approximately 9% of all otitis externa cases are contributed by otomycosis.¹ Overzealous usage of antibiotics has contributed tremendously to the recent surge in the number of otomycosis cases.² In the same vein, other contributing factors include frequent swimming, immunocompromised conditions such as diabetes mellitus, pregnancy, self-imposed injury of the ear canal by using cotton buds, post canal wall down mastoidectomy, and tympanic membrane perforation.³ The most common identified fungal pathogens include *Aspergillus niger* and *Candida albicans*. Otomycosis is diagnosed clinically from history and physical examination. Patients habitually complain of otalgia, pruritus, otorrhoea and even hearing loss. Otoscopic examination typically

Learning points

- ▶ Otomycosis although common should not be taken lightly as persistent symptoms may have detrimental effects such as hearing loss, tympanic membrane perforation, otomastoiditis and even osteomyelitis.
- ▶ Regular mechanical debridement under microscopic guidance with topical antifungal drops is deemed necessary.
- ▶ Otomycosis may be early sign of a more serious and debilitating condition such as diabetes mellitus.

presents with whitish, cottonlike or blackish greasy spores.

Treatment of otomycosis includes debridement along with antifungal medications. Persistent otomycosis leads to hearing loss, tympanic membrane perforation, fungal otomastoiditis, invasive temporal bone infection and meningoencephalitis.⁴ It is worth noting that tympanic membrane perforation is one of the overlooked complications of otomycosis. Fungal inoculation of the most medial aspect of the external auditory canal or direct extension of fungal infection into the tympanic membrane leads to mycotic thrombosis of the adjacent blood vessel leading to avascular thrombosis, hence tympanic membrane perforation.⁵

Contributors JS: writing, editing, literature review. PN: editing, literature review.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Jeyasakthy Saniasiaya <http://orcid.org/0000-0003-1974-4379>

REFERENCES

- 1 Ho T, Vrabec JT, Yoo D, et al. Otomycosis: clinical features and treatment implications. *Otolaryngol Head Neck Surg* 2006;135:787–91.
- 2 Munguia R, Daniel SJ. Ototoxicity of topical antifungals and otomycosis: a review. *Int J Pediatr Otorhinolaryngol* 2008;72:453–9.
- 3 Stren JC, Lucente FE. Otomycosis. *Ear Nose Throat J* 1998;77:804–10.
- 4 Rutt AL, Sataloff RT. Aspergillus otomycosis in an immunocompromised patient. *Ear Nose Throat J* 2008;87:622–3.
- 5 Hurst WB. Outcome of 22 cases of perforated tympanic membrane caused by otomycosis. *J Laryngol Otol* 2001;115:879–80.

Figure 1 Blackish-white hyphae occupying left external auditory canal.

© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Saniasiaya J, Narayanan P. *BMJ Case Rep* 2021;14:e245388. doi:10.1136/bcr-2021-245388

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow