Uncommon cause of respiratory distress—late-presenting congenital diaphragmatic hernia

Suvi Alikärri,¹ Vilhelmiina Parikka ,^{2,3} Mari Fihlman,⁴ Armatias Raitio ,^{3,5}

¹Department of Surgery, Porvoo Hospital, Porvoo, Finland

²Department of Paediatrics and Adolescent Medicine, Turku University Hospital, Turku, Finland

³Department of Clinical Medicine, University of Turku, Turku, Varsinais-Suomi, Finland

⁴Department of Anaesthesiology and Intensive Care, Turku University Hospital, Turku, Finland

⁵Department of Paediatric Surgery, Turku University Hospital, Turku, Finland

Correspondence to
Vilhelmiina Parikka;
vilpar@utu.fi

Accepted 22 August 2021

DESCRIPTION

A previously healthy 11-month-old boy was brought to the paediatric emergency unit due to breathing difficulties, tiredness and momentary loss of consciousness. His oxygen saturation was 95%. In a clinical examination, breathing sounds were not heard on the left side, and he was suffering from respiratory distress. Chest radiograph showed tracheal deviation, a mediastinal shift to the right and air-filled left thorax cavity (figure 1). These findings were reported as tension pneumothorax.

An emergency chest drain insertion in the operating theatre was scheduled. Initially, air was aspirated with a syringe, and a chest drain was inserted using the Seldinger technique,¹ resulting in a discharge of stomach contents from the drain. Esophagogastrosocopy confirmed the drain to be in the stomach. Intraoperative C-arm fluoroscopy revealed an intrathoracic location of the stomach consistent with a diagnosis of congenital diaphragmatic hernia. After discussing with parents, the diaphragmatic defect was repaired in laparotomy and the stomach perforation was sutured. The recovery was uneventful, and the boy was discharged on the fifth postoperative day. During 2 years of follow-up, no respiratory or gastrointestinal complications have been observed (figure 2).

Congenital diaphragmatic hernia is an uncommon congenital defect with an incidence of one in 2500–3000 live births.² However, 5%–20% of the cases are diagnosed after the neonatal period.³ Cases

Figure 2 Chest radiograph 2 years after congenital diaphragmatic hernia repair.

with late presentation may suffer from respiratory and/or gastrointestinal symptoms or be completely asymptomatic.³ In this case, the presence of several air-filled cavities, indistinct diaphragm on the left side, as well as inability to visualise the normal gastric bubble in chest radiograph were signs not typical to pneumothorax, and should have raised suspicion of congenital diaphragmatic hernia. Late-presenting congenital diaphragmatic hernia should be kept in mind as a rare diagnostic possibility when interpreting chest radiographs.

Learning points

- ▶ Congenital diaphragmatic hernia is an uncommon congenital defect, which is typically diagnosed prenatally.
- ▶ In rare cases, respiratory distress might be caused by a late-presenting congenital diaphragmatic hernia.
- ▶ Careful analysis of radiographic findings coupled with high clinical suspicion is needed to avoid misdiagnosis of late-presenting congenital diaphragmatic hernia.

Twitter Armatias Raitio @ArmatiasRaitio

Contributors SA drafted the manuscript; SA, VP, MF and AR edited the text and images. All authors approved the final version of the manuscript to be published.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Figure 1 Preoperative chest radiograph with mediastinal shift.

© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Alikärri S, Parikka V, Fihlman M, et al. *BMJ Case Rep* 2021;**14**:e245826. doi:10.1136/bcr-2021-245826

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iDs

Vilhelmiina Parikka <http://orcid.org/0000-0002-9281-8154>

Arimatias Raitio <http://orcid.org/0000-0001-9114-2204>

REFERENCES

- 1 McElnay PJ, Lim E. Modern techniques to insert chest drains. *Thorac Surg Clin* 2017;27:29–34.
- 2 Torfs CP, Curry CJ, Bateson TF, *et al.* A population-based study of congenital diaphragmatic hernia. *Teratology* 1992;46:555–65.
- 3 Baglaj M, Dorobisz U. Late-presenting congenital diaphragmatic hernia in children: a literature review. *Pediatr Radiol* 2005;35:478–88.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow