When a plain CT brain looks like contrast-enhanced- never forget polycythemia

Payel Biswas,¹ Bhagya Vardhan,² Subhankar Chatterjee ,² Umesh Kumar Ojha,² Souvik Dubey ³

¹Radiodiagnosis, Apollo Gleneagles Hospital, Kolkata, West Bengal, India

²General Medicine, Patliputra Medical College & Hospital, Dhanbad, Jharkhand, India

³Neuromedicine, Bangur Institute of Neurosciences, Institute of Postgraduate Medical Education & Research and SSKM Hospital, Kolkata, West Bengal, India

Correspondence to

Dr Souvik Dubey;
drsouvik79@gmail.com

Accepted 17 August 2021

DESCRIPTION

A 26-year-old man, suffering from complex congenital cyanotic heart disease (CCHD) with Eisenmenger complex, presented with seizures. Non-contrast CT (NCCT) scan of brain showed chronic infarct with encephalomalacic changes in the left high parietal region with vividly hyperdense intracranial arteries and venous sinuses (figure 1). Haemoglobin value was 253 g/L with 81% haematocrit.

Hyperdensity of the cerebral vasculature on NCCT results commonly from acute stroke ('dense artery sign' in acute arterial infarction), subarachnoid haemorrhage or cerebral venous sinus thrombosis (CVST). However, diffuse hyperdensity involving intracranial arteries simultaneously with dural venous sinuses on NCCT is reported in hemoconcentration and polycythemia.^{1,2} While polycythemia vera has been discussed to be associated with this striking finding, association of secondary polycythemia in an adult with complex CCHD is unique.¹⁻³

Flowing blood measures 40 HU on CT scan at 45% haematocrit. As the normal adult grey matter measures 39 HU and normal haematocrit is 42%–52%, normal cerebral vessels appear isodense or minimally hyperdense on NCCT.² A linear relationship is reported between haemoglobin level and the contrast of dural sinuses compared with cerebral grey matter, indicating that high haemoglobin level causes increased density of cerebral vessels on NCCT.¹ However, increased attenuation of venous sinuses is characteristically found in CVST. While attenuation values of more than 70 HU suggest

CVST, less than 70 HU favours polycythemia.⁴ In our case, CVST was ruled out on MR venography and the attenuation values ranged from 60 HU to

Patient's perspective

My son has been suffering from some congenital heart disease diagnosed at the age of 2 years, but could not be operated. He had multiple convulsions on the day of admission. Doctors ordered CT scan, which was abnormal. After performing further sophisticated imaging, the doctors assured us that there was no active bleed in the brain. They also informed us that the convulsions and abnormal CT imaging was due to excessive red cell mass secondary to his heart disease. My son is doing well after getting blood thinner (aspirin) and antiepileptics (given by patient's mother).

Learning points

- ▶ Seizures in congenital cyanotic heart disease can be due to hypoxic spells, cerebral venous sinus thrombosis (CVST), embolic infarct or abscess due to infective endocarditis and vascular congestion secondary to polycythemia.
- ▶ Polycythemia and CVST can mimic each other with respect to neurological symptoms clinically and radiologically on non-contrast CT brain. Moreover, CVST itself being a complication of polycythemia, these two must be differentiated to guide therapeutic decision-making.

Figure 1 Non-contrast CT images of the brain show diffusely hyperdense cerebral arteries, venous sinuses and cisterns (A to E). Gliotic foci are also noted in the high parietal lobe on the left side (F, G).

© BMJ Publishing Group Limited 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Biswas P, Vardhan B, Chatterjee S, et al. *BMJ Case Rep* 2021;**14**:e246106. doi:10.1136/bcr-2021-246106

Images in...

68 HU, correlating with hyperdense sinuses without thrombosis. The gliotic area in the left parietal region was probably due to old venous infarct. As polycythemia mimics CVST on NCCT brain, shares few similar neurological symptoms clinically and CVST being a complication of polycythemia, these two must be differentiated to guide further management.^{4 5}

Acknowledgements We are thankful to Dr Umesh Kumar Ojha, Professor and Head, Department of Medicine, Patliputra Medical College and Hospital, Dhanbad, India, for his constant support and encouragement.

Contributors PB, SC and SD generated the idea. PB did the literature search and wrote the first draft which was critically reviewed and edited by BV, SC, UKO and SD. All authors agreed upon the final version of the manuscript. All authors were involved in the management of patients and decision-making.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer-reviewed.

ORCID iDs

Subhankar Chatterjee <http://orcid.org/0000-0002-3555-4412>

Souvik Dubey <http://orcid.org/0000-0003-1733-3429>

REFERENCES

- 1 Ben Salem D, Osseby GV, Rezaizadeh-Bourdariat K, *et al.* [Spontaneous hyperdense intracranial vessels seen on CT scan in polycythemia cases]. *J Radiol* 2003;84:605–8.
- 2 Varwate P, Ilangoan G, Balganesan H, *et al.* Non-Enhanced CT Mimicking Contrast Enhanced CT - A Case Report on Polycythemia. *J Radiol Case Rep* 2019;13:10–16.
- 3 Gayathri S, Prasad A, Sachdeva N, *et al.* Unique features of polycythemia observed on plain non contrast CT scan of head. *J Pediatr Neurosci* 2010;5:27–9.
- 4 Black DF, Rad AE, Gray LA, *et al.* Cerebral venous sinus density on noncontrast CT correlates with hematocrit. *AJNR Am J Neuroradiol* 2011;32:1354–7.
- 5 Healy JF, Nichols C. Polycythemia mimicking venous sinus thrombosis. *AJNR Am J Neuroradiol* 2002;23:1402–3.

Copyright 2021 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow