

# Hypermobile Ehlers-Danlos syndrome: a video demonstration of Beighton score

Anusuya Sadhasivamohan,<sup>1</sup> Vijayasankar Palaniappan,<sup>2</sup> Kaliaperumal Karthikeyan<sup>3</sup>

<sup>1</sup>Dermatology, Venereology and Leprosy, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India

<sup>2</sup>Dermatology, Sri Manakula Vinayagar Medical College and Hospital, Puducherry, India

<sup>3</sup>Dermatology, Venereology & Leprosy, Sri Manakula Vinayagar Medical College and Hospital, Cuddalore, Tamil Nadu, India

## Correspondence to

Dr Vijayasankar Palaniappan; vijayasankarpalaniappan@gmail.com


Accepted 3 September 2022

## DESCRIPTION

An adolescent boy was brought by his parents with an 8-year history of increased elasticity of skin and flexibility of multiple joints. He also had associated bilateral knee joint pain for the last 6 months. The child's grandfather also had similar abnormal joint hypermobility. The medical history revealed a malunion of supracondylar fracture of the right humerus, which led to a gun-stock deformity. Cutaneous examination showed hyperelastic skin with a congenital melanocytic nevus over the right forearm. The joints were hypermobile with a Beighton score (BS) of 8 (figure 1, video 1). Hyperextension of the right elbow was not performed due to the gunstock deformity. There was no scoliosis or joint dislocation. Routine haematological and urine investigations did not reveal any abnormality. Neurological, cardiac and ophthalmological evaluation was found to be normal. A diagnosis of hypermobile Ehlers-Danlos Syndrome (hEDS) was established as the child satisfied the 2017 International Classification of the Ehlers-Danlos Syndromes criteria. Oral analgesics were prescribed for chronic knee joint pain along with knee stabilising braces. Health education regarding proper posture maintenance and mild strengthening exercise was provided to the patient.

EDS is a rare autosomal dominant connective tissue disorder affecting the skin and joints. There is a defect in the collagen metabolism that leads to the deposition of disordered collagen. Depending on the type of collagen affected, it has been classified

into various phenotypes. Hypermobile EDS is one of the subtypes characterised by generalised hypermobility of joints, recurrent joint dislocations, chronic joint pain and osteoarthritic changes. Skin is usually soft, mildly hyperextensible with easy



**Figure 1** (A–H) performance of various manoeuvres of Beighton scoring system in our patient.



**Video 1** A video demonstration of Beighton score in a hypermobile Ehlers-Danlos syndrome patient.



© BMJ Publishing Group Limited 2022. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Sadhasivamohan A, Palaniappan V, Karthikeyan K. *BMJ Case Rep* 2022;**15**:e252690. doi:10.1136/bcr-2022-252690

**Table 1** Beighton score for the assessment of generalised joint hypermobility

No.	Instructions given	Components	Left	Right
1	Keep your arms straight and wrist flexed, I will bend your thumb to the front of your forearm	Flexion of the thumb towards the flexor aspect of the forearm—passive	1	1
2	I shall extend your elbow backward	Hyperextension of elbow beyond 10°—passive	1	1
3	I shall extend your knee backward	Hyperextension of knee beyond 10°—passive	1	1
4	Keep your forearm on the flat surface, I will bend your little finger to the back of your hand	Dorsiflexion and hyperextension of the little finger beyond 90°—passive	1	1
5	Can you bend and touch the floor with your palms, keeping your knees straight	Forward flexion of the trunk with palms touching the floor, with knees straight—active	1	
Total score			9	

bruising. Patients may also have autonomic dysfunction with orthostatic intolerance and gastrointestinal symptoms.<sup>1</sup>

BS is an internationally recognised screening tool introduced in 1973 for the assessment of generalised joint hypermobility

### Learning points

- ▶ Beighton score is used to assess generalised joint hypermobility in heritable connective tissue disorders.
- ▶ A score of 5 and above out of 9 is considered generalised joint hypermobility.

(GJH). This nine-point scoring system requires the performance of five manoeuvres, which include four bilateral passive and one active performance (table 1). According to Beighton, the original cut-off value for GJH was  $\geq 5$ ,<sup>1-3</sup> However, since the mobility of joints decreases with age, the International Consortium on the Ehlers-Danlos Syndrome Committee proposed the following criterion of BS for GJH in hEDS as (1) BS  $\geq 6$  for adolescents and prepubertal children, (2)  $\geq 5$  for men and women from puberty to 50 years of age and (3)  $\geq 4$  above 50 years of age. The limitations of this scoring system are as follows: (1) Examination of only a few joints and (2) 'All or none' nature, that is, cannot assess the degree of joint hypermobility. This case reiterates the importance of the BS in the diagnosis of joint hypermobility disorders.<sup>1</sup>

**Contributors** AS: preparation of the manuscript. VP: preparation of the video. KK: overview of the manuscript and video.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Consent obtained from parent(s)/guardian(s).

**Provenance and peer review** Not commissioned; externally peer reviewed.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

### REFERENCES

- 1 Malfait F, Francomano C, Byers P, *et al*. The 2017 International classification of the Ehlers-Danlos syndromes. *Am J Med Genet C Semin Med Genet* 2017;175:8–26.
- 2 Malek S, Reinhold EJ, Pearce GS. The Beighton score as a measure of generalised joint hypermobility. *Rheumatol Int* 2021;41:1707–16.
- 3 Smits-Engelsman B, Klerks M, Kirby A. Beighton score: a valid measure for generalized hypermobility in children. *J Pediatr* 2011;158:119–23.

Copyright 2022 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/> BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

#### Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at [support@bmj.com](mailto:support@bmj.com).

Visit [casereports.bmj.com](http://casereports.bmj.com) for more articles like this and to become a Fellow