OPEN ACCESS

Abdominal heterotopic pregnancy post-IVF double embryo transfer

Chee Wai Ku,^{1,2} Isabella Ong ,³ Jerry Kok Yen Chan,^{1,2,3} Tat Xin Ee^{1,2}

¹Department of Reproductive Medicine, KK Women's and Children's Hospital, Singapore

²Duke-NUS Medical School, Singapore

³Yong Loo Lin School of Medicine, National University of Singapore, Singapore

Correspondence to
Dr Chee Wai Ku;
kucheewai@gmail.com

Accepted 16 December 2021

SUMMARY

Uterosacral ligament (USL) is an uncommon site of implantation for abdominal ectopic pregnancies. This is the first case of USL heterotopic pregnancy post invitro fertilisation (IVF). The patient presented 6 weeks after a double embryo transfer with acute onset abdominal pain and was diagnosed with a suspected live tubal ectopic pregnancy with a viable intrauterine pregnancy on ultrasound. A diagnostic laparoscopy revealed an ectopic pregnancy implanted on the left USL which was resected and confirmed on histology. The patient was discharged well on postoperative day 2 with a viable intrauterine pregnancy. This case highlights the importance of considering non-tubal heterotopic pregnancies in the context of risk factors including IVF with double embryo transfer presenting with abdominal pain.

BACKGROUND

Heterotopic pregnancies in the general population are rare, occurring 1 in 3889 to 1 in 30000 pregnancies.¹ The incidence increases among invitro fertilisation (IVF) patients, occurring approximately 1 in 100 to 1 in 1000 IVF pregnancies.^{2–4} There is a correlation between the number of embryos transferred and the risk of ectopic pregnancy.^{4–6} Fresh embryo transfer (ET) cycles have also been shown to increase the risk of heterotopic pregnancies post-IVF ET compared with frozen ET cycles.⁷ Other risk factors for heterotopic pregnancies include abnormal tubal anatomy due to previous tubal ectopic pregnancy, previous tubal surgery and endometriosis.⁸ The most common location for

heterotopic pregnancies was tubal, accounting for almost 90% of cases.⁹ 10.8%–11.8% of heterotopic pregnancies implant at non-tubal sites, and 1.3%–2.7% are abdominal.^{6 10 11}

In this case report, we present a case of abdominal heterotopic pregnancy implanted in the uterosacral ligament (USL) after double frozen ET.

CASE PRESENTATION

The patient is a primigravida in her 30s who presented 6 weeks post-ET with a 1 day history of sudden onset, sharp, colicky abdominal pain associated with nausea and non-bloody, non-bilious vomiting, and two episodes of diarrhoea. She had no vaginal bleeding and no contact with sick patients. On physical examination, she was alert, afebrile and had a heart rate of 90 beats per minute, but was hypotensive with blood pressure of 84/54 mm Hg. Her abdomen was soft with mild tenderness in the left iliac fossa with no rebound tenderness or guarding. On speculum examination, the cervical os was closed with no per vaginal bleeding or discharge.

She had a history of unexplained primary subfertility. The patient underwent hysterosalpingo-foam sonography which showed patency of both tubes, and normal endometrial cavity with no endometrial polyps or submucosal lesions. However, there were multiple intramural fibroids seen, ranging in size from 0.8 to 4.3×3.1×3.4 cm. She underwent an antagonist cycle with recombinant follicle-stimulating hormone 225 IU/L and gonadotropin-releasing hormone antagonist 0.25 mg. Eighteen eggs and eventually six blastocysts were obtained. Fresh transfer of one blastocyst was unsuccessful; hence, she went on to transfer two blastocysts on the frozen cycle. Two weeks later, her beta-human chorionic gonadotropin (β-hCG) level was 4682.3 mIU/mL. The review 4 weeks post-ET showed an intrauterine gestational sac (IUGS) and yolk sac with crown-rump length of 7 mm (corresponding to approximately 6 weeks' gestation) and positive foetal heart on ultrasound. There was no adnexal mass seen on the scan, and she was asymptomatic at that point in time.

INVESTIGATIONS

Bedside transvaginal ultrasound (TVUS) showed a viable intrauterine pregnancy (IUP) with a live ectopic seen in the left adnexa. The findings were suspicious for a heterotopic pregnancy. A second diagnostic TVUS performed in the radiology department confirmed the suspected diagnosis of heterotopic pregnancy with a live extrauterine pregnancy

Figure 1 Ultrasound scan showing a heterotopic pregnancy. (A) Transabdominal and (B) transvaginal ultrasound showing both intrauterine and ectopic pregnancy. (C) Left adnexal mass corresponding to likely tubal ectopic pregnancy. (D) Size of foetal pole corresponding to crown-rump length 1.8 cm. (E) Live ectopic pregnancy with positive foetal cardiac activity.

© BMJ Publishing Group Limited 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Ku CW, Ong I, Chan JKY, et al. *BMJ Case Rep* 2022;**15**:e246649. doi:10.1136/bcr-2021-246649

Figure 2 Resection of uterosacral ligament ectopic pregnancy. (A) Haemoperitoneum seen on entry into abdomen. (B) Normal right tube and ovary. (C) Normal left tube and ovary. (D–F): Mass with bleeding in pouch of Douglas. (G) Resected ectopic pregnancy from left uterosacral ligament.

corresponding to 8.3 weeks' gestation (figure 1). There was also a viable IUP seen corresponding to 8.9 weeks' gestation. There were also two uterine fibroids which measured 5.6×5.3×4.9 cm and 2.9 cm. There was no free fluid in the pouch of Douglas, but a heterogeneous mass was seen around the ectopic pregnancy suspected to be a blood clot.

The initial haemoglobin level was 95 g/L. A group and cross-match for two units of blood was also performed.

TREATMENT

The patient underwent a diagnostic four-port laparoscopy for a suspected left tubal ectopic pregnancy. Intraoperative findings showed a uterus of 10 weeks' size and multiple fibroids. Haemoperitoneum was noted on entry, suspicious of a ruptured ectopic pregnancy (figure 2A). However, both tubes were normal with no ectopic pregnancy seen, and ovaries were also normal bilaterally (figure 2B,C). Clots were evacuated and the pouch of Douglas was explored. Fresh bleeding was noted over the left USL with suspected products of conception (POC) (figure 2D–F). The ureter was identified and the POC were carefully dissected off the left USL (figure 2G). The specimen was retrieved in a bag and sent for histology. Interrupted sutures were applied, and topical haemostatic agents, including oxidised regenerated cellulose and human gelatine-thrombin matrix sealant, were applied over the left USL to secure haemostasis. An adhesion barrier was applied. The abdomen was then desufflated and the umbilical port closed with braided absorbable sutures. Two units of packed cells were transfused intraoperatively. The estimated blood loss was 2000 mL. A bedside transabdominal ultrasound done immediately postoperatively confirmed a viable IUP.

OUTCOME AND FOLLOW-UP

The patient remained haemodynamically stable in the ward with a haemoglobin level of 103 g/L on postoperative day 1 and was discharged well on the second postoperative day. Ultrasound for viability prior to discharge confirmed that the IUP was still viable. She was given a single dose of intramuscular progesterone 100 mg and kept on oral progestogen 10 mg two times per day. Histopathological findings of the resected specimen confirmed

POC, with vascularised chorionic villi seen. She was last seen in the clinic on 12 July 2021 with a viable IUP and a low-risk Down syndrome screening result.

DISCUSSION

Abdominal heterotopic pregnancies are rare occurrences with few cases reported. Here we present a case of a primigravida in her 30s with a history of unexplained subfertility who had an abdominal heterotopic pregnancy post-IVF double ET. She presented with acute onset abdominal pain and hypotension. A heterotopic tubal pregnancy was initially suspected on ultrasound imaging. The diagnostic laparoscopy revealed an abdominal heterotopic pregnancy implanted on the left USL, which was surgically resected. The patient was discharged well with a viable IUP.

Heterotopic pregnancies present an interesting diagnostic and therapeutic dilemma. Herein, we aim to present contemporary evidence-based management of heterotopic pregnancies, with particular focus on abdominal heterotopic pregnancies (figure 3).

Diagnosis

Diagnosis of a non-tubal heterotopic pregnancy can be challenging as both presentation and ultrasound findings are similar to tubal heterotopic pregnancies.¹² Patients with heterotopic pregnancies often present with abdominal pain and/or bleeding, and an adnexal mass.^{3 13–17} There were also asymptomatic cases detected only on ultrasound.^{18 19} A comprehensive history should be taken, including risk factors associated with heterotopic pregnancy, such as IVF with multiple ET, history of pelvic inflammatory disease, tubal surgery and endometriosis.⁸ Serum β -hCG has low utility due to the viable IUP in heterotopic pregnancies.²⁰ Accurate ultrasound diagnosis of non-tubal ectopic pregnancies is also challenging.¹² The Royal College of Obstetricians & Gynaecologists proposed an ultrasound criteria for abdominal ectopic pregnancies which includes: (1) absence of an IUGS, (2) absence of both an evidently dilated tube and a complex adnexal mass, (3) a gestational cavity surrounded by loops of bowel and separated by peritoneum and (4) a wide mobility similar to fluctuation of the sac particularly evident with pressure of the transvaginal probe towards the posterior cul-de-sac.^{21 22}

The index of suspicion for this patient must be elevated in view of the high β -hCG levels 2 weeks after ET. With such a high β -hCG level, double ET done and a single IUGS seen on scan, a conscious effort must be made to rule out a heterotopic pregnancy. Patients should be monitored for heterotopic pregnancies post-IVF ET with TVUS by trained sonographers as it has a high sensitivity for detection.^{2 23 24} The timing of the ultrasound scan is of great importance. This case was first seen at 6 weeks of gestation with only a single IUGS seen and subsequently presented 2 weeks later with heterotopic pregnancy diagnosed on ultrasound. Other cases reported also had initial ultrasound findings of a single IUGS at 5–7 weeks' gestation, and repeat ultrasound 1–2 weeks later revealed the ectopic pregnancy.^{3 18} At later gestation, the presence of an ectopic pregnancy may become more evident. Hence, it would be prudent to monitor patients with risk factors for heterotopic pregnancy by serial ultrasound scans.

If the diagnosis is still uncertain, and the patient is stable, other diagnostic imaging may be considered, such as MRI.²¹ MRI can be performed in stable patients to confirm the diagnosis and identify if the placenta has been implanted over

Figure 3 Approach to women with pain and bleeding in early pregnancy.

structures such as large vessels or bowel. It could also help to guide perioperative and operative considerations.²⁵ MRI of the pelvis has been used to precisely identify the location of the ectopic and to guide subsequent management.¹⁹

Management

Traditionally, management of the ectopic pregnancy would include conservative, medical or surgical methods. When managing heterotopic pregnancies, the IUP should be considered unless it is non-viable or undesired.²¹

Expectant management is rarely used except in certain stable patients, where the IUP is non-viable, or the ectopic is not implanted on essential structures such as major vessels.^{10 21} However, it is not routine management due to possible rupture of the ectopic as the abdominal pregnancy progresses. Five cases of heterotopic pregnancy managed conservatively have been reported. These cases were detected at an advanced gestational age and required hospitalisation for the remainder of the pregnancy with continuous monitoring and imaging at intervals. Hence, at advanced gestation, conservative management may be an option, with subsequent delivery of the IUP and abdominal ectopic at the same time.^{15 26–29}

Medical treatment with methotrexate should only be considered if the IUP is non-viable or undesired due to its teratogenic effects.^{30 31} However, it is also not routine management due to the high risk of treatment failure and potential adverse consequences of haemorrhage or rupture. Intracardiac potassium chloride or local hyperosmolar glucose injection with aspiration of the gestational sac is another possible intervention for stable patients. Four cases managed with intracardiac

potassium chloride injection were reported, of which three resulted in live births.¹⁹

Surgical resection of the ectopic pregnancy, via laparoscopy or laparotomy, remains the first-line treatment for both haemodynamically stable and unstable patients. Abdominal heterotopic pregnancy diagnosed at an early gestational age in stable patients should be managed laparoscopically. Laparotomy should be reserved for patients who are haemodynamically unstable and/or are at an advanced gestational age (more than 20 weeks' gestation).^{10 21 25} Seven cases diagnosed at 5–10 weeks' gestation were managed via laparotomy,^{14 16 17 32–34} and two diagnosed at 6–9 weeks' gestation were managed via laparoscopy.^{3 18}

Preoperatively, testing for haemoglobin and cross-matching for blood products are essential due to the propensity for blood loss during surgery.^{16 17 32} Fluid resuscitation and blood transfusions intraoperatively are usually required for resection of abdominal ectopic pregnancies.^{14 32} In this case, the ectopic was implanted on the USL. The principles of surgery remain largely similar to those involving the USL, such as deep infiltrating endometriosis surgeries. Identification of surrounding structures, such as the bowel, ureters, rectum, cervix and vagina, and major vessels would be essential for reduction of postoperative complications.^{35 36} In cases where there is a viable IUP, no intra-uterine devices should be used for manipulation, and laparoscopic manipulation of the uterus should be avoided.²¹ Complete resection of POC should be ensured. Haemostasis can be secured via sutures, electrocoagulation or use of haemostatic agents. The use of tranexamic acid should be considered.³⁷ Adhesion barriers can be applied to reduce abdominal adhesions.³⁸

Postoperatively, given that surgery may increase the risk of miscarriage, an ultrasound scan to monitor the viability of the IUP should be performed.¹⁸ Progesterone (intramuscular and/or oral) could also be given to support the pregnancy.^{18 33}

Follow-up

Follow-up for resolution can be challenging as monitoring of serum β -hCG levels is unreliable with a viable IUP.³ Thus, patients should be followed up with serial ultrasound scans, with close follow-up initially due to the increased risk of miscarriage.¹⁸ Spontaneous miscarriage after surgery usually occurs in the following 2–4 weeks.^{14 39} In a retrospective study of 64 heterotopic pregnancies, 14.1% of patients had a miscarriage, and it occurred within 3 weeks of the treatment. The only significant risk factor for miscarriage reported was early gestation at time of treatment, while the method of treatment (laparoscopic surgery, ultrasound-guided potassium chloride injection or gestational sac aspiration and observation) was not a significant risk factor.⁴⁰ In other studies examining the outcomes of heterotopic pregnancies managed laparoscopically, two patients had missed miscarriage within 8–14 days of the procedure, while two had miscarriages after rupture of membranes at 15–16 weeks' gestation.⁴¹

The timing and mode of delivery, after treatment for abdominal ectopic pregnancy, should be guided by obstetric indications. Among the cases of abdominal heterotopic pregnancy, some patients were delivered by caesarean section at 36–38 weeks' gestation due to patient's choice or undisclosed indications.^{33 34} Normal vaginal delivery at 40 weeks' gestation has also been reported.¹⁹ In two retrospective reviews of a total of 29 women who underwent laparoscopic surgery for heterotopic pregnancies, 23 women had live births. Across the two studies, 12 women delivered via normal vaginal delivery, while 13 had caesarean sections. Indications for caesarean section included failure to progress, cephalopelvic disproportion and previous caesarean section.^{41 42}

Learning points

- Clinicians should have a high index of suspicion for heterotopic pregnancies particularly in patients who have multiple risk factors such as having undergone in-vitro fertilisation with double embryo transfer, high beta-human chorionic gonadotropin levels, history of tubal disease or surgery, including pelvic inflammatory disease and endometriosis. While the location of the ectopic is most commonly tubal, clinicians should also consider other locations such as abdominal sites of implantation.
- The timing of the ultrasound scan is important as heterotopic pregnancies may be missed if done too early. Hence, even when a prior ultrasound scan showed a single intrauterine gestation sac, heterotopic pregnancies cannot be ruled out in patients with relevant risk factors, presenting with abdominal pain and per vaginal bleeding.
- Surgical management is recommended, via either laparoscopy or laparotomy for resection of products of conception. Do anticipate blood loss with resection of the abdominal ectopic pregnancies, with adequate blood products made available intraoperatively.
- Patients should be followed up closely with repeat ultrasound done for viability of the intrauterine pregnancy, given the increased risk of miscarriage in the next 2–4 weeks after treatment.

Contributors CWK, TXE and JKYC planned the paper. CWK and IO prepared the manuscript. CWK, IO, TXE and JKYC reviewed and edited the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s).

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

Case reports provide a valuable learning resource for the scientific community and can indicate areas of interest for future research. They should not be used in isolation to guide treatment choices or public health policy.

ORCID iD

Isabella Ong <http://orcid.org/0000-0002-3014-5193>

REFERENCES

- 1 Schroeppe TJ, Kothari SN. Heterotopic pregnancy: a rare cause of hemoperitoneum and the acute abdomen. *Arch Gynecol Obstet* 2006;274:138–40.
- 2 Li XH, Ouyang Y, Lu GX, XH L, GX L. Value of transvaginal sonography in diagnosing heterotopic pregnancy after in-vitro fertilization with embryo transfer. *Ultrasound Obstet Gynecol* 2013;41:563–9.
- 3 Knopman JM, Talebian S, Keegan DA, et al. Heterotopic abdominal pregnancy following two-blastocyst embryo transfer. *Fertil Steril* 2007;88:1437.e13–1437.e15.
- 4 Clayton HB, Schieve LA, Peterson HB, et al. Ectopic pregnancy risk with assisted reproductive technology procedures. *Obstet Gynecol* 2006;107:595–604.
- 5 Gardner DK, Lane M. Culture and selection of viable blastocysts: a feasible proposition for human IVF? *Hum Reprod Update* 1997;3:367–82.
- 6 Tal J, Haddad S, Gordon N, et al. Heterotopic pregnancy after ovulation induction and assisted reproductive technologies: a literature review from 1971 to 1993. *Fertil Steril* 1996;66:1–12.
- 7 Londra L, Moreau C, Strobino D, et al. Ectopic pregnancy after in vitro fertilization: differences between fresh and frozen-thawed cycles. *Fertil Steril* 2015;104:110–8.
- 8 Yoder N, Tal R, Martin JR. Abdominal ectopic pregnancy after in vitro fertilization and single embryo transfer: a case report and systematic review. *Reprod Biol Endocrinol* 2016;14:69.
- 9 Nabi U, Yousaf A, Ghaffar F, et al. Heterotopic Pregnancy - A Diagnostic Challenge. Six Case Reports and Literature Review. *Cureus* 2019;11:e6080.
- 10 Rojansky N, Schenker JG. Heterotopic pregnancy and assisted reproduction--an update. *J Assist Reprod Genet* 1996;13:594–601.
- 11 Bouyer J, Coste J, Fernandez H, et al. Sites of ectopic pregnancy: a 10 year population-based study of 1800 cases. *Hum Reprod* 2002;17:3224–30.
- 12 Lin EP, Bhatt S, Dogra VS. Diagnostic clues to ectopic pregnancy. *Radiographics* 2008;28:1661–71.
- 13 Reece EA, Petrie RH, Sirmans MF, et al. Combined intrauterine and extrauterine gestations: a review. *Am J Obstet Gynecol* 1983;146:323–30.
- 14 Wu Z, Zhang X, Xu P, et al. Clinical analysis of 50 patients with heterotopic pregnancy after ovulation induction or embryo transfer. *Eur J Med Res* 2018;23:17.
- 15 Bassil S, Pouly JL, Canis M, et al. Advanced heterotopic pregnancy after in-vitro fertilization and embryo transfer, with survival of both the babies and the mother. *Hum Reprod* 1991;6:1008–10.
- 16 Fisch B, Powsner E, Heller L, et al. Heterotopic abdominal pregnancy following in-vitro fertilization/embryo transfer presenting as massive lower gastrointestinal bleeding. *Hum Reprod* 1995;10:681–2.
- 17 Abdalla H, Ahuja KK, Morris N, et al. Combined intra-abdominal and intrauterine pregnancies after gamete intrafallopian transfer. *Lancet* 1986;2:1153–4.
- 18 Pisarska MD, Casson PR, Moise KJ, et al. Heterotopic abdominal pregnancy treated at laparoscopy. *Fertil Steril* 1998;70:159–60.
- 19 Yeh J, Aziz N, Chueh J. Nonsurgical management of heterotopic abdominal pregnancy. *Obstet Gynecol* 2013;121:489–95.
- 20 Marcus SF, Macnamee M, Brinsden P. Heterotopic pregnancies after in-vitro fertilization and embryo transfer. *Hum Reprod* 1995;10:1232–6.
- 21 Gynaecologists RCoA. Diagnosis and management of ectopic pregnancy: Green-top guideline No. 21. *BJOG* 2016;123:e15–55.
- 22 Gerli S, Rossetti D, Baiocchi G, et al. Early ultrasonographic diagnosis and laparoscopic treatment of abdominal pregnancy. *Eur J Obstet Gynecol Reprod Biol* 2004;113:103–5.
- 23 Cookingham LM, Goossen RP, Sparks AET, et al. Successful treatment algorithm for evaluation of early pregnancy after in vitro fertilization. *Fertil Steril* 2015;104:932–7.

- 24 Rein MS, Di Salvo DN, Friedman AJ. Heterotopic pregnancy associated with in vitro fertilization and embryo transfer: a possible role for routine vaginal ultrasound. *Fertil Steril* 1989;51:1057–8.
- 25 Worley KC, Hnat MD, Cunningham FG. Advanced extrauterine pregnancy: diagnostic and therapeutic challenges. *Am J Obstet Gynecol* 2008;198:297.e1–297.e7.
- 26 Dahniya MH, Shoukry IF, Balami WI, *et al.* Simultaneous advanced extrauterine and intrauterine pregnancy. *Int J Gynaecol Obstet* 1990;31:61–5.
- 27 Lenglet JE, Bekker MN, Akkerman C, *et al.* Prenatal ultrasound and MRI predict placental localization in a combined intrauterine and extrauterine twin pregnancy. *Prenat Diagn* 2006;26:376–8.
- 28 Shojai R, Chaumoitre K, Chau C, *et al.* Advanced combined abdominal and intrauterine pregnancy: a case report. *Fetal Diagn Ther* 2007;22:128–30.
- 29 Hathaway HR, Vasquez EC. Combined pregnancies with survival of both babies, report of a case. *Obstet Gynecol* 1961;18:352–4.
- 30 American College of Obstetricians and Gynecologists' Committee on Practice Bulletins—Gynecology. ACOG practice Bulletin No. 193: tubal ectopic pregnancy. *Obstet Gynecol* 2018;131:e91–103.
- 31 Practice Committee of American Society for Reproductive Medicine. Medical treatment of ectopic pregnancy: a Committee opinion. *Fertil Steril* 2013;100:638–44.
- 32 Dmowski WP, Rana N, Ding J, *et al.* Retroperitoneal subpancreatic ectopic pregnancy following in vitro fertilization in a patient with previous bilateral salpingectomy: how did it get there? *J Assist Reprod Genet* 2002;19:90–3.
- 33 Zhaoxia L, Honglang Q, Danqing C. Ruptured heterotopic pregnancy after assisted reproduction in a patient who underwent bilateral salpingectomy. *J Obstet Gynaecol* 2013;33:209–10.
- 34 Cormio G, Santamato S, Putignano G, *et al.* Concomitant abdominal and intrauterine pregnancy after in vitro fertilization in a woman with bilateral salpingectomy. A case report. *J Reprod Med* 2003;48:747–9.
- 35 Chapron C, Dubuisson JB. Laparoscopic treatment of deep endometriosis located on the uterosacral ligaments. *Hum Reprod* 1996;11:868–73.
- 36 Siddique SA, Gutman RE, Schön Ybarra MA, *et al.* Relationship of the uterosacral ligament to the sacral plexus and to the pudendal nerve. *Int Urogynecol J Pelvic Floor Dysfunct* 2006;17:642–5.
- 37 Gynecologists ACoOa. Topical hemostatic agents at time of obstetric and gynecologic surgery: ACOG Committee opinion summary, number 812. *Obstet Gynecol* 2020;136:870–1.
- 38 Huang C, Ding D-C. Outcomes of adhesion barriers in gynecologic surgeries: a retrospective study at a medical center. *Medicine* 2019;98:e18391.
- 39 Goldman GA, Fisch B, Ovadia J, *et al.* Heterotopic pregnancy after assisted reproductive technologies. *Obstet Gynecol Surv* 1992;47:217–21.
- 40 Na ED, Jung I, Choi DH, *et al.* The risk factors of miscarriage and obstetrical outcomes of intrauterine normal pregnancy following heterotopic pregnancy management. *Medicine* 2018;97:e12233.
- 41 Ping HL, Min ZH, Bi GJ, *et al.* Management and outcome of heterotopic pregnancy. *Ann Clin Lab Res* 2018;06.
- 42 Eom JM, Choi JS, Ko JH, *et al.* Surgical and obstetric outcomes of laparoscopic management for women with heterotopic pregnancy. *J Obstet Gynaecol Res* 2013;39:1580–6.

Copyright 2022 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <https://www.bmj.com/company/products-services/rights-and-licensing/permissions/>
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

Customer Service

If you have any further queries about your subscription, please contact our customer services team on +44 (0) 207111 1105 or via email at support@bmj.com.

Visit casereports.bmj.com for more articles like this and to become a Fellow